Doctopic: Experiments JID:PLB AID:137761 /SCO

Physics Letters B ••• (••••) ••••••

[m5G; v1.333] P.1 (1-7)



Please cite this article as: S. Diehl, A. Kim, K. Joo et al., A multidimensional study of the structure function ratio $\sigma_{LT'}/\sigma_0$ from hard exclusive π^+ electro-production off protons in the GPD regime, Physics Letters B, https://doi.org/10.1016/j.physletb.2023.137761

	Doctopic: Experiments		IPRESS	
	JID:PLB AID:137761 /SCO			[m5G; v1.333] P.2 (1-7)
	S. Diehl, A. Kim, K. Joo et al.			Physics Letters $B \bullet \bullet \bullet (\bullet \bullet \bullet \bullet) \bullet \bullet \bullet \bullet \bullet \bullet$
1	¹ Florida State University, Tallahassee, FL 32306, USA			
2	^m The George Washington University, Washington, DC 200	052, USA		
3	ⁿ GSI Helmholtzzentrum fur Schwerionenforschung GmbH	I, D-64291 Darmstadt, Germany		
4	^o INFN, Sezione di Ferrara, 44100 Ferrara, Italy			
4	^p INFN, Laboratori Nazionali di Frascati, 00044 Frascati, Ita	aly		
5	^q INFN, Sezione di Genova, 16146 Genova, Italy			
6	^r INFN, Sezione di Roma Tor Vergata, 00133 Rome, Italy			
7	^s INFN, Sezione di Torino, 10125 Torino, Italy			
8	' INFN, Sezione di Pavia, 27100 Pavia, Italy			
9	^a Universit e Paris-Saciay, CNRS/IN2P3, IJCLab, 91405 Orsa	ay, France		
10	W James Madison University Harrisonhurg VA 22807 USA	4		
14	^x Kyungnook National University Daegy 41566 Republic o	n of Koraa		
	J I amar University 4400 MIK Rhyd PO Box 10046 Regum	ont TX 77710 USA		
12	^z Massachusetts Institute of Technology. Cambridge, MA 02	2139-4307. USA		
13	^{aa} Mississippi State University. Mississippi State. MS 39762	2-5167. USA		
14	^{ab} National Research Centre Kurchatov Institute - ITEP, Mo.	oscow 117259, Russia		
15	^{ac} University of New Hampshire, Durham, NH 03824-3568	8, USA		
16	^{ad} New Mexico State University, PO Box 30001, Las Cruces,	, NM 88003, USA		
17	^{ae} Norfolk State University, Norfolk, VA 23504, USA			
17	^{af} Ohio University, Athens, OH 45701, USA			
18	^{ag} Old Dominion University, Norfolk, VA 23529, USA			
19	^{ah} II Physikalisches Institut der Universitaet Giessen, 35392	2 Giessen, Germany		
20	^{al} University of Richmond, Richmond, VA 23173, USA			
21	^a Universita' di Roma Tor Vergata, 00133 Rome, Italy			
22	^{ak} Skobeltsyn Institute of Nuclear Physics, Lomonosov Mos	scow State University, 119234 Moscow, R	lussia	
23	^{an} University of South Carolina, Columbia, SC 29208, USA			
24	an Thomas Jefferson National Accelerator Facility Newport	t News VA 23606 USA		
24	^{ao} Universidad Técnica Federico Santa María Casilla 110-V	V Valparaíso Chile		
25	^{ap} Università degli Studi dell'Insubria 22100 Como Italy	v, vaparaiso, enne		
26	^{aq} Universit'a degli Studi di Brescia, 25123 Brescia, Italy			
27	^{ar} University of California Riverside, 900 University Avenue	e, Riverside, CA 92521, USA		
28	as University of Glasgow, Glasgow, G12 800, United Kingdo	lom		
29	^{at} University of York, York YO10 5DD, United Kingdom			
30	^{au} University of Virginia, Charlottesville, VA 22901, USA			
31	av College of William and Mary, Williamsburg, VA 23187-8	8795, USA		
51	^{aw} Yerevan Physics Institute, 375036 Yerevan, Armenia			

ARTICLE INFO

ABSTRACT

Article history: Received 26 October 2022 Received in revised form 12 January 2023 Accepted 7 February 2023 Available online xxxx Editor: M. Doser Keywords: Exclusive single pion Eletroproduction GPD CLAS12

A multidimensional extraction of the structure function ratio $\sigma_{LT'}/\sigma_0$ from the hard exclusive $\vec{e}p \rightarrow \vec{e}p$ $e'n\pi^+$ reaction above the resonance region has been performed. The study was done based on beamspin asymmetry measurements using a 10.6 GeV incident electron beam on a liquid-hydrogen target and the CLAS12 spectrometer at Jefferson Lab. The measurements focus on the very forward regime (t/Q^2) \ll 1) with a wide kinematic range of x_B in the valence regime (0.17 < x_B < 0.55), and virtualities Q² ranging from 1.5 GeV^2 up to 6 GeV^2 . The results and their comparison to theoretical models based on Generalized Parton Distributions demonstrate the sensitivity to chiral-odd GPDs and the directly related tensor charge of the nucleon. In addition, the data is compared to an extension of a Regge formalism at high photon virtualities. It was found that the Regge model provides a better description at low Q^2 , while the GPD model is more appropriate at high Q^{2} .

© 2023 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/). Funded by SCOAP³.

Generalized Parton Distributions (GPDs) [1–3] provide direct access to the three-dimensional structure of the nucleon by cor-relating information about the transverse position and the longi-tudinal momentum of the quarks and gluons within the nucleon. Besides deeply virtual Compton scattering (DVCS), also deeply vir-tual meson production (DVMP) can be used to access GPDs. The factorization of the DVMP process into a perturbatively calculable hard-scattering part and two soft hadronic matrix elements, pa-rameterized by GPDs and a meson distribution amplitude (DA) as shown in Fig. 1 has been proven for longitudinally polarized virtual photons at large photon virtuality Q^2 , large energy W and fixed Bjorken-x [4,5]. The contribution of transversely polarized virtual photons for which factorisation is not explicitly proven, is typically treated as a higher twist-effect in current phenomenological mod-els [6].

Previous experimental results based on the hard exclusive scat-tering of 27.6 GeV/c un-polarized electron and positron beams off



Fig. 1. Hard exclusive electro-production of a pion on the proton in very forward kinematics $(-t/Q^2 \ll 1)$, described by GPDs [6,22].

Doctopic: Experiments JID:PLB AID:137761 /SCO S. Diehl, A. Kim, K. Joo et al.

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118 119

120

121

122

123

1 polarized and un-polarized hydrogen targets at HERMES [7–9], up 2 to 6 GeV/c polarized and un-polarized electron beams at CLAS 3 [10-17] and hall A at JLAB [18-20] and based on 160 GeV/c polar-4 ized muon beams at COMPASS [21], as well as theoretical studies 5 [6,22–25] of hard exclusive pseudoscalar meson electro-production, especially π^0 and *n* electroproduction [12–14,17,6,22,26,27], have 6 7 shown that the asymptotic leading-twist approximation is not suf-8 ficient to describe the experimental results from the existing meaq surements. It was found that there are strong contributions from 10 transversely polarized virtual photons that have to be considered 11 by including contributions from chiral-odd GPDs (H_T , \tilde{H}_T , E_T , and 12 \widetilde{E}_T) in addition to the chiral-even GPDs (*H*, \widetilde{H} , *E* and \widetilde{E}), which 13 depend on the momentum fraction of the parton x, the skewness 14 ξ and the four-momentum transfer to the nucleon *t*.

¹⁵ While chiral-even GPDs can be related to the well known nu-¹⁶ cleon form factors [28], only a few phenomenological constraints ¹⁷ exist for the chiral-odd GPDs. For example, the first moment of ¹⁸ $2\tilde{H}_T + E_T$ can be interpreted as the proton's transverse anomalous ¹⁹ magnetic moment [29], while in the forward limit, H_T becomes ²⁰ the transversity structure function h_1 , which is directly related to ²¹ the still unknown tensor charge of the nucleon [28]. ²² In exclusive meson production experiments GPDs are typically

In exclusive meson production experiments, GPDs are typically accessed through differential cross sections and beam and target polarization asymmetries [30–32]. The focus of this work is on the extraction of the structure function ratio $\sigma_{LT'}/\sigma_0$ from beam-spin asymmetry measurements. In the one-photon exchange approximation the beam-spin asymmetry is defined as [30,31]:

$$BSA = \frac{\sqrt{2\epsilon(1-\epsilon)}\frac{\sigma_{LT'}}{\sigma_0}\sin\phi}{1+\sqrt{2\epsilon(1+\epsilon)}\frac{\sigma_{LT}}{\sigma_0}\cos\phi + \epsilon\frac{\sigma_{TT}}{\sigma_0}\cos2\phi},\tag{1}$$

where the structure functions σ_L and σ_T , which contribute to $\sigma_0 = \sigma_T + \epsilon \sigma_L$, correspond to coupling to longitudinal and transverse virtual photons, and ϵ describes the flux ratio of longitudinally and transversely polarized virtual photons. σ_{LT} , σ_{TT} and the polarized structure function $\sigma_{LT'}$ describe the interference between their amplitudes. ϕ is the azimuthal angle between the electron scattering plane and the hadronic reaction plane.

 $\sigma_{LT'}$ can be expressed through the convolutions of GPDs with sub-process amplitudes (see Eq. (9)) and contains the products of chiral-odd and chiral-even terms [6]. For the π^+ channel, the imaginary parts of chiral-odd GPDs in $\sigma_{LT'}$ are significantly amplified by the pion pole term, where the contributions of GPDs are largely imaginary and those of the pion pole are real and can be accurately calculated. Due to this feature, polarized π^+ observables show an increased sensitivity to chiral-odd GPDs like H_T and can therefore be used to probe fundamental observables like the tensor charge δ_T for up (u) and down (d) quarks of the nucleon by

$$\delta_T^{u,d} = \int_{\xi-1}^1 dx H_T^{u,d}(x,\xi,t=0),$$
(2)

with the longitudinal momentum transfer ξ [26]. Due to the missing pion pole contribution, this sensitivity is much lower for exclusive π^0 and η production. In addition, π^+ observables are especially suited to access H_T , in contrast to π^0 and η production, due to the flavour composition of the charged pions.

An alternative description of hard exclusive pion production is based on Regge models. In these models, the interaction is mediated by the exchange of trajectories in the *t* channel. While Regge models were initially extensively studied for photoproduction $(Q^2 = 0)$ [33], an extension to the deeply virtual regime has been implemented within the Laget model (JML), which is based on Reggeized π^+ and ρ^+ meson exchanges in the *t*-channel

[34,35] and unitarity cuts [36,37]. The *t*-channel exchange of the 67 pion and the ρ rely on the canonical VGL [38] description, sup-68 plemented by the *t*-dependent electromagnetic form factor intro-69 duced in Ref. [39]. Alone these pole terms lead to a vanishing BSA. 70 The elastic $\pi - N$ [37] and inelastic $\rho - N$ unitarity cuts [36,37] 71 provide the phase necessary to get a non-zero BSA, through their 72 interference with the Regge poles. The JML model, which provides 73 74 a unified description at the real photon point, as well as in the 75 virtual photon sector, nicely reproduces the recent CLAS [40] and 76 HERMES [41] data on un-polarized π^+ electro-production cross 77 sections.

Altogether, two theoretical descriptions are available for hard exclusive π^+ electro-production. While the JML model starts at the real photon point and extends to the deeply virtual regime, a firm QCD foundation exists for the GPD model within the Bjorken regime and its applicability must be tested in the accessible Q^2 range.

Previous measurements of the hard exclusive π^+ production BSA (*i.e.* [42]) only provided a binning in -t and ϕ , while the virtuality Q^2 and the Bjorken scaling variable x_B where integrated over the complete accessible range due to limited statistics. In addition, only a limited range in Q^2 could be accessed due to the low electron beam energies that were available for these studies. For a precise comparison to theoretical models and especially for a study of higher-twist effects, a multidimensional study in t, ϕ , x_B and Q^2 with fine binning is needed to reduce uncertainties and to access the kinematic dependencies of the involved GPDs. In addition, a fully multidimensional study can provide a better comparison between the theoretical models and the data and help to investigate the validity of the two models.

For the present study, hard exclusive π^+ electro-production was measured at Jefferson Lab with CLAS12 (CEBAF Large Acceptance Spectrometer for operation at 12 GeV) [43]. Beam-spin asymmetries in forward kinematics were extracted over a wide range in Q^2 , x_B and ϕ . The incident electron beam was longitudinally polarized and had an energy of 10.6 GeV and an average current of 40-55 nA, impinging on a 5-cm-long un-polarized liquid-hydrogen target placed at the center of the solenoid magnet of CLAS12. The CLAS12 forward detector consists of six identical sectors within a toroidal magnetic field. The momentum and the charge of the particles were determined by 3 regions of drift chambers from the curvature of the particle trajectories in the magnetic field. The electron identification was based on a leadscintillator electromagnetic sampling calorimeter in combination with a Cherenkov counter. Positive pions were identified by timeof-flight measurements. Based on the high statistics of CLAS12, a precise, multidimensional study of the cross section ratio $\sigma_{LT'}/\sigma_0$ becomes possible for the first time.

For the selection of deeply inelastic scattered electrons, cuts on $Q^2 > 1.5 \text{ GeV}^2$, y < 0.75 and on the invariant mass of the hadronic final state W > 2 GeV, were applied. To select the exclusive $e'\pi^+n$ final state, events with exactly one electron and one π^+ were detected, and the missing neutron was selected via a cut on the neutron peak in the $e'\pi^+X$ missing mass spectrum. Fig. 2 shows the missing mass spectrum for $e'\pi^+X$ in the region around the missing neutron peak for selected bins of -t in the forward region, integrated over Q^2 and x_B .

124 As illustrated in Fig. 2, the signal-to-background ratio decreases with -t from \approx 4.5 at -t close to the threshold t_{min} to \approx 2 for 125 $-t \approx t_{min} + 1 \text{ GeV}^2$, making a background subtraction necessary for 126 beam-spin asymmetry extractions. The observed background be-127 haviour was found to be nearly independent of the Q^2 and x_B 128 bin. To determine the signal and background counts, the complete 129 distribution (signal + background) was fit with a Gaussian (de-130 131 scribing the signal) plus a third-order polynomial (describing the 132 background). After the combined fit, the signal and background

CLE IN PR

Doctopic: Experiments JID:PLB AID:137761 /SCO S. Diehl, A. Kim, K. Joo et al.

q



Fig. 2. Missing mass spectrum of $e'\pi^+X$ in the region of the missing neutron peak for selected bins of -t in the forward region. The raw distributions (upper histogram in each plot) were fit with a Gaussian (green curve) and a polynomial background (orange curve). For comparison, the background histogram obtained with the CERN-ROOT based background estimator applying a sensitive nonlinear iterative peak clipping algorithm [44] is shown in red and the background subtracted missing neutron peak is displayed as a black histogram fitted with a Gaussian (brown). The cut borders for the event selection are shown as vertical lines.



Fig. 3. Distribution of Q^2 versus x_B . The bin boundaries are shown as black lines and the bin numbering is given. The bin borders are also provided in the supplemental material [46].

contributions can be separated and integrated within a 2 σ region of the Gaussian distribution. As a crosscheck, another background histogram was obtained with the CERN-root based background estimator applying a sensitive nonlinear iterative peak clipping algorithm [44]. The obtained background was found to be very similar to the result from a full fit of the signal and background function (see Fig. 2), and was used to estimate the systematic uncertainty of the background subtraction.

Fig. 3 shows the Q^2 versus x_B distribution of the exclusive events, together with the binning scheme applied for the multidimensional study. For each of the nine $Q^2 - x_B$ bins, up to six bins in -t and 12 bins in ϕ were defined to extract the beam-spin asymmetry (BSA).

The BSA was determined experimentally from the number of counts with positive and negative helicity (N_i^{\pm}) , in a specific bin *i* as:

$$BSA_{i} = \frac{1}{P_{e}} \frac{N_{i}^{+} - N_{i}^{-}}{N_{i}^{+} + N_{i}^{-}},$$
(3)

where P_{e} is the average magnitude of the beam polarization. P_{e} was measured with a Møller polarimeter upstream of CLAS12 to



Physics Letters B ••• (••••) •••••

[m5G; v1.333] P.4 (1-7)

Fig. 4. Beam-spin asymmetry as a function of ϕ for representative -t bins of Q^2 – x_B bin 9 (Q² = 5.8 GeV², x_B = 0.55). The vertical error bars show the statistical uncertainty of each point. The red line shows the fit with the functional form of Eq. (1).

be 86.3%±2.6%. To obtain the signal counts, a full fit of the signal and background as described above was applied for each multidimensional bin in Q^2 , x_B , -t and ϕ and for each helicity state separately. The number of counts and their uncertainty were then given by the integral over the fit function of the signal distribution and the uncertainty of the beam-spin asymmetry was calculated based on standard error propagation.

To extract the structure function ratio $\sigma_{LT'}/\sigma_0$, the beam-spin asymmetry was plotted as a function of the azimuthal angle ϕ . Then a fit of the data with a $\sin \phi$ function was applied. The flux ratio ϵ (see Eq. (1)) was calculated for each bin based on the electron kinematics. Fig. 4 shows the beam-spin asymmetry as a function of ϕ in two different -t bins for the example of $Q^2 - x_B$ bin 9. Even in the highest Q^2 bin shown, a precise measurement of the ϕ dependence is possible. As expected, the ϕ -dependence can be well described by the assumed $\sin \phi$ shape. The impact of the denominator terms in Eq. (1) on $\sigma_{LT'}/\sigma_0$ was studied during the analysis using different extraction methods and was found to be on average 2.7% and, therefore, much smaller than the statistical and the total systematic uncertainty, and was considered as a systematic uncertainty.

The main source of systematic uncertainty is given by the back-ground subtraction. It was evaluated by comparing the two de-scribed background subtraction methods. The variation between the two methods which was in average 4.9% is considered as systematic uncertainty. The systematic effect due to the uncer-tainty of the beam polarization (3.4%) was determined based on the uncertainty of the measurement with the Møller polarimeter. To estimate the impact of acceptance and bin-migration effects, a realistic Monte Carlo simulation including all detector effects was performed. The impact of these effects was evaluated by compar-ing the injected and reconstructed asymmetries and was found to be in the order of 3.6%. Systematic uncertainties due to radiative effects have been studied based on Ref. [45], and were found to be in the order of 3.0%. Several additional sources of systematic un-certainty, including particle identification and the effect of fiducial volume definitions, were investigated and found to give a small contribution to the total systematic uncertainty (<1.5%). The total systematic uncertainty in each bin is defined as the square-root of

ARTICLE IN PRESS

Doctopic: Experiments JID:PLB AID:137761 /SCO S. Diehl, A. Kim, K. Joo et al.

q

[m5G; v1.333] P.5 (1-7) Physics Letters B ••• (••••) ••••••



Fig. 5. $\sigma_{LT'}/\sigma_0$ and its statistical uncertainty as a function of -t in the forward kinematic regime and its systematic uncertainty (grey bins). The bold dotted magenta line shows the theoretical prediction from the Regge based JML model [34]. The blue band shows the theoretical prediction from the GPD-based GK model [48,49]. The dashed brown and the dash-dotted red curve show the effect of increasing the GPD H_T by an overall factor of 1.5 and 2.0 for the mean kinematics. The dotted green curve shows the theory result under the assumption that no pion pole term is contributing. The corresponding result tables can be found in the supplemental material [46] and can be downloaded from Ref. [47].

the quadratic sum of the uncertainties from all sources. On average it was found to be on the order of 8.3%, which is smaller than the statistical uncertainty in most kinematic bins.

Fig. 5 shows the final results for $\sigma_{LT'}/\sigma_0$ in the region of -tup to 0.8 GeV² - 1.2 GeV², depending on the Q² bin $(-t/Q^2 \approx$ 0.2 - 0.4), where the leading-twist GPD framework is applicable and compares them to the theoretical predictions from the IML model [34], which is based on hadronic degrees of freedom and to the predictions from the GPD-based model by Goloskokov and Kroll (GK) [48,49]. The band on the theoretical prediction repre-sents the variation of the mean value of Q^2 and x_B within each multidimensional bin. The increasing width of these bands for bins 8 and 9, which cover a larger x_B and Q^2 range than the other bins, clearly shows the advantages of a fine multidimensional bin-ning for a precise theory comparison. The structure function ratio $\sigma_{LT'}/\sigma_0$ is clearly positive in all kinematic bins and shows a typical shape that can be explained by the contributing structure func-tions. The non- ϕ -dependent cross section $\sigma_0 = \sigma_T + \epsilon \sigma_L$ is typi-cally forward peaked due to the pion pole term contribution, while $\sigma_{LT'}$ is constrained to be zero at $t = t_{min}$ due to angular momen-tum conservation.

The GK model includes chiral-odd GPDs to calculate the contri-butions from the transversely polarized virtual photon amplitudes, with their *t*-dependence incorporated from Regge phenomenol-ogy. The GPDs are constructed from double distributions and con-strained by the latest results from lattice QCD and transversity parton distribution functions [48]. A special emphasis is given to the GPDs H_T and $\overline{E}_T = 2\widetilde{H}_T + E_T$, while contributions from other chiral-odd GPDs are neglected in the calculations, unlike chiral-even GPDs. The pion pole contribution to the amplitudes is taken into account for longitudinally and transversely polarized virtual photons.

 $\sigma_{LT'}$ can be expressed through the convolutions of GPDs with sub-process amplitudes (twist-2 for the longitudinal and twist-3 for the transverse amplitudes) and contains the products of chiral-odd and chiral-even terms [6]:

$$\sigma_{LT'} \sim \xi \sqrt{1 - \xi^2} \frac{\sqrt{-t'}}{2m} Im[\langle \overline{E}_{T-eff} \rangle^* \langle \widetilde{H}_{eff} \rangle + \langle H_{T-eff} \rangle^* \langle \widetilde{E}_{eff} \rangle], \qquad (4)$$

where *m* is the proton mass and the "eff" in the subscript describes the inclusion of the pion pole term, *i.e.*

$$\langle \widetilde{E}_{eff} \rangle = \langle \widetilde{E}_{\text{non-pole}} \rangle + c \frac{\rho_{\pi}}{t - m_{\pi}^2}$$
(5)

$$\langle \widetilde{H}_{eff} \rangle = \langle \widetilde{H} \rangle + \frac{\xi^2}{1 - \xi^2} \langle \widetilde{E}_{eff} \rangle \tag{6}$$

with a factor $c = m_p Q^2 / \xi$, the residue ρ_{π} and the pion mass m_{π} [48].

For π^+ the imaginary part of small chiral-odd GPDs in $\sigma_{LT'}$ is significantly amplified by the pion pole term, which is real and theoretically well described. The strength of this effect is illustrated in Fig. 5, which shows the comparison between the calculation with and without considering the pion pole (blue band vs green dotted line). Due to this feature, polarized π^+ observables show an increased sensitivity to chiral-odd GPDs in contrast to the exclusive π^0 and η production where the pole contribution is not present. The pion pole is well determined from cross section measurements with an uncertainty of less than 10%. Therefore, it 1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

ARTICLE IN PRESS

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121 122

123

124

125

126

127

128

129

130

131

132

cannot explain the observed overestimation of the experimental result by the theoretical prediction.

The denominator terms of the structure function ratio σ_L and σ_T can be expressed by [6]:

$$\sigma_{L} \sim (1 - \xi^{2}) \left| \langle \widetilde{H}_{eff} \rangle \right|^{2} - 2\xi^{2} Re \left[\langle \widetilde{H}_{eff} \rangle^{*} \langle \widetilde{E}_{eff} \rangle \right] - \frac{t'}{4m^{2}} \xi^{2} \left| \langle \widetilde{E}_{eff} \rangle \right|^{2}$$
(7)

$$\sigma_T \sim (1 - \xi^2) \left| \langle H_{T-eff} \rangle \right|^2 - \frac{t'}{8m^2} \left| \langle \overline{E}_{T-eff} \rangle \right|^2. \tag{8}$$

Due to the quark flavour composition of the pions, π^+ production is typically dominated by H_T , while the contribution from \overline{E}_T is significantly smaller. In contrast to this, neutral pseudoscalarmesons like π^0 and η show a significantly stronger contribution from \overline{E}_T , except at very small values of -t where H_T dominates. Since chiral even GPDs are much better known than their chiral odd counterparts, the strongest uncertainty for the theoretical prediction is expected from the so far poorly known GPD H_T for which the dependence on the measured structure function ratio is given in Eq. (9).

$$\sum_{\substack{23\\26\\26}}^{23} \frac{\sigma_{LT'}}{\sigma_0} \sim \frac{Im[\langle H_{T-eff} \rangle^* \langle \widetilde{E}_{eff} \rangle]]}{|\langle H_{T-eff} \rangle|^2 + \epsilon \sigma_L}.$$
(9)

27 The comparison between the experimental results and the the-28 oretical predictions shows that the magnitude of the GK model calculations is overestimated, while the t-dependence of the mea-29 30 sured $\sigma_{LT'}/\sigma_0$ values is, especially if the variation with Q^2 and x_B is considered, much better, but not perfectly reproduced. This 31 32 discrepancy of the magnitude might be due to the interplay of 33 the pion pole term with the poorly known chiral-odd GPD H_T . 34 Based on Eq. (9) the results especially hint on an underestimation 35 of H_T . To show the sensitivity of $\sigma_{LT'}/\sigma_0$ on the GPD H_T , Fig. 5 also contains calculations under the assumption that the GPD H_T 36 is increased by an overall factor of 1.5 (brown dashed line) and 37 by a factor of 2.0 (red dash-dotted line). Due to the amplification 38 39 by the pion pole term, a strong sensitivity to such a variation can 40 be observed. After the modification of the GPD H_T , a significantly 41 better agreement between the theoretical predictions and the ex-42 perimental result is observed.

However, a change of H_T will help as far as $\sigma_{LT'}/\sigma_0$ is con-43 cerned, but the consequences for other observables remain to be 44 45 checked. Especially observables with transversely polarized targets like the $\sin \phi_S$ modulation of the A_{UT} moment for hard exclusive 46 47 π^+ production, for which measurements based on HERMES data exist [48] and various modulations of A_{UT} and A_{LT} for ρ^0 pro-48 49 duction [50] show strong contributions from the transversity GPDs 50 and need to be considered for the determination of H_T . Altogether, 51 a new global fit of the GPDs to all existing data, e.g. [10,17,11–13], 52 as well as the aforementioned HERMES results and additional up-53 coming CLAS12 results on other mesons becomes necessary. Here, the new multidimensional, high precision π^+ beam-spin asymme-54 55 try data from this work and its high sensitivity to the GPD H_T 56 due to the amplification by the pion pole, will allow a much bet-57 ter determination of this so far poorly known GPD. Based on the 58 improvements in the knowledge of H_T , it will become possible to 59 extract the tensor charge of the proton, which is a fundamental 60 quantity and so far only poorly constrained.

The JML model, which turns out to reproduce available measurements of un-polarized electro-production cross-sections with a focus on Q^2 up to 5 GeV² and W up to 4 GeV [41,40], provides a reasonable description of the sign and the shape of $\sigma_{LT'}/\sigma_0$ at low and medium Q^2 and x_B values, but shows extrapolation problems for the highest Q^2 and x_B bins for which no explicit tuning could be performed based on previous data. The predicted theoretical $\sigma_{LT'}/\sigma_0$ values also fall short by a factor of two on average to reproduce the experimental values. However, a better agreement can be observed in the region of the lowest investigated Q^2 values, while the difference increases for higher values of Q^2 . The observed effects may originate from missing ingredients in the model. For instance, only the dominant singular unitary part of the re-scattering integrals is taken into account, while the effect of the principal part may be significant in the interference with the pole amplitudes. However, the observed difference in magnitude may also reflect the smallness of the theoretical transverse amplitude, which also misses the experimental value by a factor two at lower *W* [34].

The BSA measurement provides us with an access to the small box diagram contributions (either chiral odd GPD's or unitarity rescatterings), through their interference with the dominant pion pole amplitude. As Q^2 increases, the differences between the two theoretical approaches, as well as their departure from experiment, may tell us that they are used beyond their domain of validity: lack of other unitarity cuts in the Regge approach, energy not large enough to safely replace the hadronic basis by the quark basis in the GPD approach. The experiment presented here calls for improvements of the models along these lines.

In summary, we have performed a multidimensional measurement of the structure function ratio $\sigma_{LT'}/\sigma_0$ for $\vec{e}p \rightarrow e'n\pi^+$ at large photon virtuality, above the resonance region. The comparison in very forward kinematics showed that, especially, the magnitude of $\sigma_{LT'}/\sigma_0$ is overestimated in all Q^2 and x_B bins by the most advanced GPD-based model [48], indicating that a new global fit for the dominating GPD H_T becomes necessary to obtain a better fit for the dominant GPD H_T and the directly related tensor charge of the proton. Also the Regge-based JML model shows difficulties to fully reproduce the data and underestimates $\sigma_{LT'}/\sigma_0$ in the investigated Q^2 and x_B region. However, especially at low Q^2 , the JML model shows a slightly better agreement than the GK model, while the situation changes for high Q^2 where the GPD-based model provides a better reproduction of the data, especially after the GPD H_T is adjusted.

We acknowledge the outstanding efforts of the staff of the Accelerator and the Physics Divisions at Jefferson Lab in making this experiment possible. We owe much gratitude to P. Kroll for many fruitful discussions concerning the interpretation of our results. This work was supported in part by the U.S. Department of Energy, the National Science Foundation (NSF), the Italian Istituto Nazionale di Fisica Nucleare (INFN), the French Centre National de la Recherche Scientifique (CNRS), the French Commissariat pour l'Energie Atomique, the UK Science and Technology Facilities Council, the National Research Foundation (NRF) of Korea, the Helmholtz-Forschungsakademie Hessen für FAIR (HFHF), the Ministry of Science and Higher Education of the Russian Federation and the National Natural Science Foundation of China. The Southeastern Universities Research Association (SURA) operates the Thomas Jefferson National Accelerator Facility for the U.S. Department of Energy under Contract No. DE-AC05-060R23177.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

TICLE IN PRESS

Physics Letters B ••• (••••) •••••

Appendix A. Supplementary material

Supplementary material related to this article can be found online at https://doi.org/10.1016/j.physletb.2023.137761.

References

q

- [1] A.V. Radyushkin, Phys. Rev. D 56 (1997) 5524.
- [2] J.C. Collins, L. Frankfurt, M. Strikman, Phys. Rev. D 56 (1997) 2982.
- [3] S.J. Brodsky, et al., Phys. Rev. D 50 (1994) 3134.
- [4] A.V. Radyushkin, Phys. Lett. B 385 (1996) 333.
- [5] J. Collins, L. Frankfurt, M. Strikman, Phys. Rev. D 56 (1997) 2982.
- [6] S.V. Goloskokov, P. Kroll, Eur. Phys. J. A 47 (2011) 112.
- [7] A. Airapetian, et al., HERMES Collaboration, Phys. Lett. B 535 (2002) 85.
- [8] A. Airapetian, et al., HERMES Collaboration, Phys. Lett. B 659 (2008) 486.
- [9] A. Airapetian, et al., HERMES Collaboration, Phys. Lett. B 682 (2010) 345.
- [10] R. De Masi, et al., CLAS Collaboration, Phys. Rev. C 77 (2008) 042201.
- [11] I. Bedlinskiy, et al., CLAS Collaboration, Phys. Rev. Lett. 109 (2012) 112001.
- [12] I. Bedlinskiy, et al., CLAS Collaboration, Phys. Rev. C 90 (2014) 025205.
- [13] I. Bedlinskiy, et al., CLAS Collaboration, Phys. Rev. C 95 (2017) 035202.
- [14] A. Kim, et al., CLAS Collaboration, Phys. Lett. B 768 (2017) 168.
- [15] P. Bosted, et al., CLAS Collaboration, Phys. Rev. C 95 (2017) 035207.
- [16] P. Bosted, et al., CLAS Collaboration, Phys. Rev. C 95 (2017) 035206.
- [17] B. Zhao, et al., CLAS Collaboration, Phys. Lett. B 789 (2019) 426.
 - [18] E. Fuchey, et al., Hall A DVCS Collaboration, Phys. Rev. C 83 (2011) 025201.
- [19] M. Defurne, et al., Hall A DVCS Collaboration, Phys. Rev. Lett. 117 (2016)
- 262001. [20] M. Mazouz, et al., Hall A DVCS Collaboration, Phys. Rev. Lett. 118 (2017) 222002.
- [21] M.G. Alexeev, et al., COMPASS Collaboration, Phys. Lett. B 805 (2020) 135454.
- [22] G.R. Goldstein, J.O. Hernandez, S. Liuti, Phys. Rev. D 91 (11) (2015) 114013.
 - [23] M. Diehl, W. Kugler, Eur. Phys. J. C 52 (2007) 933.
- [24] G. Duplančić, D. Müller, K. Passek-Kumerički, Phys. Lett. B 771 (2017) 603.
- [25] M. Siddikov, I. Schmidt, Phys. Rev. D 99 (2019) 116005.

- [26] S. Ahmad, G.R. Goldstein, S. Liuti, Phys. Rev. D 79 (2009) 054014.
- [27] G.R. Goldstein, J.O. Hernandez, S. Liuti, Phys. Rev. D 84 (2011) 034007.
- [28] G.R. Goldstein, J.O.G. Hernandez, S. Liuti, Phys. Rev. D 91 (2015) 114013.
- [29] M. Burkardt, Phys. Rev. D 72 (2005) 094020;
- M. Burkardt, Phys. Lett. B 639 (2006) 462.
- [30] D. Drechsel, L. Tiator, J. Phys. G 18 (1992) 449.
- [31] T. Arens, O. Nachtmann, M. Diehl, P.V. Landshoff, Z. Phys. C 74 (1997) 651.
- [32] M. Diehl, S. Sapeta, Eur. Phys. J. C 41 (2005) 515.
- [33] M. Guidal, I.M. Laget, M. Vanderhaeghen, Nucl. Phys. A 627 (1997) 645.
- [34] J.M. Laget, Prog. Part. Nucl. Phys. 111 (2020) 103737, https://doi.org/10.1016/j.
- ppnp.2019.103737. [35] J.M. Laget, Phys. Rev. C 104 (2020) 025202, https://doi.org/10.1103/PhysRevC.
- 104 025202
- [36] J.M. Laget, Phys. Lett. B 685 (2010) 146, https://doi.org/10.1016/j.physletb.2010. 01.052
- [37] J.M. Laget, Phys. Lett. B 695 (2011) 199, https://doi.org/10.1016/j.physletb.2010.
- [38] M. Vanderhaeghen, M. Guidal, J.M. Laget, Phys. Rev. C 57 (1998) 1454.
- [39] J.M. Laget, Phys. Rev. D 70 (2004) 054023.
- [40] K. Park, et al., CLAS Collaboration, Eur. Phys. J. A 49 (2013) 16, https://doi.org/ 10.1140/epja/i2013-13016-9.
- [41] A. Airapetian, et al., HERMES Collaboration, Phys. Lett. B 6590 (2008) 486, https://doi.org/10.1016/j.physletb.2007.11.079.
- [42] S. Diehl, et al., CLAS Collaboration, Phys. Rev. Lett. 125 (2020) 182001, https:// doi.org/10.1103/PhysRevLett.125.182001.
- [43] V.D. Burkert, et al., CLAS Collaboration, Nucl. Instrum, Methods, Sect. A 959 (2020) 163419, https://doi.org/10.1016/j.nima.2020.163419.
- [44] M. Morhac, et al., Nucl. Instrum. Methods, Sect. A 401 (1997) 113.
- [45] A. Afanasev, I. Akushevich, V. Burkert, K. Joo, Phys. Rev. D 66 (2002) 074004.
- [46] See supplemental material for the result tables (link).
- [47] CLAS physics database, https://clas.sinp.msu.ru/cgi-bin/jlab/db.cgi.
- [48] S.V. Goloskokov, P. Kroll, Eur. Phys. J. C 65 (2010) 137.
- [49] B. Berthou, D. Binosi, N. Chouika, et al., Eur. Phys. J. C 78 (2018) 478, https:// doi.org/10.1140/epic/s10052-018-5948-0.
- [50] S.V. Goloskokov, P. Kroll, Eur. Phys. J. C 74 (2014) 2725.