## Physics Letters B 781 (2018) 719-722

Contents lists available at ScienceDirect

Physics Letters B

www.elsevier.com/locate/physletb

# Baryon masses and $\sigma$ terms in SU(3) BChPT $\times 1/N_c$

I.P. Fernando<sup>a</sup>, J.M. Alarcón<sup>b</sup>, J.L. Goity<sup>a,b,\*</sup>

<sup>a</sup> Department of Physics, Hampton University, Hampton, VA 23668, USA

<sup>b</sup> Thomas Jefferson National Accelerator Facility, Newport News, VA 23606, USA

### ARTICLE INFO

Article history: Received 9 April 2018 Received in revised form 20 April 2018 Accepted 25 April 2018 Available online 27 April 2018 Editor: B. Grinstein

*Keywords:* Sigma terms Nucleon mass Baryon masses Gell-Mann–Okubo mass formula

## ABSTRACT

Baryon masses and nucleon  $\sigma$  terms are studied with the effective theory that combines the chiral and  $1/N_c$  expansions for three flavors. In particular the connection between the deviation of the Gell-Mann–Okubo relation and the  $\sigma$  term associated with the scalar density  $\bar{u}u + \bar{d}d - 2\bar{s}s$  is emphasized. The latter is at lowest order related to a mass combination whose low value has given rise to a  $\sigma$  term puzzle. It is shown that while the nucleon  $\sigma$  terms have a well behaved low energy expansion, that mass combination is affected by large higher order corrections non-analytic in quark masses. Adding to the analysis lattice QCD baryon masses, it is found that  $\sigma_{\pi N} = 69(10)$  MeV and  $\sigma_s$  has natural magnitude within its relatively large uncertainty.

© 2018 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/). Funded by SCOAP<sup>3</sup>.

## 1. Introduction

Barvon mass dependencies on quark masses, quantified by the different  $\sigma$ -terms, are among the fundamental observables in baryon chiral dynamics. In particular, they give information on the baryon matrix elements of scalar quark densities, for which there is no alternative way for their determination. The definition of  $\sigma$  terms is through the Feynman–Hellmann theorem.<sup>1</sup> which, for three flavors, through the physical baryon masses gives access to only two such terms, namely those associated with the SU(3) octet quark mass combinations  $m_3 = m_u - m_d$  and  $m_8 = \frac{1}{\sqrt{3}}(\hat{m} - m_s)$ , where  $\hat{m}$  is the average of the *u* and *d* quark masses. The  $\sigma$  terms associated with the singlet component  $m_0 = \frac{1}{3}(2\hat{m} + m_s)$  require knowledge of baryon masses for unphysical quark masses, which is made possible through lattice QCD (LQCD) calculations. On the other hand, the pion-nucleon  $\sigma$  term  $\sigma_{\pi N} \equiv \frac{\hat{m}}{2m_N} \langle N | \bar{u}u + \bar{d}d | N \rangle$  is accessible through its connection to pion-nucleon scattering via a low energy theorem [1–3]. Such a determination of  $\sigma_{\pi N}$  had a long evolution through the availability of increasingly accurate data and the development of combined methods of dispersion theory and chiral perturbation theory [4–11]. The values obtained for  $\sigma_{\pi N}$  range from 45 MeV [4–6] to 64 MeV [7–12], where the difference between the results of the different dispersive analyses resides mostly in the different values of the S-wave  $\pi N$  scattering lengths  $a^{1/2,3/2}$  used in the subtractions, cf. [12]. In addition to the results from the analyses of  $\pi N$  scattering, LQCD calculations extrapolated to or at the physical point obtain different results, with values consistent with the recent  $\pi N$  results [13] and smaller,  $\sigma_{\pi N} \approx 40$  MeV [14–17]. The relatively large range of values obtained for  $\sigma_{\pi N}$  keeps it as an active topic of study, and in part motivates the present work. An additional motivation is the relevance of scalar quark operator matrix elements, quantities that are relevant in studies of direct dark matter detection [18–20], and of lepton flavor violation through  $\mu$ –e conversion in scattering with nuclei [21].

A puzzle that has been emphasized for a long time [22] is the relation between  $\sigma_{\pi N}$  in the isospin symmetry limit and the nucleon's  $\hat{\sigma} \equiv \sqrt{3} \frac{\hat{m}}{m_8} \sigma_8$ , namely  $\sigma_{\pi N} = \hat{\sigma} + 2 \frac{\hat{m}}{m_8} \sigma_s$ , which for a natural size value of  $\sigma_s$  should give  $\sigma_{\pi N} \sim \hat{\sigma}$ . The origin of the puzzle is the relation:  $\sigma_8 = \frac{1}{3}(2m_N - m_\Sigma - m_\Xi)$  (or other combinations related via the Gell-Mann–Okubo (GMO) relation) valid at linear order in quark masses, which gives  $\hat{\sigma} \sim 25$  MeV. If that relation is a reasonable approximation to the value of  $\hat{\sigma}$ , the implication is that, contrary to expectations,  $m_s$  must give a very large contribution to the nucleon mass even for the smaller values of  $\sigma_{\pi N}$ . The puzzle is particularly striking for the larger values that have been obtained for  $\sigma_{\pi N}$ , which would imply  $\sigma_s \sim 500$  MeV. Indeed, this is clearly impossible if one considers that  $\sigma_s$  is OZI suppressed with respect to  $\sigma_{\pi N}$ .

https://doi.org/10.1016/j.physletb.2018.04.054





<sup>\*</sup> Corresponding author.

E-mail address: goity@jlab.org (J.L. Goity).

<sup>&</sup>lt;sup>1</sup> The following notation will be used:  $\sigma_i(B) = m_i \frac{\partial}{\partial m_i} m_B$ , where  $m_i$  indicates a quark mass (i = u, d, s) or combination thereof (i = 0, 3, 8), and B is a given baryon. When B is not explicitly indicated it is assumed to be a nucleon.

<sup>0370-2693/© 2018</sup> The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/). Funded by SCOAP<sup>3</sup>.

This work analyzes the  $\sigma$  terms through the octet and decuplet baryon masses in the combined chiral and  $1/N_c$  expansions  $BChPT \times 1/N_c$ . The emphasis is in that the effective theory can give at NNLO (one chiral loop) a natural description of baryon masses, including LQCD results, along with the axial couplings which have been obtained in LOCD at different guark masses. In particular, the resolution of the  $\sigma$  term puzzle is explained by the fact that  $\Delta \sigma_8 \equiv \sigma_8 - \frac{1}{3}(2m_N - m_\Sigma - m_\Xi)$  receives large non-analytic in quark mass corrections dominated by  $m_s$ . It will also be shown that  $\sigma_8$  itself, and thus  $\hat{\sigma}$ , has a natural low energy expansion and therefore the origin of the puzzle resides in the large non-analytic correction to the mass combination  $\frac{1}{3}(2m_N - m_\Sigma - m_\Xi)$ . In fact, a big part of that large correction stems from the contribution of decuplet baryons in the loop, as it was found in Refs. [13,23]. By analyzing LQCD baryon masses [24], it is found that as expected  $\sigma_{\pi N} \sim \hat{\sigma}$ , with the results  $\sigma_{\pi N} = 69(8)(6)$  MeV, where the errors are respectively the statistical and theoretical (expected NNNLO corrections) ones, and  $|\sigma_s| \lesssim 50$  MeV. The connection between the deviation from the GMO relation,  $\Delta_{GMO} \equiv 3m_{\Delta} + m_{\Sigma} - 2(m_N + m_{\Xi})$ , and  $\Delta \sigma_8$ , both calculable at NNLO and given solely in terms of non-analytic loop contributions, is of particular importance in the present work.

#### 2. BChPT $\times 1/N_c$ analysis of masses and $\sigma$ terms

The combined BChPT ×  $1/N_c$  [25–29] implements the consistency of the effective theory with both the approximate chiral symmetry and the expansion in  $1/N_c$  of QCD. The expansion requires a link between the chiral and the  $1/N_c$  expansions: in practice the natural link is the  $\xi$  expansion where  $\mathcal{O}(p) = \mathcal{O}(1/N_c) = \mathcal{O}(\xi)$ , which is closely related to the so called small scale expansion [30,31] even when that one did not strictly implement the constraints of the  $1/N_c$  expansion. Consistency with  $1/N_c$  power counting demands the imposition of a dynamical SU(6) spin-flavor symmetry, which is broken by sub-leading corrections in  $1/N_c$  and requires the inclusion of the higher spin baryons (the decuplet in the case  $N_c = 3$ ) and relates low energy constants (LECs) in the chiral Lagrangian. The details on the calculations of baryon masses concerning the present work can be found in [29].

The chiral Lagrangian to  $O(\xi^3)$ , including electromagnetic corrections to the baryon masses is given by [29]:

$$\mathcal{L}_{B} = \mathbf{B}^{\dagger} \Big( i D_{0} + \mathring{g}_{A} u^{ia} G^{ia} - \frac{C_{HF}}{N_{c}} \hat{S}^{2} - \frac{1}{2\Lambda} c_{2} \hat{\chi}_{+} + \frac{c_{3}}{N_{c} \Lambda^{3}} \hat{\chi}_{+}^{2} + \frac{h_{1}}{N_{c}^{3}} \hat{S}^{4} + \frac{h_{2}}{N_{c}^{2} \Lambda} \hat{\chi}_{+}^{2} \hat{S}^{2} + \frac{h_{3}}{N_{c} \Lambda} \chi_{+}^{0} \hat{S}^{2} + \frac{h_{4}}{N_{c} \Lambda} \chi_{+}^{a} \{S^{i}, G^{ia}\} + \alpha \hat{Q} + \beta \hat{Q}^{2} \Big) \mathbf{B},$$
(1)

where terms not directly relevant to the baryon masses have been omitted. The spin-flavor singlet piece of the baryon masses,  $M_0 = O(N_c)$ , provides the large mass expansion parameter for HBChPT. In addition to the well known chiral building blocks, **B** represents the baryon spin-flavor multiplet field,  $\hat{S}^2$  is the square of the baryon spin operator,  $G^{ia}$  are the spin-flavor generators of SU(6), and  $\hat{Q}$  is the electric charge operator. No baryon-spin dependent electromagnetic effects are included. The term proportional to  $C_{HF}$ gives the leading order mass splitting between the spin 1/2 and 3/2 baryons.  $\mathring{g}_A$  is identified with  $\frac{6}{5}g_A^N$  at the LO, whose physical value is  $g_A^N = 1.2723(23)$ . The term  $h_1$  is only relevant if baryons with higher spin than 3/2 appear, which requires  $N_c \geq 5$ . The rest of the terms describe the quark mass effects. The combination  $\hat{\chi}_+ = N_c \ \chi_+^0 + \ \chi_+$ , where  $\chi_+^0 = \frac{1}{3}$ Tr  $\chi_+$  and  $\ \chi_+$  is the traceless piece of  $\chi_+$ , assures that the nucleon mass dependency on  $m_s$  is at most  $\mathcal{O}(N_c^0)$  (OZI).  $\Lambda$  is an arbitrary scale, which is conveniently chosen to be  $m_\rho$ . The baryon mass formula then reads (neglecting isospin breaking for now) [29]:

$$m_{B} = M_{0} + \frac{C_{HF}}{N_{c}} \hat{S}^{2} - \frac{c_{1}}{\Lambda} 2B_{0}(\sqrt{3}m_{8}Y + N_{c}m_{0}) - \frac{c_{2}}{\Lambda} 4B_{0}m_{0}$$
  
$$- \frac{c_{3}}{N_{c}\Lambda^{3}} \left( 4B_{0}(\sqrt{3}m_{8}Y + N_{c}m_{0}) \right)^{2}$$
  
$$- \frac{h_{1}}{N_{c}^{2}\Lambda} \hat{S}^{4} - \frac{h_{2}}{N_{c}\Lambda} 4B_{0}(\sqrt{3}m_{8}Y + N_{c}m_{0}) \hat{S}^{2} - \frac{h_{3}}{N_{c}\Lambda} 4B_{0}m_{0} \hat{S}^{2}$$
  
$$- \frac{h_{4}}{N_{c}\Lambda} \frac{4B_{0}m_{8}}{\sqrt{3}} \left( 3\hat{I}^{2} - \hat{S}^{2} - \frac{1}{12}N_{c}(N_{c} + 6) \right)$$
  
$$+ \frac{1}{2}(N_{c} + 3)Y - \frac{3}{4}Y^{2} + \delta m_{B}^{loop}, \qquad (2)$$

where  $\delta m_B^{\text{loop}}$  can be obtained with some work using the results in [29], where the details on the mass renormalization and results for general  $N_c$  can be found.

Setting  $c_3 = 0$ ,<sup>2</sup> the terms analytic in quark masses in Eqn. (2) lead to the exact GMO and Equal Spacing mass relations, which are unchanged at generic  $N_c$ . On the other hand at generic  $N_c$  the mass relation for  $\sigma_8$  at tree level reads:

$$\Delta \sigma_8 = \sigma_8 - \frac{1}{9} \left( \frac{5N_c - 3}{2} m_N - (2N_c - 3)m_\Sigma - \frac{N_c + 3}{2} m_\Xi \right).$$
(3)

The dominant contributions to  $\Delta_{GM0}$  and  $\Delta\sigma_8$  are calculable nonanalytic contributions.  $\Delta_{GM0}$  is  $\mathcal{O}(\xi^4)$  and in large  $N_c$  limit it is  $\mathcal{O}(1/N_c)$ . On the other hand,  $\sigma_8$  is  $\mathcal{O}(\xi)$  and it has a prefactor  $N_c$ , and  $\Delta\sigma_8$  is  $\mathcal{O}(\xi^2)$  also with a prefactor  $N_c$ .  $c_3$  gives a contribution to the  $\Delta_{GM0}$  which is  $\mathcal{O}(\xi^5)$ , and to  $\Delta\sigma_8$  at  $\mathcal{O}(\xi^4)$ , both being beyond the accuracy of the present work.  $\Delta_{GM0}^3$  and  $\Delta\sigma_8$ are thus determined by the meson masses and by the LECs  $\mathring{g}_A/F_{\pi}$ , and  $C_{HF}$ .  $\Delta_{GM0}$  depends rather smoothly on  $C_{HF}$ , and drives to a large extent the determination of  $\mathring{g}_A/F_{\pi}$ . One finds the interesting fact that the ratio  $\Delta\sigma_8/\Delta_{GM0}$ , which is independent of  $\mathring{g}_A/F_{\pi}$ , is also almost entirely independent of the value of  $C_{HF}$  in a very wide range around its actual value. For  $N_c = 3$ ,  $\sigma_8/\Delta_{GM0} \sim -13.5$ , which translates into  $\Delta\hat{\sigma}/\Delta_{GM0} \sim 1.68$ .

The analysis of the physical octet and decuplet baryon masses suffice to make the main point of this work. In this case, the LECs  $c_2$ ,  $c_3$  and  $h_1$  are set to vanish, because at the order of the calculation they are redundant. A fit is carried out including strong and electromagnetic isospin breaking. This requires using the meson masses with isospin breaking, which include  $\eta - \pi^0$  mixing (required to have a consistent renormalization of the baryon masses) and the electromagnetic mass shifts where Dashen's theorem is used, which should be sufficient for the current application. The electromagnetic addition to  $\Delta_{\text{GMO}}$  is equal to  $-\frac{4}{3}\beta$ , while the strong isospin breaking has negligible effect, and the electromagnetic contribution to the p-n mass difference is equal to  $\alpha + \beta$ . The result of the fit to physical masses is shown in Table 1, Fit 1.

The information given by LQCD, where the baryon masses have been obtained with  $M_K$  approximately constant and varying  $m_u = m_d$  in a range where 213 MeV  $< M_\pi < 430$  MeV [24], is very

<sup>&</sup>lt;sup>2</sup> The 27-plet SU(3) breaking produced by this term is  $O(\xi^5)$ , and thus for the current purposes it can be neglected.

 $<sup>^3</sup>$   $\Delta_{GMO}$  corresponds to having removed the EM corrections, otherwise it is denoted by  $\Delta_{GMO}^{phys}.$ 

Table 1

Results from fits to baryon masses. Fit 1 uses only the physical octet and decuplet masses, Fit 2 uses the physical and the LQCD masses from Ref. [24] with  $M_{\pi} \lesssim 300$  MeV, and Fit 3 uses only those LQCD masses and imposes the value of  $\Delta_{GMO}^{\text{phys}}$  determined by the physical masses. The renormalization scale  $\mu$  and the scale  $\Lambda$  are taken to be equal to  $m_{\rho}$ . \* indicates an input. An estimated theoretical error of 6 MeV is indicated for  $\hat{\sigma}$  and  $\sigma_{\pi N}$ .

Fit	$rac{\dot{g}_A}{F_\pi}$ MeV $^{-1}$	$rac{M_0}{N_c}$ MeV	C <sub>HF</sub> MeV	<i>c</i> <sub>1</sub>	C2	h <sub>2</sub>	h <sub>3</sub>	h4	α MeV	eta MeV
1	0.0126(2)	364(1)	166(23)	-1.48(4)	0	0	0.67(9)	0.56(2)	-1.63(24)	2.16(22)
2	0.0126(3)	213(1)	179(20)	-1.49(4)	-1.02(5)	-0.018(20)	0.69(7)	0.56(2)	-1.62(24)	2.14(22)
3	0.0126*	262(30)	147(52)	-1.55(3)	-0.67(8)	0	0.64(3)	0.63(3)	$-1.63^{*}$	2.14*
	nhvs				•					
Fit	$\Delta_{GMO}^{phys}$	$\sigma_8$		$\Delta \sigma_8$	σ	$\sigma_{\pi N}$	$\sigma_s$		$\sigma_3$	$\sigma_{u+d}(p-n)$
	MeV	MeV		MeV	MeV	MeV	MeV		MeV	MeV
1	25.6(1.1)	-583(24)		-382(13)	70(3)(6)	-	-		-1.0(3)	-1.6(6)
2	25.5(1.5)	-582(55)		-381(20)	70(7)(6)	69(8)(6)	-3(32)		-1.0(4)	-1.6(8)
3	25.8*	-615(80)		-384(2)	74(1)(6)	65(15)(6)	-121(15)	5)	-	-



**Fig. 1.** Left panel: summary of the determinations of  $\sigma_{\pi N}$  from  $\pi N$  scattering (blue), from LQCD (red), and from this work showing the combined fit and theoretical error. Right panel: N and  $\Delta$  masses from Fit 2 of Table 1: physical and LQCD masses from [32]. The squares are the results from the fit and the error bands correspond to 68% confidence interval. (For interpretation of the colors in the figure(s), the reader is referred to the web version of this article.)

useful for testing the effective theory, and necessary for calculating  $\sigma_{\pi N}$ . Two different fits that include LQCD baryon masses were performed, shown in Table 1. One fit (Fit 2) combines the physical and LQCD masses, up to  $M_{\pi} \sim 300$  MeV, and the other (Fit 3) uses only LQCD and the physical value of  $\Delta_{GMO}$ , which is important for controlling the value of  $\mathring{g}_A/F_{\pi}$ . In these fits the LEC  $c_2$  which gives the baryon mass dependencies on the singlet quark mass component  $m_0$  becomes significant, and its presence is responsible for the significant change in  $M_0$  compared to the physical fit.  $M_0$  is very precisely determined by the physical masses; Fit 3 shows that it is much less precise if only LQCD masses are used. The constant  $\beta$  can be estimated by the relation  $2\beta = m_p - m_n - (m_{\Xi^0} - m_{\Xi^-})$ , valid to LO in quark masses, which gives  $\beta = 2.78 \pm 0.1$  MeV. The fit indicates that higher order terms in quark masses affect the extraction of  $\beta$ . The theoretical error for  $\hat{\sigma}$  and  $\sigma_{\pi N}$  accounting for higher order corrections was estimated by explicitly expanding in  $\xi$  and identifying the size of the contributions; the magnitude of the theoretical error was then estimated to be  $\sim 1/3$  the size of the last term in the expansion.

The observations derived from the effective theory and from the fits are the following:

- i) The value of  $\mathring{g}_A/F_{\pi}$  is to a large extent fixed by  $\Delta_{GMO}$ , and it corresponds to a value of  $g_A^N$  at LO which is roughly a factor 0.75 of the physical one; this agrees with what is observed in the analysis of the axial vector couplings [29] provided by LQCD calculations at different values of quark masses [32].
- ii) The octet baryons contribute 43% of  $\Delta_{GM0}$ , and 33% of  $\Delta\sigma_8$ , which shows the importance of the decuplet contributions.
- iii) The first fit determines  $\sigma_8$ . Using the natural renormalization scale  $\mu = m_\rho$ , the different contributions to  $\sigma_8$  are primarily

given by the terms  $c_1$  (~ -870 MeV),  $h_4$  (~ 110 MeV) and the loop contributions (~ 190 MeV), where the latter two are the NLO contributions. This seems to be a well behaved expansion. On the other hand the mass combination on the RHS of Eqn. (3) has the corresponding pattern -870 MeV, 110 MeV and 570 MeV, the latter loop contribution given by the additional contribution due to  $\Delta \sigma_8 \sim -380$  MeV. The NLO terms in the mass combination are very large and reduce the LO one.

- iv) The correction  $\Delta\sigma_8$  becomes quite large for  $M_K > 350$  MeV, being about 70% of  $\sigma_8$  for the physical  $M_K$ . As mentioned earlier,  $\Delta\sigma_8$  and  $\Delta_{GMO}$  are determined only in terms of  $\mathring{g}_A/F_{\pi}$ ,  $C_{HF}$  and the meson masses. The ratio  $\Delta\sigma_8/\Delta_{GMO}$  does not depend on  $\mathring{g}_A/F_{\pi}$ , and has virtually no dependence on  $C_{HF}$ . The ratio is also modestly dependent on  $M_K$ , going from  $\sim -11$  to  $\sim -14$  when  $M_K$  is increased from 200 to 600 MeV.
- v) The combined fit of physical and LQCD masses, Fit 2, is compatible with Fit 1; this implies that the chiral extrapolation of the LQCD results to the physical case is consistent.
- vi) The fit to only LQCD masses and imposing the physical  $\Delta_{GMO}$ , Fit 3, serves for a consistency check, which turns out to be quite reasonable. The LQCD masses do not describe correctly the hyperfine mass shifts between the octet and decuplet, which is shown in Fig. 1 right panel, where the  $\Delta$  mass is systematically large, and this is the reason the resulting  $C_{HF}$ has some difference with the other fits. The extrapolation to the physical case turns out to be from 20 to 50 MeV larger than the physical octet masses, but less accurate for the decuplet ones where the  $\Delta$  mass, which is the worst case, comes out to be about 100 MeV larger than the physical one.

- vii) It is observed that  $\hat{\sigma}$  and  $\sigma_{\pi N}$  have both a small and approximately linear dependency on  $M_K$  in a very wide range. This in particular indicates that  $\hat{m} \sigma_s/m_s$  must remain relatively small throughout.
- viii)  $\sigma_s$  is poorly determined in the present study because the LQCD results are at approximately fixed  $m_s$ . Its range of values is however in line with the natural expectations. A LQCD calculation performed with smaller  $M_K$  than the physical one is necessary to obtain  $\sigma_s$  with better precision and also for understanding the effective theory in general.
- ix) The results obtained for  $\sigma_{\pi N}$  are consistent with the larger values obtained from  $\pi N$  analyses [7–11]. Note however that a more reliable value would require some more accurate and extensive LQCD results. Fig. 1 depicts the result for  $\sigma_{\pi N}$  from Fit 2 and its comparison with other results.
- x) The analysis also gives an estimate of the isospin-breaking  $\sigma$  terms  $\sigma_3$  and  $\sigma_{u+d}(p-n)$ . In addition one can extract the separate contributions  $\sigma_q(N)$ , q = u, d, N = p, n. The results are the following:  $\sigma_u(p) = 26.23$  MeV,  $\sigma_d(p) = 42.42$  MeV,  $\sigma_u(n) = 23.82$  MeV,  $\sigma_d(n) = 46.48$  MeV, which checks with  $\sigma_{\pi N} = \hat{m}(\sigma_u/m_u + \sigma_d/m_d)$ . The relation  $\sigma_u(p) = \sigma_d(n)$  in the isospin symmetry limit is of course satisfied, but the naive quark model relation in the isospin limit  $\sigma_u(p) = 2\sigma_d(p)$  is significantly violated due to contributions by the SU(2) singlet component of the quark masses.
- xi) Obviously, the discussion can be extended to the rest of the  $\sigma$  terms for the different baryons and their various relations [29].
- xii) One can compare with an analysis in ordinary HBChPT without the decuplet. In that case  $\Delta_{GMO}$  requires  $\mathring{g}_A/F_{\pi}$  to be significantly larger (corresponding to  $g_A^N = 1.48$  at LO), which despite the lack of the decuplet contributions leads to values of the  $\sigma$  terms which are not very different but somewhat larger than the ones obtained here ( $\hat{\sigma} \sim 83$  MeV,  $\sigma_{\pi N} \sim$ 76 MeV). The difference lies in the fact that in ordinary HBChPT the corrections to the axial currents couplings have large  $N_c$  power violating contributions, which compounded with the larger value of  $\mathring{g}_A/F_{\pi}$  required by  $\Delta_{GMO}$  lead to a failure in describing the axial couplings obtained in LQCD at different quark masses [32], in particular their observed small quark mass dependencies.
- xiii) Although the approach followed in recent work [33] should be expected to give a result for  $\sigma_{\pi N}$  similar to the one obtained here, it is actually much smaller. It is not clear to the authors whether this may be entirely due to the different set of LQCD data. However, since  $\hat{\sigma}$  is accurately obtained with only the physical masses, the result of [33] would require a large negative  $\sigma_s$ , which seems to be unlikely within the present framework.

## 3. Summary

The  $\sigma$  terms of nucleons were calculated using SU(3) BChPT × 1/N<sub>c</sub>. From the physical octet and decuplet baryon masses a value of  $\hat{\sigma}$  is obtained which is much larger than the one predicted by a tree level baryon mass combination, in agreement with similar observations in calculations that included the decuplet baryons as explicit degrees of freedom. The " $\sigma$  term puzzle" is understood as the result of large non-analytic contributions to that mass combination, while the higher order corrections to the  $\sigma$  terms have natural magnitude. The intermediate spin 3/2 baryons play an important role in enhancing  $\hat{\sigma}$  and thus  $\sigma_{\pi N}$ . The analysis carried out here shows that there is compatibility in the description of  $\Delta_{GMO}$  and the nucleon  $\sigma$  terms. The value of  $\sigma_{\pi N} = 69 \pm 10$  MeV obtained here from including LQCD baryon masses agrees with the more recent results from  $\pi N$  analyses, where the increase in value

with respect to previous analyses has been understood as a result of the values of the input scattering lengths, and strongly disfavor the values from recent LQCD evaluations. The tension between results, which includes LQCD, remains as an important problem to which the present approach can hopefully contribute with useful insights. The resolution of that tension will in turn provide a validation test of the approach.

#### Acknowledgments

This work was supported by DOE Contract No. DE-AC05-06OR23177 under which JSA operates the Thomas Jefferson National Accelerator Facility. JMA acknowledges partial support from the MINECO (Spain) and the ERDF (European Commission) through grant No. FPA2016-77313-P. IPF and JLG acknowledge partial support through National Science Foundation through grants PHY-1307413 and PHY-1613951.

#### References

- [1] M. Gell-Mann, R.J. Oakes, B. Renner, Phys. Rev. 175 (1968) 2195-2199.
- [2] T.P. Cheng, R.F. Dashen, Phys. Rev. Lett. 26 (1971) 594.
- [3] L.S. Brown, W.J. Pardee, R.D. Peccei, Phys. Rev. D 4 (1971) 2801-2810.
- [4] J. Gasser, H. Leutwyler, M.P. Locher, M.E. Sainio, Phys. Lett. B 213 (1988) 85–90.
- [5] J. Gasser, H. Leutwyler, M.E. Sainio, Phys. Lett. B 253 (1991) 252-259.
- [6] J. Gasser, H. Leutwyler, M.E. Sainio, Phys. Lett. B 253 (1991) 260-264.
- [7] M.M. Pavan, I.I. Strakovsky, R.L. Workman, R.A. Arndt, PiN Newsl. 16 (2002) 110–115, arXiv:hep-ph/0111066.
- [8] J.M. Alarcón, J. Martin Camalich, J.A. Oller, Phys. Rev. D 85 (2012) 051503, arXiv:1110.3797.
- [9] M. Hoferichter, J. Ruiz de Elvira, B. Kubis, U.-G. Meißner, Phys. Rev. Lett. 115 (2015) 092301, arXiv:1506.04142.
- [10] M. Hoferichter, J. Ruiz de Elvira, B. Kubis, U.-G. Meißner, Phys. Rep. 625 (2016) 1–88, arXiv:1510.06039.
- [11] M. Hoferichter, J. Ruiz de Elvira, B. Kubis, U.-G. Meißner, Phys. Lett. B 760 (2016) 74–78, arXiv:1602.07688.
- [12] J. Ruiz de Elvira, M. Hoferichter, B. Kubis, U.-G. Meißner, J. Phys. G 45 (2018) 024001, arXiv:1706.01465.
- [13] A. Walker-Loud, H.-W. Lin, D. Richards, R. Edwards, M. Engelhardt, et al., Phys. Rev. D 79 (2009) 054502, arXiv:0806.4549.
- [14] S. Durr, et al., Phys. Rev. Lett. 116 (2016) 172001, arXiv:1510.08013.
- [15] Y.-B. Yang, A. Alexandru, T. Draper, J. Liang, K.-F. Liu, Phys. Rev. D 94 (2016) 054503, arXiv:1511.09089.
- [16] A. Abdel-Rehim, C. Alexandrou, M. Constantinou, K. Hadjiyiannakou, K. Jansen, C. Kallidonis, G. Koutsou, A. Vaquero Aviles-Casco, Phys. Rev. Lett. 116 (2016) 252001, arXiv:1601.01624.
- [17] G.S. Bali, S. Collins, D. Richtmann, A. Schaefer, W. Soeldner, A. Sternbeck, Phys. Rev. D 93 (2016) 094504, arXiv:1603.00827.
- [18] A. Bottino, F. Donato, N. Fornengo, S. Scopel, Astropart. Phys. 13 (2000) 215–225, arXiv:hep-ph/9909228.
- [19] A. Bottino, F. Donato, N. Fornengo, S. Scopel, Astropart. Phys. 18 (2002) 205–211, arXiv:hep-ph/0111229.
- [20] J.R. Ellis, K.A. Olive, C. Savage, Phys. Rev. D 77 (2008) 065026, arXiv:0801.3656.
- [21] V. Cirigliano, R. Kitano, Y. Okada, P. Tuzon, Phys. Rev. D 80 (2009) 013002, arXiv:0904.0957.
- [22] H. Leutwyler, PoS CD15 (2015) 022, arXiv:1510.07511.
- [23] J.M. Alarcón, L.S. Geng, J. Martin Camalich, J.A. Oller, Phys. Lett. B 730 (2014) 342–346, arXiv:1209.2870.
- [24] C. Alexandrou, V. Drach, K. Jansen, C. Kallidonis, G. Koutsou, Phys. Rev. D 90 (2014) 074501, arXiv:1406.4310.
- [25] E.E. Jenkins, Phys. Rev. D 53 (1996) 2625–2644, arXiv:hep-ph/9509433.
- [26] R. Flores-Mendieta, E.E. Jenkins, A.V. Manohar, Phys. Rev. D 58 (1998) 094028, arXiv:hep-ph/9805416.
- [27] R. Flores-Mendieta, M.A. Hernandez-Ruiz, C.P. Hofmann, Phys. Rev. D 86 (2012) 094041, arXiv:1210.8445.
- [28] A. Calle Cordon, J.L. Goity, Phys. Rev. D 87 (2013) 016019, arXiv:1210.2364.
- [29] I.P. Fernando, J.L. Goity, Phys. Rev. D 97 (2018) 054010, arXiv:1712.01672.
- [30] T.R. Hemmert, B.R. Holstein, J. Kambor, Phys. Lett. B 395 (1997) 89–95, arXiv: hep-ph/9606456.
- [31] T.R. Hemmert, B.R. Holstein, J. Kambor, J. Phys. G 24 (1998) 1831–1859, arXiv: hep-ph/9712496.
- [32] C. Alexandrou, K. Hadjiyiannakou, C. Kallidonis, Phys. Rev. D 94 (2016) 034502, arXiv:1606.01650.
- [33] M.F.M. Lutz, Y. Heo, X.-Y. Guo, arXiv:1801.06417.