The g_2 Structure Function: An Experimental Overview

K. Slifer

University of New Hampshire

Abstract. We will discuss recent results for the spin structure functions, with an emphasis on g_2 . High precision g_2 data allows for tests of the Burkhardt–Cottingham sum rule, and is needed to consistently evaluate higher twist effects.

Keywords: Spin Structure, Polarized Targets

PACS: 11.55.Hx, 25.30.Bf, 29.25.Pj, 29.27.Hj

INTRODUCTION

Four independent structure functions are needed for a complete description of nucleon structure. All spin-dependent effects are contained in g_1 and g_2 , while spin-independent effects are parameterized in F_1 and F_2 . A simple physical interpretation is given in the impulse approximation of the parton model as distributions of quark momentum and spin in the nucleon:

$$\begin{aligned} F_1(x) &= \frac{1}{2} \sum e_i^2 [q_i(x) + \bar{q}_i(x)] \\ F_2(x) &= 2xF_1(x) \\ g_1(x) &= \frac{1}{2} \sum e_i^2 \Delta q_i(x) \end{aligned} \tag{1}$$

Here the summation runs over all quark flavors.

Analysis of the spin structure functions (SSF) is typically facilitated via the Cornwall-Norton (CN) moments [1, 2]:

$$\begin{aligned} \Gamma_1^{(n)}(Q^2) &= \int_0^1 dx x^{n-1} g_1(x, Q^2) \\ \Gamma_2^{(n)}(Q^2) &= \int_0^1 dx x^{n-1} g_2(x, Q^2) \end{aligned} \tag{2}$$

where x is the Bjorken scaling variable, and Q^2 represents the positive definite four momentum transfer of the virtual photon, which mediates the interaction between the incident electron probe and the nucleon target. By convention, the superscript is usually dropped in the case of $n = 1$.

The Q^2 and x -dependence of the g_1 structure function has been measured [3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13] with impressive precision over a wide kinematic range. In this proceedings, we focus on the other structure function g_2 , which has been historically

neglected due to the technical difficulties of producing transversely polarized targets, and the lack of a simple interpretation of g_2 in the classic parton model.

THE BURKHARDT-COTTINGHAM SUM RULE

The first CN moment of the g_2 structure function is predicted to vanish by the Burkhardt-Cottingham (BC) sum rule [14]:

$$\Gamma_2 = \int_0^1 dx g_2(x, Q^2) = 0 \quad (3)$$

This sum rule arises from the unsubtracted dispersion relation for the spin-dependent virtual-virtual Compton scattering amplitude S_2 . It is expected to be valid for any value of Q^2 , although the reader is referred to [15] for a detailed discussion of scenarios which would lead to violations of the relation.

Fig. 1 displays existing world data for $\Gamma_2(Q^2)$ for the proton, neutron and ^3He . All available data are consistent with the nuclear sum rule for ^3He . Similarly, we find satisfaction of the neutron sum rule, within uncertainties, across several different experiments and a large range of Q^2 . The picture is not so clear for the proton. The E155 collaboration found their data to be inconsistent with the proton BC sum rule at $Q^2 = 5 \text{ GeV}^2$, and the only other data point is from the RSS collaboration at $Q^2 = 1.3 \text{ GeV}^2$. Two upcoming JLab experiments will greatly expand our knowledge of g_2^p with measurements planned [16, 17] in the ranges $0.02 < Q^2 < 0.4$ and $2.5 < Q^2 < 6.5 \text{ GeV}^2$.

The open symbols in Fig. 1 represent the experimentally measured data which typically covers the resonance region. To evaluate Eq. 3, the contributions from $x = 1$ and $x \rightarrow 0$ must be included. The nucleon elastic contribution at $x = 1$ can be easily evaluated using the form factor parametrizations of [22, 23], and the nuclear elastic contribution for ^3He is similarly determined from [24]. All form factor parameterizations are known to high precision over the relevant region.

To estimate the $x \rightarrow 0$ contribution for the g_2 integrals, the Wandzura-Wilczek [25] relation is used:

$$g_2^{WW}(x, Q^2) \equiv -g_1(x, Q^2) + \int_x^1 \frac{g_1(y, Q^2)}{y} dy \quad (4)$$

This relation gives a prediction for the leading twist behaviour of g_2 entirely in terms of g_1 . Using Eq. 4 for the low x contribution of the integral relies on the assumption that higher twist effects asymptotically vanish as $x \rightarrow 0$, which is, of course, untested due to the difficulty in measuring structure functions in this region. This contribution to the integral represents the largest contribution to the overall systematic uncertainty.

What can the BC sum rule tell us about low x ?

Due to the extreme low counting rate, precise g_2 data at low Q^2 and low x are unlikely to be obtained anytime in the near future. However, if one were to accept the BC sum rule

FIGURE 1. World data for $\Gamma_2(Q^2)$ for the proton (top), neutron (middle) and ^3He (bottom). Open symbols represent measured data (typically the resonance region), while the full symbols include an estimate of the unmeasured contributions to the integral. Inner (outer) error bars represent statistical (total) uncertainties. Brown: E155 collaboration [4]. Red: RSS [18, 19]. Black: E94010 [20, 11]. Green: E97110 [21] (Very Preliminary). Blue: E01012 [13] (Very Preliminary).

purely on theoretical grounds, it is possible to invert Eq. 3 to provide information on g_2 in this region. The high precision of the elastic form factors and the measured resonance region data allows us to determine the $x \rightarrow 0$ contribution to $\Gamma_2(Q^2)$ with relatively small uncertainty. Fig. 2 shows the resonance region (open symbols) and elastic contribution (solid line) to Eq. 3. The solid band is the $x \rightarrow 0$ contribution inferred by assuming that Eq. 3 vanishes.

Violations of the BC Sum Rule?

As a necessary caveat to the preceding discussion, we stress that the experimental verification of the BC sum rule is still in question. In particular, Table 1 gives some perspective on the relative uncertainty of the determinations of Γ_2^n in the vicinity of $Q^2 \approx 1.0 \text{ GeV}^2$. Satisfaction of the neutron BC sum rule is found only at the 2 to 2.5 σ level of the total uncertainty. As can be seen from the table, the uncertainty is typically dominated

FIGURE 2. Full band represents an evaluation of $x \rightarrow 0$ contribution to $\Gamma_2(Q^2)$ assuming the validity of the BC sum rule. Proton (top), neutron (middle), and ^3He (bottom). Open symbols represent measured data (typically the resonance region). Inner (outer) error bars represent statistical (total) uncertainties. Brown: E155 collaboration [4]. Red: RSS [18, 19]. Black: E94010 [20, 11]. Green: E97110 [21] (Very Preliminary). Blue: E01012 [13] (Very Preliminary). Nucleon form factors from [22, 23]. Nuclear form factors from [24].

TABLE 1. Standard deviations of $\Gamma_2^n(Q^2)$ from zero. From E94010 [20] for $Q^2 < 1.0 \text{ GeV}^2$ and E01012 [13] (Very Preliminary) for $Q^2 > 1.0 \text{ GeV}^2$

$Q^2(\text{GeV}^2)$	Total Relative Uncertainty	Statistical Relative Uncertainty
0.74	1.7	6.6
0.90	2.0	7.9
1.20	2.4	2.9
1.80	1.9	2.2
2.40	1.8	2.2

by systematic effects, while the statistical uncertainties are quite small. The dominant systematics of E94010 arose from the radiative corrections. Considering the significant improvements in knowledge of structure functions at low Q^2 that has occurred in recent years, a re-examination of this uncertainty is warranted in order to provide a conclusive test of the BC sum rule.

FIGURE 3. $\Gamma_2^n(Q^2)$. Full integral for neutron. Inner (outer) error bars represent statistical (total) uncertainties. Brown: E155 collaboration [4]. Red: RSS [18, 19]. Black: E94010 [20, 11]. Green: E97110 [21] (Very Preliminary). Blue: E01012 [13] (Very Preliminary).

HIGHER TWIST MEASUREMENTS

The Operator Product Expansion (OPE) [26, 27, 28] relates the structure functions to QCD quark and gluon matrix elements, which contain information on parton interactions. This effective theory of QCD is formulated in terms of the CN moments of Eq. 2. The OPE provides the following expansion of the lowest moment $\Gamma_1(Q^2)$:

$$\Gamma_1(Q^2) = \sum_{\tau=2,4,\dots} \frac{\mu_\tau(Q^2)}{Q^{\tau-2}} \quad (5)$$

This expansion is performed in inverse powers of the *twist* τ . Successful predictions from the parton model map onto the leading twist of the OPE, while deviations from leading twist behaviour are known as higher twist effects. The parton model says nothing about the g_2 structure function, making it an excellent quantity to study higher twist.

The coefficients μ_τ are identified with nucleon matrix elements of QCD operators of twist $\leq \tau$. The first coefficient beyond leading twist is μ_4 , which can be decomposed as:

$$\mu_4 = \frac{1}{9}M^2 (\tilde{a}_2 + 4\tilde{d}_2 + 4\tilde{f}_2) \quad (6)$$

FIGURE 4. $I(Q^2)$ for proton (top) and neutron (bottom). Brown: E155 collaboration [4]. Black open square: QCDSF [29] Red: RSS [18, 19]. Black circle: E94010 [20]. Green: E97110 [21] (Very Preliminary). Blue: E01012 [13] (Very Preliminary). Magenta: E99117 [30]. Solid line: MAID model [31]. In the top panel, the filled band represents a determination of $I(Q^2)$ based on existing world data [10]. The large uncertainty reflects the lack of knowledge of g_2^p in this region.

This involves a combination of the leading twist, twist-3 and twist-4 matrix elements, respectively.

Higher Moments

The twist-2 term $\tilde{a}_2$ is purely kinematical in nature and can be evaluated [32] from the (target mass corrected) third moment of g_1 :

$$\int_0^1 x^2 g_1(x, Q^2) dx = \frac{1}{2} \tilde{a}_2 + \mathcal{O}\left(\frac{M^2}{Q^2}\right) \quad (7)$$

Similarly, the third moment of g_1 provides information on $\tilde{d}_2$:

$$\int_0^1 x^2 g_2(x, Q^2) dx = \frac{1}{3} (\tilde{d}_2 - \tilde{a}_2) + \mathcal{O}\left(\frac{M^2}{Q^2}\right) \quad (8)$$

FIGURE 5. $M_1^P(Q^2)$ from [10] and RSS [18, 19] (Preliminary). Statistical errors are shown on the data points, while systematic errors are represented by the full bands on the horizontal axis.

Eqs. 7 and 8 are often combined to extract the twist-3 matrix element via:

$$I(Q^2) = \int_0^1 dx x^2 (2g_1 + 3g_2) \quad (9)$$

$$= \tilde{d}_2(Q^2) + \mathcal{O}\left(\frac{M^2}{Q^2}\right) \quad (10)$$

However, note that $I(Q^2)$ is not equivalent to the twist-3 matrix element $\tilde{d}_2$ in any kinematic region where M^2 is not negligible compared to Q^2 [32]. World data on $I(Q^2)$ is shown in Fig. 4. In the top panel, the filled band represents a determination of $I(Q^2)$ based on existing world data [10]. The large uncertainty reflects the lack of knowledge of g_2^P in this region.

TARGET MASS CORRECTIONS

As emphasized in [32], dynamical higher twists can be extracted from the measured SSFs by using the Nachtmann moments [33, 34], which utilize the variable $\xi = 2x/(1 +$

FIGURE 6. RSS (very preliminary) R_2 data compared to the prediction of Ref. [32]

$\sqrt{1 + (2xM)^2/Q^2}$). The moments are defined:

$$M_1^n(Q^2) \equiv \frac{\tilde{a}_{n-1}}{2} \equiv \frac{1}{2} a_n E_1^n(Q^2, g) = \int_0^1 \frac{dx}{x^2} \xi^{n+1} \left[\left\{ \frac{x}{\xi} - \frac{n^2}{(n+2)^2} \frac{M^2 x \xi}{Q^2} \right\} g_1 - \frac{4n}{n+2} \frac{M^2 x^2}{Q^2} g_2 \right] \quad (11)$$

$$M_2^n(Q^2) \equiv \frac{\tilde{d}_{n-1}}{2} \equiv \frac{1}{2} d_n E_2^n(Q^2, g) = \int_0^1 \frac{dx}{x^2} \xi^{n+1} \left[\frac{x}{\xi} g_1 + \left\{ \frac{n}{n-1} \frac{x^2}{\xi^2} - \frac{n}{n+1} \frac{M^2 x^2}{Q^2} \right\} g_2 \right] \quad (12)$$

In the limit $M^2/Q^2 \rightarrow 0$, (i.e. TMC vanish) the Nachtmann moments reduce to the more familiar CN moments:

$$M_1^1(Q^2) \rightarrow \Gamma_1(Q^2) \quad (13)$$

$$2M_2^3(Q^2) \rightarrow I(Q^2) \quad (14)$$

At finite Q^2 , the ratio of Nachtmann to CN moments gives a quantitative measure of the TMC:

$$R_1(Q^2) = \frac{M_1^1(Q^2)}{\Gamma_1(Q^2)} \quad (15)$$

$$R_2(Q^2) = \frac{2M_2^3(Q^2)}{I(Q^2)} \quad (16)$$

The Nachtmann moments allow a clean separation of dynamical higher twists. In particular, Ref. [32] shows that M_2^3 gives access to the twist-3 matrix element that is accurate to $\mathcal{O}(M^8/Q^8)$.

A comparison of RSS data for M_1^1 to the analysis of Ref. [10] is shown in Fig. 5. Ref. [32] predicts that experimental determinations of $I(Q^2)$ significantly overestimate contributions to the twist-3 matrix element for $Q^2 < 10 \text{ GeV}^2$. Fig. 6 provides support for this conclusion from preliminary RSS proton data.

CONCLUSION

We have presented an overview of existing tests of the Burkhardt–Cottingham sum rule. All neutron and ^3He data support the sum rule prediction within experimental uncertainties, although the agreement is only at the 2.5σ level for the neutron. Proton data is found to be lacking. We have also presented a Nachtmann moment analysis of existing proton g_2 data, which shows that CN moments can significantly overestimate determinations of the twist-3 matrix element.

ACKNOWLEDGMENTS

I would like to thank the spokesmen of RSS: Mark Jones and Oscar Rondon, E97110, Alexandre Deur, J.P. Chen and Franco Garibaldi, and E01012: Nilanga Liyanage, J.P. Chen and Seonho Choi for kindly allowing me to present their preliminary data here. Many thanks also to Vince Sulkosky and Patricia Solvignon for providing the raw data. I would also like to thank Oscar Rondon for many instructive discussions on the topic material.

REFERENCES

1. J. M. Cornwall, and R. E. Norton, *Phys. Rev.* **177**, 2584–2586 (1969).
2. W. Melnitchouk, R. Ent, and C. Keppel, *Phys. Rept.* **406**, 127–301 (2005), hep-ph/0501217.
3. P. L. Anthony, et al., *Phys. Lett.* **B493**, 19–28 (2000), hep-ph/0007248.
4. P. L. Anthony, et al., *Phys. Lett.* **B553**, 18–24 (2003).
5. A. Airapetian, et al., *Phys. Rev.* **D75**, 012007 (2007), hep-ex/0609039.
6. M. Amarian, et al., *Phys. Rev. Lett.* **89**, 242301 (2002).
7. M. Amarian, et al., *Phys. Rev. Lett.* **93**, 152301 (2004).
8. Z. E. Meziani, et al., *Phys. Lett.* **B613**, 148–153 (2005), hep-ph/0404066.
9. M. Osipenko, et al., *Phys. Lett.* **B609**, 259–264 (2005), hep-ph/0404195.

10. M. Osipenko, et al., *Phys. Rev.* **D71**, 054007 (2005), hep-ph/0503018.
11. K. Slifer, et al., *Phys. Rev. Lett.* **101**, 022303 (2008), 0803.2267.
12. Y. Prok, et al. (2008), 0802.2232.
13. P. Solvignon, et al., *Phys. Rev. Lett.* **101**, 182502 (2008), 0803.3845.
14. H. Burkhardt, and W. N. Cottingham, *Annals Phys.* **56**, 453–463 (1970).
15. R. L. Jaffe, *Comments Nucl. Part. Phys.* **19**, 239 (1990).
16. A. Camsonne, J.P. Chen and K. Slifer, JLab Experiment E08-027. (???)
17. S. Choi, M. Jones, Z.-E. Meziani and O.A. Rondon, JLab Experiment E07-003. (???)
18. K. Slifer, et al. (2008), 0812.0031.
19. F. R. Wesselmann, et al., *Phys. Rev. Lett.* **98**, 132003 (2007), nucl-ex/0608003.
20. M. Amarian, et al., *Phys. Rev. Lett.* **92**, 022301 (2004), hep-ex/0310003.
21. A. Deur, J.P. Chen and F. Garibaldi, JLab Experiment E97-110. (???)
22. E. L. Lomon, *Phys. Rev.* **C66**, 045501 (2002), nucl-th/0203081.
23. P. Mergell, U. G. Meissner, and D. Drechsel, *Nucl. Phys.* **A596**, 367–396 (1996), hep-ph/9506375.
24. A. Amroun, et al., *Nucl. Phys.* **A579**, 596–626 (1994).
25. S. Wandzura, and F. Wilczek, *Phys. Lett.* **B72**, 195 (1977).
26. K. G. Wilson, *Phys. Rev.* **179**, 1499–1512 (1969).
27. J. Kodaira, Y. Yasui, and T. Uematsu, *Phys. Lett.* **B344**, 348–354 (1995), hep-ph/9408354.
28. B. Ehrnsperger, A. Schafer, and L. Mankiewicz, *Phys. Lett.* **B323**, 439–445 (1994), hep-ph/9311285.
29. M. Gockeler, et al., *Phys. Rev.* **D63**, 074506 (2001), hep-lat/0011091.
30. X. Zheng, et al., *Phys. Rev.* **C70**, 065207 (2004), nucl-ex/0405006.
31. D. Drechsel, O. Hanstein, S. S. Kamalov, and L. Tiator, *Nucl. Phys.* **A645**, 145–174 (1999), nucl-th/9807001.
32. Y. B. Dong, *Phys. Rev.* **C77**, 015201 (2008).
33. S. Matsuda, and T. Uematsu, *Nucl. Phys.* **B168**, 181 (1980).
34. A. Piccione, and G. Ridolfi, *Nucl. Phys.* **B513**, 301–316 (1998), hep-ph/9707478.