Proton- η' interactions at threshold

A. V. Anisovich^{a,b}, V. Burkert^c, M. Dugger^d, E. Klempt^{a,c}, V. A. Nikonov^{a,b}, B. G. Ritchie^d, A. Sarantsev^{a,b}. U. Thoma^a

> ^aHelmholtz-Institut für Strahlen- und Kernphysik. Universität Bonn. Germany ^bNRC "Kurchatov Institute", PNPI, Gatchina 188300, Russia ^c Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606 ^dArizona State University, Tempe, Arizona 85287-1504

Abstract

Abstract Recent data on photoproduction of η' mesons off protons have been included in the data base for the Bonn-Gatchina partial wave analysis. The real and imaginary parts of the *S*-wave $\eta' p \rightarrow \eta' p$ scattering amplitude in the threshold region were fit to yield the ηp scattering length and the interaction range. This new analysis found $|a_{\eta'p}| = (0.403 \pm 0.015 \pm 0.015$

ing the possibility of the existence of $\eta' N$ bound states in nuclear matter. Even the existence of an η -deuteron bound state in vacuum has been suggested [11]. A very recent determination of the η' -nucleus potential gave – with carbon [12] or niobium [13] as nuclei – a shallow potential of $-39 \pm 7_{\text{stat}} \pm 15_{\text{syst}} \text{ MeV}$ [14] for η' -mesons with momentum of 1200 - 2900 MeV/c. A new determination of the η' -nucleus potential with a mean η' momentum of $600 \,\mathrm{MeV/c}$ is in progress [15]. The scattering length and the range parameter for the η' -proton interaction also are directly related to the existence of any ηN bound states,

$$\gamma p \to \eta' p$$
 (1)

with the aim of identifying the contributions to this reaction from different N^* resonances, and to determine the $N^* \rightarrow N\eta'$ branching ratios [20]. Four resonances, $N(1895)1/2^{-}$, $N(1900)3/2^{+}$, $N(2100)1/2^{+}$, and $N(2120)3/2^{-}$ were found to provide the most significant contributions. The fit used the differential cross sections $d\sigma/d\Omega$ from the Crystal Barrel [21] and CLAS [22] experiments, along with recent beam asymmetry data from GRAAL Σ [17] and CLAS Σ [18] mentioned above. In the analysis presented here, we also used the new precise MAMI-A2 data on the $\gamma p \rightarrow \eta' p$ differential cross section [19]. In addition to the data on η' photoproduction, a large body of pion and



Figure 1: (Color online) The beam asymmetry Σ for the reaction $\gamma p \rightarrow \eta' p$. Shown are data from GRAAL [17] (first two subfigures) and recent data from CLAS [18] (next eight subfigures). The curves represent two fits: the solid (black) curve represents the main fit without a new narrow resonance, the dashed (red) curve a fit which includes a narrow $\eta' p$ threshold resonance with spin-parity $J^P = 3/2^-$. The partial-wave analysis curves for the CLAS beam asymmetries are scaled by a factor 0.94 (see Ref. [20]).

photo-induced reactions is fit in a coupled channel analysis. The data base includes the real and imaginary part of the πN scattering amplitude for the partial waves up to $J = 9/2^{\pm}$ from Ref. [23] and data on pion and photoproduction data with πN , ηN , $K \Lambda$, $K \Sigma$, $N \pi^0 \pi^0$, and $N \pi^0 \eta$ in the final state. A list of the data with references can be found on our web page (pwa.hiskp.uni-bonn.de/).

Similar fits - even though mostly restricted to the $\gamma N \rightarrow$ $\eta' N$ reaction - have recently been presented in Refs. [24– 27]. In Ref. [24], a quark model is used to fit differential cross sections for η' photoproduction off protons [21, 22] and neutrons [28]; Ref. [25] includes data on cross sections for $\pi^- p \to \eta' n$ and on $NN \to NN\eta'$ [29] (references to $\pi^- p$ and earlier NN data can be found in Ref. [25]). The authors of Ref. [27] fit CLAS 2009 differential cross sections [22] and the total cross section from MAMI-A2 collaboration [19] within an isobar model. The analyses [20, 24, 25, 27] agree that nucleon resonances should be included even though there is no consensus concerning the spin-parities of the preferred resonances. No explicit resonances were included in the analysis presented in Ref. [26], but instead final-state interactions between η' mesons and nucleons were studied within a three-flavor linear σ model. The need of final-state interactions is demonstrated even though the quality of the fit is moderate. The recent CLAS data on the $\gamma p \rightarrow \eta' p$ beam asymmetry, which include a much wider energy range than that provided by GRAAL, [18] have to date only been included in Ref. [20].

The formalism used here to fit the data is described in Ref. [20]. The fit solution presented in Ref. [20] predicted the new MAMI-A2 data [19] on the differential cross section for reaction (1) reasonably well; the inclusion of the data did not change the resonances contributing to η' photoproduction, and the changes in their $N\eta'$ decay branching ratios were small. Note that the errors given in [20] were dominantly due to a variation of the model assumptions. The new MAMI-A2 data stabilize the amplitudes close to the $\eta'p$ threshold.

Figure 1 shows the GRAAL [17] and CLAS [18] data on the beam asymmetry Σ , while Fig. 2 gives differential cross section measured at MAMI-A2 [19]. The data are compared to two fits:

1. Our standard fit is represented by the solid curves. This standard fit gives a reasonable description of the data except for the region just above the threshold. This fit predicts a vanishing beam asymmetry for the mass range where the GRAAL data exist, and a relatively flat angular distribution for the differential cross section. However, in contrast to the predictions of this standard fit, the data GRAAL data show an appreciably beam asymmetry at threshold, and the new MAMI-A2 data indicate a significant forward rise of the differential cross section in the 1899.5 to 1902.7 MeV mass range.

2. We tried to improve the standard fit by including a narrow $N\eta'$ resonance. The narrow resonance was represented by a convolution of a squared Breit-Wigner amplitude and a Gaussian function representing the resolution. The GRAAL resolution was fixed to 16 MeV (FWHM), for the MAMI-A2 data the resolution is expected to be given by the bin width (which corresponds to $\sigma \approx 3.2/\sqrt{12} \approx 1 \text{ MeV}$), and is neglected. The narrow resonance requires four additional parameters: M, Γ , and the product of helicities $A_{1/2}, A_{3/2}$ and the square root of the $N\eta'$ decay branching fraction.

Different spin-parity combinations were tested: $J^P = 1/2^{\pm}, 3/2^{\pm}, 5/2^{\pm}, 7/2^{\pm}$. Fits with $J^P = 1/2^{\pm}$ and $J^P = 7/2^{\pm}$ gave no improvement; the best fit was achieved with quantum numbers $J^P = 3/2^-$ but $5/2^-$ could also be pos-



Figure 2: (Color online) The MAMI-A2 differential cross section for $\gamma p \rightarrow \eta' p$ [19] and two BnGa fits. The dashed curve represents a fit which includes a narrow $\eta' p$ threshold resonance with spin-parity $J^P = 3/2^-$, the solid curve the main fit without a narrow resonance.

sible. For the first two bins in Fig. 1, the beam asymmetry was calculated for 1 MeV wide bins and then averaged. For the other bins, the beam asymmetries were calculated for the central masses. The dashed curves in Figs. 1 show this fit. Assuming $J^P = 3/2^-$, the fit returns

$$M_{\eta' p} = 1900 \pm 1 \,\text{MeV}; \qquad \Gamma_{\eta' p} < 3 \,\text{MeV}.$$
 (2)

When the narrow resonance is included, the χ^2 of the fit improves from 120.3 to 59.9 for the 70 data points in the first five Mainz mass bins, or from 29.5 to 11.7 for the 14 GRAAL data points. Due to the strongly rising phase space, the narrow resonance entails a small $J^P = 3/2^-$ amplitude extending over more than 10 MeV. Note that the mass resolution is given by the photon beam energy and not by the reconstruction of the final state.

The existence of such a narrow resonance in the *D*-wave is unexpected. (In the *S*-wave, a $N\eta'$ bound state just below the threshold is predicted in the linear sigma model [30]). To trace the origin of the narrow structure, we have excluded the MAMI-A2 data from the fit. If we impose the GRAAL resolution of 16 MeV (FWHM), the narrow width is confirmed. Decreasing the resolution leads to a larger natural width of the narrow resonance. Thus, the evidence



Figure 3: (Color online) Prediction for the double-polarization observables E, G, P and T for the 1901 - 1902 MeV mass window. The solid (black) curve represents the main fit without a new narrow resonance, the dashed (red) curve a fit represents the prediction for the case when a narrow $\eta' p$ threshold resonance with spin-parity $J^P = 3/2^-$ exists.

for the narrow width of the resonance rests mostly on the MAINZ $\gamma p \rightarrow \eta' p$ differential cross section [19].

The GRAAL data suggest the possible existence of a $p\eta'$ threshold resonance; the MAMI-A2 differential cross sections support this conjecture. Hence we think that the search for a narrow $N\eta'$ resonance should be continued with new data with high statistics and precision. A measurement of further polarization observables might help to find an unambiguous answer. The observables for which predictions are made include those for several polarized-photon-beam/polarized-proton-target combinations.

Figure 3 shows the predictions for the polarization observables E, H, P and T for a 2 MeV mass region at the nominal mass of the possible $\eta'p$ resonance. The observable E is the (normalized) difference between the meson photoproduction cross sections for helicity 1/2 and helicity 3/2. The observable H is the correlation between linearlypolarized photons and transversely-polarized protons. The observable P is the polarization of the outgoing proton with respect to the scattering plane. The observable T is the asymmetry in the production cross section when the target proton polarization transverse to the incident photon beam is flipped.

When a resonance is added to the fit, significant differences in one or more of these observables should arise. In particular, data for P and H should indicate the presence of a resonance (even if the resonance is narrow) if such a structure exists, and predictions for those observables are relatively sensitive to the presence of that state. However, we also recognize that experiments so near to threshold are very demanding, especially if a putative resonance is so narrow that the polarization observables can only show evidence for that resonance in a small mass window.

We now turn to the main objective of this paper: the determination of the $\eta'p$ scattering length. The scattering length is given by the $J^P = 1/2^-$. (S_{11}) -wave amplitude for $\eta'p$ production at the η' threshold (corresponding to the wave in which the $\eta'p$ orbital angular momentum vanishes). As the primary solution, we use the *standard fit* without additional narrow resonance. The upper and lower panels of Figure 4 show the real and imaginary parts, respectively, of the S_{11} partial wave in the low-energy region, from the threshold to 10 MeV above. A fit to the squared amplitude with $\beta \cdot \sqrt{s - s_0}$ yields an offset of 1896.0 \pm 1.0 MeV, fully consistent with the sum of proton and η' mass: 1896.05 \pm 0.06 MeV.

The results shown in Figs. 4a and b were fit with a function

$$A = \frac{a k}{1 - i k a + R k^2 a / 2 + d k^4 a}$$
(3)
$$k = \frac{\sqrt{(s - (M_p + M_{\eta'})^2)}}{4 s}$$

where $a = a_{p\eta'}$ is the $\eta' p$ scattering length, R the interaction range and d a parameter representing higher-order terms. The fit to Figs. 4a and b give consistent results for the modulus of the $\eta' p$ scattering length.

The S_{11} amplitude has one arbitrary phase which cannot be defined from experiment. The phase could be defined as zero at the pion-production threshold or, alternatively, at the η' production threshold.

We fit the amplitude in the range from threshold to 10 MeV above the threshold, the parameter d was fixed to zero or was used as a free fit parameter. From these fits, we obtained a modulus of the scattering length in the range $0.367 > |a_{p\eta'}| > 0.344$ fm with a statistical error (σ) of less than ± 0.013 fm. Then we performed fits with an additional narrow resonance, again with d = 0 and with das free fit parameter. In this case the fit returned values between $0.462 > |a_{p\eta'}| > 0.394$ fm and a statistical error of ± 0.020 fm. The modulus of the $\eta' p$ scattering length depends only weakly on the existence of a narrow *D*-wave resonance at the $\eta' p$ threshold. Since we do not know if a narrow resonance exists or not, we quote

$$|a_{p\eta'}| = (0.403 \pm 0.020 \pm 0.060) \,\mathrm{fm} \tag{4}$$

as our final result, where the first uncertainty is statistical, the second one is nature. The phase relative to the η' production threshold is

$$\delta = (1.5 \pm 0.5)^{\circ}; \tag{5}$$

With this definition of the phase, the imaginary part of the amplitude nearly vanishes. This scattering length would lead to a value for the real part of the η' potential of



Figure 4: (Color online) Real part (upper panel) and imaginary part (lower panel) of the $\eta' N \rightarrow \eta' N$ S-wave scattering amplitude A (solid curve) using Eqn. (3). The phase is rotated to have a vanishing imaginary part at the $\eta' p$ threshold. The error band is calculated from the variance of the partial-wave analysis coefficients.

 $33\pm5\,{\rm MeV}$ and a very small ($\approx 1\,{\rm MeV})$ imaginary part, making the search for $\eta'\text{-nucleus}$ bound states particularly attractive.

The phase shift between the πN and the $\eta' p$ threshold is rather stable. If the phase is defined relative to the pion-production threshold, we find

$$\delta = (87 \pm 2)^{\circ}.\tag{6}$$

These values are consistent with the numbers obtained at COSY [16]: $\operatorname{Re}(a_{p\eta'}) = 0 \pm 0.43$ fm and $\operatorname{Im}(a_{p\eta'}) = 0.37 \stackrel{+0.40}{-0.16}$ fm.

The phase of about 90° implies that the real part of the scattering length is small compared to imaginary part. The existence of a $\eta' p$ (S-wave) bound state would require the real part to be larger than the imaginary part [32]. This value for the scattering length does not, however, exclude the existence of a $\eta' N$ bound state in a nuclear environment or the existence of a D-wave $\eta' N$ resonance.

The range parameter is a complex number which remained essentially undetermined. We find $R = (3.2 \pm 2.1) + i(1.6 \pm 1.2)$ fm for fits with d = 0 and $R = (10 \pm 5) + i(5 \pm 5)$ fm when d is varied freely. There was wide variation of the estimates for the value of (and uncertainty for) d, so no specific value for that quantity is quoted here.

In all fits, $N\eta$ decay modes of most resonances are admitted. The opening of the $\eta'p$ threshold can be seen in the data as pointed out in Ref. [19]. The data are shown in Fig. 5 and compared to the BnGa fit. The drop of the cross section at about 1890 MeV is due to the opening of the η' threshold and indicates the presence of a resonance



Figure 5: (Color online) Total cross section for $\gamma p \rightarrow \eta p$ Ref. [19]. The significant drop at the $\eta'p$ threshold near 1890 MeV indicates the presence of strong contributions from the $N(1895)1/2^{-}$ resonance. The data shown come from two running periods. The cross sections for the high-mass data above 1880 MeV have been scaled with a factor 1.06.

at about this mass with quantum number $J^P = 1/2^-$.

Summarizing, we have studied the photoproduction reaction $\gamma p \to \eta' p$. The GRAAL data on the beam asymmetry for this reaction suggest the possible existence of a narrow $\eta' p$ resonance at 1900 \pm 1 MeV and a width of less than 3 MeV. The $\eta' p$ scattering length has been determined. Its magnitude is found to be $|a_{p\eta'}| = (0.403 \pm$ 0.020 ± 0.0600) fm, its phase relative to the πN threshold is $\delta = (87 \pm 2)^{\circ}$. This value does not depend on the exis tence or not of the narrow $\eta' p$ resonance. The range of the interaction could not be determined with a reasonable accuracy.

We would like to thank V. Metag for illuminating discussions on η' interactions in nuclei. This work was supported by the Deutsche Forschungsgemeinschaft (SFB/TR110) and by the RSF grant 16-12-10267. Work at Arizona State University was supported by the U.S. National Science Foundation under award PHY-1306737. This material is in part based upon work supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under contract DE-AC05-06OR23177.

References

- [1] D. Coffman et al. [MARK-III Collaboration], Phys. Rev. D 38, 2695 (1988).
- T. Feldmann, P. Kroll and B. Stech, Phys. Rev. D 58, 114006 [2](1998).
- R. Escribano, Eur. Phys. J. C 65, 467 (2010). [3]
- [4]F. G. Cao, Phys. Rev. D 85, 057501 (2012).
- R. Aaij et al. [LHCb Collaboration], JHEP 1501, 024 (2015). [5][6]
- L. A. Harland-Lang, V. A. Khoze, M. G. Ryskin and A. G. Shuvaev, Phys. Lett. B 770, 88 (2017).
- [7]G. 't Hooft, Phys. Rev. D 14, 3432 (1976) Erratum: [Phys. Rev. D 18, 2199 (1978)].
- V. Bernard and U. G. Meissner, Nucl. Phys. A 489, 647 (1988).
- [9] G. E. Brown and M. Rho, Phys. Rev. Lett. 66, 2720 (1991).

- [10] T. Hatsuda and S. H. Lee, Phys. Rev. C 46, no. 1, R34 (1992).
- [11] T. Sekihara, H. Fujioka and T. Ishikawa, "Possible $\eta'd$ bound state and its s-channel formation in the $\gamma d \rightarrow \eta d$ reaction," arXiv:1712.06257 [nucl-th].
- [12] M. Nanova et al. [CBELSA/TAPS Collaboration], Phys. Lett. B 727, 417 (2013).
- [13]M. Nanova et al. [CBELSA/TAPS Collaboration], Phys. Rev. C 94, no. 2, 025205 (2016).
- [14] V. Metag, M. Nanova and E. Y. Paryev, Prog. Part. Nucl. Phys. **97**, 199 (2017).
- [15] M. Nanova et al. [CBELSA/TAPS Collaboration], "The η '-Carbon potential at low meson momenta," in preparation.
- E. Czerwinski et al., Phys. Rev. Lett. 113, 062004 (2014).
- [17] P. Levi Sandri et al., Eur. Phys. J. A 51, no. 7, 77 (2015).
- [18] P. Collins et al. [CLAS Collaboration], Phys. Lett. B 771, 213 (2017).
- [19] V. L. Kashevarov et al., Phys. Rev. Lett. 118, no. 21, 212001 (2017).
- A. V. Anisovich et al., " $N^* \to N\eta'$ decays from photoproduc-[20]tion of η' -mesons off protons", submitted to Physics Letters B.
- [21] V. Crede et al. [CBELSA/TAPS Collaboration], Phys. Rev. C 80, 055202 (2009).
- [22]M. Williams et al. [CLAS Collaboration], Phys. Rev. C 80, 045213 (2009).
- [23]R. A. Arndt, W. J. Briscoe, I. I. Strakovsky and R. L. Workman, Phys. Rev. C 74, 045205 (2006).
- X. H. Zhong and Q. Zhao, Phys. Rev. C 84, 065204 (2011). [24]
- [25]F. Huang, H. Haberzettl and K. Nakayama, Phys. Rev. C 87, 054004 (2013)
- [26] S. Sakai, A. Hosaka and H. Nagahiro, Phys. Rev. C 95, no. 4, 045206 (2017).
- V. A. Tryasuchev, A. G. Kondratyeva and A. A. Kiziridi, Russ. [27]Phys. J. 60, no. 5, 782 (2017).
- [28] I. Jaegle et al. [CBELSA/TAPS Collaboration], Eur. Phys. J. A 47, 11 (2011).
- [29]P. Klaja et al., Phys. Lett. B 684, 11 (2010).
- [30] S. Sakai and D. Jido, Hyperfine Interact. 234, no. 1-3, 71 (2015). A. M. Green and S. Wycech, Phys. Rev. C 71, 014001 (2005) [31]Erratum: [Phys. Rev. C 72, 029902 (2005)]
- [32]Q. Haider and L. C. Liu, Int. J. Mod. Phys. E 24, no. 10, 1530009 (2015).