

Spectral functions of nucleon form factors: Three-pion continua at low energies¹

N. Kaiser¹ and E. Passemar^{2,3,4}

- 1) Physik-Department T39, Technische Universität München, D-85747 Garching, Germany
- 2) Department of Physics, Indiana University, Bloomington, IN 47405, USA
- 3) Center for Exploration of Energy and Matter, Indiana University, Bloomington, IN 47408, USA
- 4) Theory Center, Thomas Jefferson National Accelerator Facility, Newport News, VA 23606, USA

Abstract

We study the imaginary parts of the isoscalar electromagnetic and isovector axial form factors of the nucleon close to the 3π -threshold in covariant baryon chiral perturbation theory. At the two-loop level, the contributions arising from leading and next-to-leading order chiral πN -vertices, as well as pion-induced excitations of virtual $\Delta(1232)$ -isobars, are calculated. It is found that the heavy baryon treatment overestimates substantially these 3π -continua. From a phenomenological analysis, that includes the narrow $\omega(783)$ -resonance or the broad a_1 -resonance, one can recognize small windows near threshold, where chiral 3π -dynamics prevails. However, in the case of the isoscalar electromagnetic form factors $G_{E,M}^s(t)$, the radiative correction provided by the $\pi^0\gamma$ -intermediate state turns out to be of similar size.

1 Introduction

The structure of the nucleon as revealed in elastic electron-nucleon and (anti)neutrino-nucleon scattering is encoded in four electromagnetic form factors $G_{E,M}^{p,n}(t)$ and two axial form factors $G_{A,P}(t)$, with t the squared momentum-transfer. Dispersion theory is a tool to interpret (and cross check) these scattering data in a largely model independent way [1, 2]. The nucleon form factors are assumed to satisfy unsubtracted dispersion relations and their absorptive parts are often parametrized by a few vector meson poles. However, such an approach is not in conformity with general constraints from unitarity and analyticity. In particular, the singularity structure of the $\pi\pi N$ -triangle diagram leads to a pronounced enhancement of the isovector electromagnetic spectral functions on the left wing of the $\rho(770)$ -resonance. The two-pion intermediate state can actually be treated exactly (in the energy region $2m_\pi < \sqrt{t} < 1\text{ GeV}$) in terms of the pion charge form factor $F_\pi(t)$ and the p-wave πN partial wave amplitudes $f_\pm^1(t)$ in the crossed t -channel $\pi\pi \rightarrow \bar{N}N$. For the latter quantities improved results have been obtained in recent dispersion analyses of πN -scattering [3, 4] based on solutions of the Roy-Steiner equations. The calculation of the isovector electromagnetic spectral functions $\text{Im}G_{E,M}^v(t)$ in chiral perturbation theory up to two loops [5] is able to account (step by step) for the strong enhancement above the 2π -threshold (which originates from a logarithmic singularity at $t_c = 4m_\pi^2 - m_\pi^4/M^2 = 3.978 m_\pi^2$ on the second Riemann sheet), but an additional (adjusted) $\rho(770)$ -resonance contribution is necessary to reproduce reasonably the empirical spectral functions. On the other hand, the isoscalar electromagnetic form factors $G_{E,M}^s(t)$ are usually represented by sums of a few vector meson poles ($\omega, \phi, s_1, s_2, s_3$) [2, 6]. The

¹This work has been supported in part by DFG and NSFC (CRC110), the DFG cluster of excellence ‘Origin and Structure of the Universe’ (www.universe-cluster.de), the U.S. Department of Energy (contract DE-AC05-06OR23177) and the National Science Foundation (PHY-1714253).

effects of the 3π -continuum on the spectral functions $\text{Im}G_{E,M}^s(t)$ close to threshold have been calculated in ref. [7] using heavy baryon chiral perturbation theory, and it was concluded that these are small against the tails of an $\omega(783)$ -resonance with constant width. In that work [7] the axial spectral function $\text{Im}G_A(t)$ near the 3π -threshold was also computed and the analogous calculation for the induced pseudoscalar form factor $G_P(t)$ has been performed in ref. [8].

The purpose of the present paper is an improved calculation of the 3π -continua in covariant baryon chiral perturbation theory. This seems appropriate in view of the expected size of relativistic corrections: $\sqrt{t}/M > 3m_\pi/M = 0.44$. In addition to leading order chiral πN -vertices, we consider also the next-leading order ones (involving the low-energy constants c_1, c_2, c_3, c_4) and we treat the pion-induced excitation of the low-lying $\Delta(1232)$ -resonance, described by a Rarita-Schwinger spinor. Our paper is organized as follows. In section 2 we recapitulate the Cutkosky cutting rule applied to two-loop diagrams with a 3π -absorptive part and we present the Lorentz-invariant 3π -phase space integral in explicit form together with all kinematical variables. The formulas to project out individual nucleon form factors from the transition matrix elements of the vector and axial-vector currents are also given. Section 3 is devoted to the presentation and discussion of the calculated 3π -spectral function $\text{Im}G_{E,M}^s(t)$ and $\text{Im}G_A(t)$, separated into contributions from leading order chiral πN -vertices, next-to-leading order ones, and the inclusion of explicit Δ -isobars. In each case we give also convenient formulas, which refer to the non-relativistic approximation. In section 4 we perform a simple phenomenological analysis by considering the narrow $\omega(783)$ -resonance for $\text{Im}G_{E,M}^s(t)$ and the broad $a_1(1260)$ -resonance for $\text{Im}G_A(t)$. In the first case this draws our attention to electromagnetic effects and are thus compelled to compute the radiative correction to $\text{Im}G_{E,M}^s(t)$ provided by the $\pi^0\gamma$ -intermediate state. The paper ends with a summary and conclusions in section 5.

2 Calculation of three-pion spectral functions

We follow the (standard) notations for the electromagnetic and axial form factors of the nucleon as introduced in section 2 of ref. [7], and remind that these form factors are assumed to satisfy unsubtracted dispersion relations of the form:

$$F(t) = \frac{1}{\pi} \int_{t_0}^{\infty} dt' \frac{\text{Im}F(t')}{t' - t - i\epsilon}. \quad (1)$$

The threshold t_0 for hadronic intermediate states is $t_0 = 4m_\pi^2$ for the isovector electromagnetic form factors $G_{E,M}^v(t)$ and the scalar form factor $\sigma_N(t)$, while $t_0 = 9m_\pi^2$ for the isoscalar electromagnetic form factors $G_{E,M}^s(t)$ and the isovector axial form factors $G_{A,P}(t)$. The measurable electromagnetic form factors of the proton and neutron are composed of the isoscalar and isovector ones as:

$$G_{E,M}^{p,n}(t) = G_{E,M}^s(t) \pm G_{E,M}^v(t), \quad (2)$$

where the normalizations $G_E^{s,v}(0) = 1/2$, $G_M^s(0) = 0.440$ and $G_M^v(0) = 2.353$ hold at $t = 0$.

Figure 1 shows a generic two-loop diagram with a 3π -intermediate state contributing to the nucleon transition matrix element of the vector (or axial-vector) current. The three pions have four-momenta l_1, l_2, l_3 and a, b, c are their (cartesian) isospin-indices. Exploiting (perturbative) unitarity in the form of the Cutkosky cutting rules, one obtains for the imaginary (or absorptive)




Figure 1: Generic two-loop diagram $\gamma^* \rightarrow 3\pi \rightarrow \bar{N}N$ generating the three-pion spectral function.

part of the corresponding two-loop amplitude:

$$\text{Im } T_{2\text{-loop}} = -\frac{1}{2} \int d\Phi_3 A \cdot B, \quad (3)$$

where A denotes the S-matrix for $\gamma^* \rightarrow 3\pi$ and B the S-matrix for $3\pi \rightarrow \bar{N}N$ (in the subthreshold region $\sqrt{t} < 2M$). The integral $\int d\Phi_3$ goes over the Lorentz-invariant three-pion phase space, whose volume is determined by the kinematical variable $t = (l_1 + l_2 + l_3)^2 = (p_2 - p_1)^2$ and the mass m_π . In the center-of-mass frame the phase space integration can be expressed as a four-dimensional integral over two energies (ω_1, ω_2) and two angular variables (x, φ) by:

$$\int d\Phi_3 H(\dots) = \frac{1}{64\pi^4} \iint_{z^2 < 1} d\omega_1 d\omega_2 \int_{-1}^1 dx \int_0^\pi d\varphi H(\dots), \quad (4)$$

where z is determined by energy and momentum conservation as

$$|\vec{l}_1| |\vec{l}_2| z = \omega_1 \omega_2 - \sqrt{t} (\omega_1 + \omega_2) + \frac{t + m_\pi^2}{2}, \quad |\vec{l}_{1,2}| = \sqrt{\omega_{1,2}^2 - m_\pi^2}, \quad (5)$$

and $\sqrt{t} > 3m_\pi$ denotes the three-pion invariant mass. The directional cosines

$$x = \hat{l}_1 \cdot \vec{v}, \quad y = \hat{l}_2 \cdot \vec{v} = xz + \sqrt{(1-x^2)(1-z^2)} \cos \varphi, \quad (6)$$

refer to a unit-vector \vec{v} , which is introduced by the momentum of the nucleon.

Before eqs.(3,4) can be applied to calculate spectral functions, one has to project the (individual) electric and magnetic form factors out of the (isoscalar) current matrix element

$$\bar{u}_2 V^\mu u_1 = \bar{u}_2 \left[\gamma^\mu F_1(t) + \frac{i}{2M} \sigma^{\mu\nu} (p_2 - p_1)_\nu F_2(t) \right] u_1, \quad (7)$$

where $u_{1,2}$ free are Dirac-spinors. This is done by multiplying V^μ with on-shell projectors $\gamma \cdot p_{1,2} + M$ and taking suitable Dirac-traces:

$$G_E(t) = F_1(t) + \frac{t}{4M^2} F_2(t) = \frac{(p_1 + p_2)_\mu}{4M(4M^2 - t)} \text{tr} \{ V^\mu (\gamma \cdot p_1 + M) (\gamma \cdot p_2 + M) \}, \quad (8)$$

$$G_M(t) = F_1(t) + F_2(t) = \frac{1}{4t} \text{tr} \left\{ V^\mu (\gamma \cdot p_1 + M) \left[\gamma_\mu + \frac{2M}{t - 4M^2} (p_1 + p_2)_\mu \right] (\gamma \cdot p_2 + M) \right\}, \quad (9)$$

where $M = 939 \text{ MeV}$ denotes the (average) nucleon mass. The axial and pseudoscalar form factors are projected out of the isovector axial-current matrix element (proportional to $\tau_d/2$):

$$\bar{u}_2 A^\mu u_1 = \bar{u}_2 \left[\gamma^\mu G_A(t) + \frac{(p_2 - p_1)^\mu}{2M} G_P(t) \right] \gamma_5 u_1, \quad (10)$$

in a similar way:

$$G_A(t) = -\frac{1}{4(4M^2 - t)} \text{tr} \left\{ A^\mu (\gamma \cdot p_1 + M) \left[\gamma_\mu + \frac{2M}{t} (p_2 - p_1)_\mu \right] \gamma_5 (\gamma \cdot p_2 + M) \right\}, \quad (11)$$

$$G_P(t) = \frac{M}{t(4M^2 - t)} \text{tr} \left\{ A^\mu (\gamma \cdot p_1 + M) \left[M \gamma_\mu + \left(\frac{6M^2}{t} - 1 \right) (p_2 - p_1)_\mu \right] \gamma_5 (\gamma \cdot p_2 + M) \right\}. \quad (12)$$

In our calculation the S-matrices $A(\gamma^* \rightarrow 3\pi)$ and $B(3\pi \rightarrow \bar{N}N)$ are built from chiral vertices and hence the integrand $H(\dots)$ in eq.(4) becomes a rational function of the Lorentz scalar products:

$$l_1 \cdot l_2 = \sqrt{t}(\omega_1 + \omega_2) - \frac{t + m_\pi^2}{2}, \quad p_1 \cdot p_2 = M^2 - \frac{t}{2}, \quad (13)$$

$$l_1 \cdot p_{1,2} = \frac{1}{2} \left(\mp \sqrt{t} \omega_1 - ix \sqrt{4M^2 - t} |\vec{l}_1| \right), \quad (14)$$

$$l_2 \cdot p_{1,2} = \frac{1}{2} \left(\mp \sqrt{t} \omega_2 - iy \sqrt{4M^2 - t} |\vec{l}_2| \right). \quad (15)$$

We note that in the nonrelativistic limit $M \rightarrow \infty$ the nucleon propagators become complex-valued distributions:

$$\frac{-1}{ix - \epsilon} = \pi \delta(x) + iP \frac{1}{x}, \quad \frac{-1}{(ix - \epsilon)(iy + \epsilon)} = P \frac{1}{x} P \frac{1}{y} + \pi^2 \delta(x) \delta(y) + i\pi \left[P \frac{1}{x} \delta(y) - \delta(x) P \frac{1}{y} \right], \quad (16)$$

and the angular integrations $\int_{-1}^1 dx \int_0^\pi d\varphi$ can (and must) be performed analytically. For example, the outcomes of the two distributions in eq.(16) are π^2 and $2\pi(1 - z^2)^{-1/2} \arccos(-z)$.

3 Results of digrammatic calculation and discussion

In this section we present the results for the spectral functions $\text{Im}G_{E,M}^s(t)$ and $\text{Im}G_A(t)$ as calculated from leading-order chiral πN -vertices, next-to-leading order ones, and pion-induced excitations of virtual $\Delta(1232)$ -isobars. Since one works at all three stages with the same couplings of the external sources to three pions, we recapitulate these first. The momentum-dependent (anomalous) coupling of the virtual photon to three pions $\pi^a(l_1), \pi^b(l_2), \pi^c(l_3)$ reads [9]:

$$\frac{\epsilon_{abc}}{4\pi^2 f_\pi^3} \epsilon_{\mu\nu\alpha\beta} l_1^\nu l_2^\alpha l_3^\beta, \quad (17)$$

where the charge factor e has been dropped, and $f_\pi = 92.2 \text{ MeV}$ is the pion decay constant. On the other hand the momentum-dependent coupling of the axial source (with isospin-index d) to three pions $\pi^a(l_1), \pi^b(l_2), \pi^c(l_3)$ is given by [10]:

$$\frac{1}{f_\pi} \left[\delta_{ad} \delta_{bc} (l_2 + l_3 - l_1)^\mu + \delta_{bd} \delta_{ac} (l_1 + l_3 - l_2)^\mu + \delta_{cd} \delta_{ab} (l_1 + l_2 - l_3)^\mu \right]. \quad (18)$$




Figure 2: Two-loop diagrams contributing to the current matrix elements under consideration.




Figure 3: Spectral functions $\text{Im}G_E^s(t)$ (red), $\text{Im}G_M^s(t)$ (blue), and $\text{Im}G_A(t)$ (green) calculated with leading order chiral πN -vertices. The dashed lines correspond to the nonrelativistic approximation.

3.1 Leading-order chiral vertices

The four relevant two-loop diagrams that need to be evaluated are shown in Fig. 2. In addition to the well-known pseudovector πN -coupling and the (vectorial) Weinberg-Tomozawa vertex, one encounters for the axial form factor $G_A(t)$ the chiral contact-vertex with three pions $\pi^a(l_1), \pi^b(l_2), \pi^c(l_3)$ absorbed on a nucleon (see right half of diagram (d) in Fig. 2):

$$-\frac{g_A}{4f_\pi^3} [\tau_a \delta_{bc} \gamma \cdot (l_2 + l_3) + \tau_b \delta_{ac} \gamma \cdot (l_1 + l_3) + \tau_c \delta_{ab} \gamma \cdot (l_1 + l_2)] \gamma_5. \quad (19)$$

The rational integrand-functions $H(\dots)$ resulting from the Dirac-traces turn out to be quite lengthy, and are therefore not given here. The four-dimensional phase space integration in eq.(4) has been performed numerically, setting $g_A = 1.3$ (to have a strong πN -coupling constant $g_{\pi NN} = g_A M / f_\pi = 13.24$) and taking an average pion mass $m_\pi = 138$ MeV. The obtained leading order spectral functions $\text{Im}G_E^s(t)$, $\text{Im}G_M^s(t)$ and $\text{Im}G_A(t)$ in the low-energy region $3m_\pi < \sqrt{t} < 6m_\pi$ are shown in Fig. 3 by red, blue and green lines, respectively. A logarithmic scale is used to make visible the very small values in the threshold region. The dashed lines in Fig. 3 correspond to the nonrelativistic approximation. For the leading terms in the $1/M$ -expansion of the electromagnetic spectral functions one can actually give convenient formulas:

$$\begin{aligned} \text{Im}G_E^s(t) &= \frac{g_A}{8M(8\pi)^4 f_\pi^6 \sqrt{t}} \int_{2m_\pi}^{\sqrt{t}-m_\pi} dw (w^2 - 4m_\pi^2)^{3/2} \lambda(w, t) \\ &\quad + \frac{3g_A^3 t}{(4\pi)^5 f_\pi^6} \iint_{z^2 < 1} d\omega_1 d\omega_2 |\vec{l}_1| |\vec{l}_2| \sqrt{1 - z^2} \arccos(-z), \end{aligned} \quad (20)$$

$$\text{Im}G_M^s(t) = \frac{g_A M}{4(8\pi)^4 f_\pi^6 t^{3/2}} \int_{2m_\pi}^{\sqrt{t}-m_\pi} dw \sqrt{w^2 - 4m_\pi^2} \left[w^2 - 4m_\pi^2 + g_A^2 (5w^2 - 8m_\pi^2) \right] \lambda(w, t), \quad (21)$$

with the abbreviation $\lambda(w, t) = [t - (w + m_\pi)^2][t - (w - m_\pi)^2]$ and w denotes a 2π -invariant mass. The prefactor M in $\text{Im}G_M^s(t)$ originates from the magnetic coupling $i\vec{\sigma} \times (\vec{p}_2 - \vec{p}_1) / 2M$. The analogous (nonrelativistic) formula for $\text{Im}G_A(t)$ can be found in eq.(26) of ref. [7] and that for $\text{Im}G_P(t)$ in eqs.(4,5) of ref. [8]. Note that eq.(20) includes in the first line also the leading $1/M$ -correction to $\text{Im}G_E^s(t)$ coming from the two diagrams proportional to g_A . By inspection of Fig. 3 one observes that the heavy baryon treatment (used in ref. [7]) leads to an overestimation of the spectral functions near the 3π -threshold by about a factor of 2 to 3. This noteworthy property of the chiral 3π -continua points to sizeable relativistic corrections of magnitude $\sqrt{t}/M > 3m_\pi/M = 0.44$.

3.2 Second-order chiral vertices

Next, we compute the spectral functions with vertices from the second-order chiral πN -Lagrangian. The pertinent S-matrix for the absorption of two pions $\pi^a(l_1), \pi^b(l_2)$ on a nucleon reads:

$$-\frac{2i}{f_\pi^2} \delta_{ab} \left[2c_1 m_\pi^2 + \frac{c_2}{4M^2} (p + p') \cdot l_1 (p + p') \cdot l_2 + c_3 l_1 \cdot l_2 \right] + \frac{ic_4}{f_\pi^2} \epsilon_{abe} \tau_e \sigma_{\mu\nu} l_1^\mu l_2^\nu, \quad (22)$$

where p and p' denote in-going and out-going nucleon four-momenta. This 2π -contact vertex enters now diagrams (a) and (b) in Fig. 2. Note that due to the contraction with ϵ_{abc} only the




Figure 4: Spectral functions $\text{Im}G_E^s(t)$, $\text{Im}G_M^s(t)$, and $\text{Im}G_A(t)$ calculated with second-order chiral πN -vertices. The dashed lines correspond to the nonrelativistic approximation.

(last) c_4 -term contributes to $\text{Im}G_{E,M}^s(t)$, whereas all four c_i -terms contribute to $\text{Im}G_{A,P}(t)$. For the numerical evaluation of $\int d\Phi_3 H(\dots)$, we choose the (rounded) values $c_1 = -0.8 \text{ GeV}^{-1}$, $c_2 = 3.3 \text{ GeV}^{-1}$, $c_3 = -4.7 \text{ GeV}^{-1}$ and $c_4 = 3.4 \text{ GeV}^{-1}$ of the second-order low-energy constants c_i . Similar values are often employed in chiral NN-potentials and they are consistent with recent determinations from πN -dispersion relation analyses. With this fixed input the results for the spectral functions $\text{Im}G_E^s(t)$, $\text{Im}G_M^s(t)$ and $\text{Im}G_A(t)$ are shown in Fig. 4. One sees that these (formally) subleading corrections are roughly of similar size as the leading order terms displayed in Fig. 3. A more detailed comparison reveals that the c_4 -contribution to $\text{Im}G_E^s(t)$ is suppressed for $3m_\pi < \sqrt{t} < 5m_\pi$ and this suppression is more pronounced for $\text{Im}G_M^s(t)$. On the other hand the combined c_i -contributions to $\text{Im}G_A(t)$ exceed the leading order axial spectral function already for $\sqrt{t} > 3.7m_\pi$. The latter feature is explained by the large value of the low-energy constant c_3 . The dashed lines in Fig. 4 refer to the nonrelativistic approximation, which again leads to an overestimation by about a factor 2. In the nonrelativistic limit the following integral-representations can be derived for the electromagnetic spectral functions:

$$\text{Im}G_E^s(t) = \frac{g_A c_4}{2(8\pi)^4 f_\pi^6 \sqrt{t}} \int_{2m_\pi}^{\sqrt{t}-m_\pi} dw (w^2 - 4m_\pi^2)^{3/2} \lambda(w, t), \quad (23)$$

$$\text{Im}G_M^s(t) = \frac{g_A c_4 M}{2(4\pi)^5 f_\pi^6 t} \int_{2m_\pi}^{\sqrt{t}-m_\pi} dw (w^2 - 4m_\pi^2)^{3/2} \sqrt{\lambda(w, t)} \left[t - w^2 - m_\pi^2 - \frac{\lambda(w, t)}{3t} \right], \quad (24)$$

and for the axial spectral functions:

$$\begin{aligned} \text{Im}G_A(t) = & \frac{g_A}{(4f_\pi)^4 \pi^2 \sqrt{t}} \int_{2m_\pi}^{\sqrt{t}-m_\pi} dw \sqrt{w^2 - 4m_\pi^2} \left\{ \frac{2c_4}{3} (w^2 - 4m_\pi^2) (t - w^2 - m_\pi^2) \right. \\ & \left. + \frac{\lambda(w, t)}{t} \left[c_3 (2m_\pi^2 - w^2) - 4c_1 m_\pi^2 + \frac{c_2 + c_4}{6} (4m_\pi^2 - w^2) \right] \right\}, \end{aligned} \quad (25)$$

$$\begin{aligned}
\text{Im}G_P(t) = & \frac{g_A M^2}{64\pi^2 f_\pi^4 (t - m_\pi^2) t^{3/2}} \int_{2m_\pi}^{\sqrt{t-m_\pi}} dw \sqrt{w^2 - 4m_\pi^2} \left\{ \left[c_3(2m_\pi^2 - w^2) - 4c_1 m_\pi^2 \right. \right. \\
& + \frac{c_2 + c_4}{6} (4m_\pi^2 - w^2) \left. \right] \left[\frac{3m_\pi^2}{t} (w^2 - m_\pi^2)^2 - 3w^2 m_\pi^2 - m_\pi^4 - \lambda(w, t) \right] \\
& \left. + \frac{c_4}{6} (w^2 - 4m_\pi^2) [t(4w^2 + 5m_\pi^2) - 4t^2 - w^2 m_\pi^2 + m_\pi^4] \right\}, \tag{26}
\end{aligned}$$

where the latter expression includes also pion-pole diagrams (axial source $\rightarrow \pi \rightarrow 3\pi \rightarrow \bar{N}N$) involving the chiral $\pi\pi$ -interaction. As a good check, one can verify that the combination $\text{Im}G_A(t) + (t/4M^2)\text{Im}G_P(t)$, related to the divergence of the isovector axial-current, scales as m_π^2 . Note that the dw -integrals in eqs.(23,25,26) can be solved in terms of square-root and logarithmic functions.

3.3 Inclusion of explicit $\Delta(1232)$ -isobars

The sizeable magnitude of the low-energy constants $c_{2,3,4}$ is explained by large contributions from the $\Delta(1232)$ -resonance, which strongly couples to the πN -system. The covariant description of the $\Delta(1232)$ -isobar with spin and isospin 3/2 requires a Rarita-Schwinger spinor field Ψ_α . In this formulation the spin-3/2 propagator (vector-index β to α) takes the (common) form [10]:

$$\frac{i}{3} \frac{\gamma \cdot P + M_\Delta}{M_\Delta^2 - P^2} \left(3g_{\alpha\beta} - \gamma_\alpha \gamma_\beta - \frac{2P_\alpha P_\beta}{M_\Delta^2} + \frac{P_\alpha \gamma_\beta - \gamma_\alpha P_\beta}{M_\Delta} \right), \tag{27}$$

with P the four-momentum of the propagating $\Delta(1232)$ -isobar. In order to keep the two-loop calculations tractable, we choose minimal forms of the vertices for the coupling of an in-going pion to ΔN and $\Delta\Delta$, which read:

$$\Delta_\alpha N \pi^a(l_1): \quad -\frac{3g_A}{2\sqrt{2}f_\pi} l_1^\alpha T_a, \quad \Delta_\alpha \Delta_\beta \pi^b(l_2): \quad \frac{3g_A}{10f_\pi} g^{\alpha\beta} \gamma \cdot l_2 \gamma_5 \Theta_b. \tag{28}$$

The isospin transition operator T_a satisfies the relation $T_a T_b^\dagger = (2\delta_{ab} - i\epsilon_{abc}\tau_c)/3$, and for the isospin-3/2 operator Θ_a (a 4×4 matrix) the reduction formula $T_a \Theta_b T_c^\dagger = (5i\epsilon_{abc} - \delta_{ab}\tau_c + 4\delta_{ac}\tau_b - \delta_{bc}\tau_a)/3$ is relevant. The coupling constants in eq.(28) obey the ratios $g_{\pi N\Delta}/g_{\pi NN} = 3/\sqrt{2}$ and $g_{\pi\Delta\Delta}/g_{\pi NN} = 1/5$ as inferred from large- N_c QCD [11]. One should note that extended versions of the vertices in eq.(28) with further off-shell parameters have been proposed [10, 11], but these parameters are not well determined. Since no direct empirical information is available, the relation $g_{\pi\Delta\Delta} = g_{\pi NN}/5$ is commonly used [11].

Employing the just described formulation of vertices and propagators, we have derived the integrand-functions $H(\dots)$ for $\text{Im}G_{E,M}^s(t)$ and $\text{Im}G_A(t)$ from the diagrams with single and double virtual $\Delta(1232)$ -excitation (analogous to diagram (c) in Fig. 2). The corresponding numerical results are shown by the full lines in Fig. 5. By comparison with Fig. 4 one observes that the spectral functions get appreciably reduced by the energy-dependent Δ -propagators. The suppression factor is about 2 to 3 for $\text{Im}G_E(t)$ and $\text{Im}G_A(t)$, whereas it amounts to about 7 to 8 for $\text{Im}G_M(t)$. Of course, the physics here and in subsection 3.2 is somewhat different. The c_i -parameters represent more than the Δ -intermediate state ($c_2^{(\Delta)} = -c_3^{(\Delta)} = 2c_4^{(\Delta)} \simeq 2.9 \text{ GeV}^{-1}$) and there are partly compensating effects from single and double Δ -isobar excitation. It is also instructive to present formulas which refer to the nonrelativistic approximation. For doing that we take first the limit




Figure 5: Spectral functions $\text{Im}G_E^s(t)$, $\text{Im}G_M^s(t)$, and $\text{Im}G_A^s(t)$ calculated from diagrams with single and double propagation of Δ -isobars. The dashed lines refer to the nonrelativistic approximation.

of ΔN -degeneracy, $M_\Delta = M$, and then expand in $1/M$. This way one obtains for the electric spectral function:

$$\text{Im}G_E^s(t) = \frac{3g_A^3 t}{(4\pi)^5 f_\pi^6} \left(1 + \frac{5}{2}\right) \iint_{z^2 < 1} d\omega_1 d\omega_2 |\vec{l}_1| |\vec{l}_2| \sqrt{1-z^2} \arccos(-z), \quad (29)$$

where the factor $(1 + 5/2)$ displays the separate contributions from ΔN and $\Delta\Delta$. Likewise, one finds for the magnetic spectral function:

$$\text{Im}G_M^s(t) = \frac{g_A^3 M}{4(8\pi)^4 f_\pi^6 t^{3/2}} \int_{2m_\pi}^{\sqrt{t}-m_\pi} dw \sqrt{w^2 - 4m_\pi^2} (8m_\pi^2 - 5w^2) \lambda(w, t), \quad (30)$$

which is opposite to the term proportional to g_A^3 in eq.(21). This opposite sign and the factor $(1 + 5/2)$ can be deduced from the spin- and isospin-algebra involved in the (nonrelativistic) three-pion to nucleon coupling, which has to be spin-independent (spin-dependent) for the electric (magnetic) form factor. The dashed lines in Fig. 5 correspond to the nonrelativistic approximations written in eqs.(29,30) as well as to a more complicated formula for $\lim_{M_\Delta=M \rightarrow \infty} \text{Im}G_A^s(t)$. One can see that the proposed nonrelativistic approximation strongly overestimates the results for spectral functions with $\Delta(1232)$ -excitation based on fully relativistic kinematics. In the case of $\text{Im}G_M^s(t)$ there is even a difference in sign.

4 Phenomenological analysis and $\pi^0\gamma$ intermediate state

In this section we want find out the low-energy region, where the 3π -continua calculated in covariant baryon chiral perturbation theory could become physically relevant. For that purpose we compare with the spectral functions produces by the respective lowest-lying vector-meson resonance. For the isoscalar electromagnetic form factors $G_{E,M}^s(t)$ this is obviously the narrow ω -meson with mass $m_\omega = 783 \text{ MeV}$ and decay width $\Gamma_\omega = 8.5 \text{ MeV} = (7.6 + 0.7 + 0.2) \text{ MeV}$ [12]. In this decomposition of Γ_ω the first two entries refer to the dominant decay modes $\omega \rightarrow \pi^+\pi^0\pi^-$ and




Figure 6: Isoscalar electromagnetic spectral functions $\text{Im}G_{E,M}^s(t)$ assuming $\omega(783)$ -meson dominance.

$\omega \rightarrow \pi^0\gamma$. The reasonable assumption of $\omega(783)$ -meson dominance in the region $3m_\pi < \sqrt{t} < 7m_\pi$ leads to the following complex-valued form factors:

$$G_{E,M}^s(t) = \frac{(0.5, 0.44)m_\omega^2}{m_\omega^2 - t - im_\omega\Gamma_\omega(t)}, \quad (31)$$

with $\Gamma_\omega(t)$ an energy-dependent ω -meson decay width. Modelling the two dominant decay modes by appropriate contact-couplings, one gets:

$$\Gamma_\omega(t) = \frac{h^2}{m_\omega t} \int_{2m_\pi}^{\sqrt{t}-m_\pi} dw [(w^2 - 4m_\pi^2)\lambda(w, t)]^{3/2} + \frac{h'^2}{m_\omega t} (t - m_\pi^2)^3, \quad (32)$$

with the parameters $h = 2.72 \text{ GeV}^{-3}$ and $h' = 0.040 \text{ GeV}^{-1}$ adjusted to the partial decay widths.

The resulting imaginary parts $\text{Im}G_{E,M}^s(t)$ are shown in Fig. 6. The resonance curves for the electric and magnetic form factor are almost equal, due to similar normalizations $0.5 \simeq 0.44$. One sees that in the region $3m_\pi < \sqrt{t} < 5m_\pi$ the 3π -only contributions from the $\omega(783)$ -resonance fall below the (combined) chiral 3π -continua, whereas the additional $\pi^0\gamma$ -mode introduces appreciable strength in the threshold region. In view of this striking effect, one is compelled to compute the radiative correction to the isoscalar electromagnetic spectral functions coming from the $\pi^0\gamma$ -intermediate state. The pertinent S-matrix for $\pi^0 \rightarrow \gamma\gamma$ reads: $(-i\alpha_{\text{em}}/\pi f_\pi)\epsilon_{\mu\nu\alpha\beta}k_1^\alpha k_2^\beta$, where (k_1, μ) and (k_2, ν) pertain to out-going photons. The one-loop calculation of both diagrams $\gamma^* \rightarrow \pi^0\gamma \rightarrow \bar{N}N$ requires only one angular integration $(m_\pi^2 - t)/(32\pi t) \int_{-1}^1 dx$, such that the $\pi^0\gamma$ -contribution to the isoscalar electromagnetic spectral functions can be given in analytical form:

$$\text{Im}G_E^s(t) = \frac{\alpha_{\text{em}}g_A(t - m_\pi^2)^2}{(4\pi f_\pi)^2(4M^2 - t)} \left\{ -\frac{1}{4} + \frac{\kappa_v}{3} \left(\frac{t - m_\pi^2}{16M^2} - 1 + \frac{m_\pi^2}{4t} \right) + \frac{(1 + \kappa_v)M^2}{\sqrt{t}(4M^2 - t)} \arccos \frac{\sqrt{t}}{2M} \right\}, \quad (33)$$

$$\text{Im}G_M^s(t) = \frac{\alpha_{\text{em}}g_A M^2(t - m_\pi^2)^2}{(4\pi f_\pi)^2(4M^2 - t)t} \left\{ \frac{1}{2} + \frac{\kappa_v}{3} \left(2 - \frac{t + 2m_\pi^2}{8M^2} + \frac{m_\pi^2}{t} \right) + \frac{4M^2 - (2 + \kappa_v)t}{2\sqrt{t}(4M^2 - t)} \arccos \frac{\sqrt{t}}{2M} \right\}, \quad (34)$$




Figure 7: Contributions to the isoscalar electromagnetic spectral functions $\text{Im}G_{E,M}^s(t)$ from the $\pi^0\gamma$ -intermediate state.

with $\kappa_v = \kappa_p - \kappa_n = 3.706$ the (large) isovector anomalous magnetic moment and $\alpha_{\text{em}} = 1/137$. Note that one averages here over proton and neutron form factors, while the $\pi^0 N$ -coupling $\sim \tau_3$ introduces an opposite sign for the magnetic moment term. The curves resulting from the expressions in eqs.(33,34), with threshold behavior $\text{Im}G_{E,M}^s(t) \sim (t - m_\pi^2)^2$, are drawn in Fig. 7. One observes that in the region $3m_\pi < \sqrt{t} < 4m_\pi$ the radiative corrections due to the $\pi^0\gamma$ -intermediate state exceed the chiral 3π -continua. This behavior is explained kinematically by the fast decrease of the 3π -phase space towards the threshold $\sqrt{t} = 3m_\pi$, while the $\pi^0\gamma$ -phase space remains open down to $\sqrt{t} = m_\pi$.

In analogy to eq.(31) the nucleon axial form factor $G_A(t)$ dominated by the a_1 -resonance reads:

$$G_A(t) = \frac{g_A m_{a_1}^2}{m_{a_1}^2 - t - im_{a_1} \Gamma_{a_1}(t)}, \quad (35)$$

with the axial-vector coupling constant $g_A = 1.27$ [12]. The mass and width of the broad a_1 -meson are still under debate, due to conflicting results from different experiments. A very recent partial wave analysis of diffractive dissociation data ($\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$) by the COMPASS collaboration [13] finds the (central) values $m_{a_1} = 1.3 \text{ GeV}$ and $\Gamma_{a_1} = 0.38 \text{ GeV}$. On the other hand the values extracted from τ -lepton decays in ref.[14] are $m_{a_1} = 1.2 \text{ GeV}$ and $\Gamma_{a_1} = 0.48 \text{ GeV}$, while a later reanalysis in ref. [15] gave a somewhat lower a_1 -mass of $m_{a_1} = 1.12 \text{ GeV}$.

The full lines in Fig. 8 show the axial spectral function $\text{Im}G_A(t)$ using the specific form [15] of $\Gamma_{a_1}(t)$, which follows from integrating (interfering) Breit-Wigner functions for the $\rho(770)$ -resonance over the 3π -phase space. The dashed lines were obtained with the phenomenological parametrization of $\Gamma_{a_1}(t)$ from ref. [16], which describes separately the regions below and above the $\rho\pi$ -threshold $t = (m_\rho + m_\pi)^2$. The light and dark pair of curves refer to the parameter sets $(m_{a_1}, \Gamma_{a_1}) = (1.3, 0.38) \text{ GeV}$ [13] and $(m_{a_1}, \Gamma_{a_1}) = (1.2, 0.48) \text{ GeV}$ [14], which are clearly distinguished by their shifted peaks. By comparison with the full (green) lines in Figs. 3 and 4 one can recognize an energy window near threshold, $3m_\pi < \sqrt{t} < 5m_\pi$, in which the chiral 3π -continua do prevail. However, such tiny contributions to the axial spectral function are presumably irrelevant for physical observables.




Figure 8: Axial spectral function $\text{Im}G_A(t)$ assuming a_1 -meson dominance. The two sets of (light and dark green) curves correspond to masses and widths of $(m_{a_1}, \Gamma_{a_1}) = (1.3, 0.38)$ GeV [13] and $(m_{a_1}, \Gamma_{a_1}) = (1.2, 0.48)$ GeV [14].

5 Summary and conclusions

In this work we have studied the imaginary parts of the isoscalar electromagnetic and isovector axial form factors of the nucleon close to the 3π -threshold. The contributions to $\text{Im}G_{E,M}^s(t)$ and $\text{Im}G_A(t)$ arising from chiral πN -vertices at leading and next-to-leading order, as well as pion-induced $\Delta(1232)$ -excitations have been calculated and compared with each other. It was found that the heavy baryon approach overestimates the chiral 3π -continua. From a phenomenological analysis, that included the narrow $\omega(783)$ -resonance or the broad $a_1(1260)$ -resonance, one could recognize small windows near threshold, where chiral 3π -dynamics prevails. However, for $\text{Im}G_{E,M}^s(t)$ the radiative correction provided by the $\pi^0\gamma$ -intermediate state becomes actually more relevant in the region close to threshold. Altogether, one may conclude that the chiral 3π -continua for nucleon form factors are too weak to influence physical observables in a significant way.

References

- [1] P. Mergell, U. G. Meißner and D. Drechsel, *Dispersion theoretical analysis of the nucleon electromagnetic form-factors*, *Nucl. Phys.* **A596** (1996) 367–396, [[hep-ph/9506375](#)].
- [2] I. T. Lorenz, U.-G. Meißner, H. W. Hammer and Y. B. Dong, *Theoretical Constraints and Systematic Effects in the Determination of the Proton Form Factors*, *Phys. Rev.* **D91** (2015) 014023, [[1411.1704](#)].
- [3] M. Hoferichter, J. Ruiz de Elvira, B. Kubis and U.-G. Meißner, *Roy-Steiner-equation analysis of pion-nucleon scattering*, *Phys. Rept.* **625** (2016) 1–88, [[1510.06039](#)].

- [4] M. Hoferichter, B. Kubis, J. Ruiz de Elvira, H. W. Hammer and U. G. Meißner, *On the $\pi\pi$ continuum in the nucleon form factors and the proton radius puzzle*, *Eur. Phys. J.* **A52** (2016) 331, [[1609.06722](#)].
- [5] N. Kaiser, *Spectral functions of isoscalar scalar and isovector electromagnetic form-factors of the nucleon at two loop order*, *Phys. Rev.* **C68** (2003) 025202, [[nucl-th/0302072](#)], and references therein.
- [6] I. T. Lorenz, H. W. Hammer and U.-G. Meißner, *The size of the proton - closing in on the radius puzzle*, *Eur. Phys. J.* **A48** (2012) 151, [[1205.6628](#)].
- [7] V. Bernard, N. Kaiser and U.-G. Meißner, *Nucleon electroweak form-factors: Analysis of their spectral functions*, *Nucl. Phys.* **A611** (1996) 429–441, [[hep-ph/9607428](#)].
- [8] N. Kaiser, *Induced pseudoscalar form-factor of the nucleon at two loop order in chiral perturbation theory*, *Phys. Rev.* **C67** (2003) 027002, [[nucl-th/0301034](#)].
- [9] J. Bijnens, *Chiral perturbation theory and anomalous processes*, *Int. J. Mod. Phys.* **A8** (1993) 3045–3105.
- [10] V. Bernard, N. Kaiser and U.-G. Meißner, *Chiral dynamics in nucleons and nuclei*, *Int. J. Mod. Phys.* **E4** (1995) 193–346, [[hep-ph/9501384](#)], appendices A and E.
- [11] D. Siemens, V. Bernard, E. Epelbaum, H. Krebs and U.-G. Meißner, *The reaction $\pi N \rightarrow \pi\pi N$ in chiral effective field theory with explicit $\Delta(1232)$ degrees of freedom*, *Phys. Rev.* **C89** (2014) 065211, [[1403.2510](#)].
- [12] PARTICLE DATA GROUP collaboration, M. Tanabashi et al., *Review of Particle Physics*, *Phys. Rev.* **D98** (2018) 030001.
- [13] COMPASS collaboration, R. Akhunzyanov et al., *Light isovector resonances in $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$ at 190 GeV/c*, [1802.05913](#); J. M. Friedrich, private communication.
- [14] D. Gomez Dumm, A. Pich and J. Portoles, *$\tau \rightarrow \pi\pi\nu_\tau$ decays in the resonance effective theory*, *Phys. Rev.* **D69** (2004) 073002, [[hep-ph/0312183](#)].
- [15] D. G. Dumm, P. Roig, A. Pich and J. Portoles, *$\tau \rightarrow \pi\pi\nu_\tau$ decays and the $a_1(1260)$ off-shell width revisited*, *Phys. Lett.* **B685** (2010) 158–164, [[0911.4436](#)].
- [16] J. H. Kühn and A. Santamaria, *Tau decays to pions*, *Z. Phys.* **C48** (1990) 445–452.