

Baryon σ terms in SU(3) BChPT $\times 1/N_c$

Ishara P. Fernando*

Hampton University & Thomas Nelson Community College, Hampton, VA, USA

E-mail: ishara@jlab.org

José L. Goity

Hampton University, Hampton, VA & Jefferson Lab, Newport News, VA, USA

E-mail: goity@jlab.org

ChPT and the $1/N_c$ expansion provide systematic frameworks for the strong interactions at low energy. A combined framework of both expansions has been developed and applied for baryons with three light-quark-flavors. The small scale expansion of the combined approach is identified as the ξ -expansion, in which the power counting of the expansions is linked according to $\mathcal{O}(p) = \mathcal{O}(1/N_c) = \mathcal{O}(\xi)$. The physical baryon masses as well as lattice QCD baryon masses for different quark mass masses are analyzed to $\mathcal{O}(\xi^3)$ in that framework. σ terms are addressed using the Feynman Hellmann theorem. For the nucleon, a useful connection between the deviation of the Gell-Mann-Okubo relation and the σ term σ_{8N} associated with the scalar density $\bar{u}u + \bar{d}d - 2\bar{s}s$ is identified. In particular, the deviation from the tree level relation $\sigma_{8N} = \frac{1}{3}(2m_N - m_\Sigma - m_\Xi)$, which gives rise to the so called σ -term puzzle, is studied in the ξ -expansion. A large correction non-analytic in ξ results for that relation, making plausible the resolution of the puzzle. Issues with the determination of the strangeness σ terms are discussed, emphasizing the need for lattice calculations at smaller m_s for better understanding the range of validity of the effective theory. The analysis presented here leads to $\sigma_{\pi N} = 69(10)$ MeV and $\sigma_{\pi \Delta} = 60(10)$ MeV.

The 9th International workshop on Chiral Dynamics

17-21 September 2018

Durham, NC, USA

*Speaker.

1. Introduction

Combining $BChPT$ and the $1/N_c$ expansion [1, 2, 3, 4] in baryons with three light quark flavors leads to an improvement in the description of baryon masses and currents [4, 5, 6, 7] to one-loop. A link between the chiral and the $1/N_c$ expansions is necessary in order to establish an unambiguous power counting: the counting where $\mathcal{O}(p) = \mathcal{O}(1/N_c) = \mathcal{O}(\xi)$, closely related to the small scale expansion [8, 9], is in practice the most effective one. In this framework, the effective Lagrangians to $\mathcal{O}(\xi^3)$ can be found in Ref. [5, 10]. The chiral Lagrangian relevant to the discussion of masses up to $\mathcal{O}(\xi^3)$ and including electromagnetic contributions is given by [5, 10]:

$$\begin{aligned} \mathcal{L}_{\mathbf{B}}^{\text{mass}} = & \mathbf{B}^\dagger \left(iD_0 + \hat{g}_A u^{ia} G^{ia} - \frac{C_{\text{HF}}}{N_c} \hat{S}^2 + \frac{c_1}{2\Lambda} \hat{\chi}_+ + \frac{c_2}{\Lambda} \chi_+^0 + \frac{c_3}{N_c \Lambda^3} \hat{\chi}_+^2 \right. \\ & \left. + \frac{h_1 \Lambda}{N_c^3} \hat{S}^4 + \frac{h_2}{N_c^2 \Lambda} \hat{\chi}_+ \hat{S}^2 + \frac{h_3}{N_c \Lambda} \chi_+^0 \hat{S}^2 + \frac{h_4}{N_c \Lambda} \chi_+^a \{S^i, G^{ia}\} + \alpha \hat{Q} + \beta \hat{Q}^2 \right) \mathbf{B}, \end{aligned} \quad (1.1)$$

where $\hat{g}_A$ is the axial coupling constant identified at LO with $\frac{6}{5}g_A^N$, where $g_A^N = 1.2724(23)$. The low energy constants (LECs) C_{HF} , c_{1-4} , h_{1-4} and α, β can be fixed [5] by fitting the baryon masses to the experimental data and to results from lattice QCD (LQCD) calculations [11] at varying quark masses. Using standard notation, $\hat{\chi}_+ = \tilde{\chi}_+ + \chi_+^0$, where $\chi_+^0 = \frac{1}{3}\text{Tr}\chi_+$, provide the quark mass dependent terms. $\hat{Q}$ is the electric charge operator. The electromagnetic contribution to the $p-n$ mass difference is $\alpha + \beta$, whereas the electromagnetic contribution to the Gell-Mann-Okubo (GMO) formula is $-\frac{4}{3}\beta$. Up to $\mathcal{O}(\xi^3)$ the baryon mass formula, neglecting isospin breaking, reads:

$$\begin{aligned} m_{\mathbf{B}} = & M_0 + \frac{C_{\text{HF}}}{N_c} \hat{S}^2 - \frac{c_1}{\Lambda} 2B_0 (\sqrt{3}m_8 \hat{Y} + N_c m_0) - \frac{c_2}{\Lambda} 4B_0 m_0 \\ & - \frac{h_1 \Lambda}{N_c^3} \hat{S}^4 - \frac{h_2}{N_c^2 \Lambda} 4B_0 (\sqrt{3}m_8 \hat{Y} + N_c m_0) \hat{S}^2 - \frac{h_3}{N_c \Lambda} 4B_0 m_0 \hat{S}^2 \\ & - \frac{h_4}{N_c \Lambda} \frac{4B_0 m_8}{\sqrt{3}} \left(3\hat{I}^2 - \hat{S}^2 - \frac{1}{12} N_c (N_c + 6) + \frac{1}{2} (N_c + 2) \hat{Y} - \frac{3}{4} \hat{Y}^2 \right) + \delta m_B^{\text{loop}}, \end{aligned} \quad (1.2)$$

where M_0 is the $\mathcal{O}(N_c)$ spin-flavor singlet piece of the baryon masses, $\hat{S}$, $\hat{I}$ and $\hat{Y}$ are respectively the baryon spin, isospin and hypercharge operators, the term proportional to C_{HF} gives the LO hyperfine mass splittings between different spin baryons, and m_0 and m_8 are the singlet and octet components of the quark masses. δm_B^{loop} gives the one-loop contributions $\mathcal{O}(\xi^2)$ and $\mathcal{O}(\xi^3)$. It is straightforward to generalize 1.3 to include isospin breaking. In the following the definitions are used: $m_0 = \frac{1}{3}(2\hat{m} + m_s)$, $m_3 = m_u - m_d$ and $m_8 = \frac{1}{\sqrt{3}}(\hat{m} - m_s)$, where $\hat{m} = \frac{1}{2}(m_u + m_d)$. More details on the self energy one-loop corrections obtained in $BChPT \times 1/N_c$ can be found in these proceedings [7].

2. σ -terms

The matrix elements of scalar quark densities are of high interest. At zero momentum they are related via the Feynman-Hellmann theorem to the slope of the hadron mass with

respect to the corresponding quark mass,¹

$$\sigma_{f\mathbf{B}}(m_f) = m_f \frac{\partial}{\partial m_f} m_{\mathbf{B}} = \frac{m_f}{2m_{\mathbf{B}}} \langle \mathbf{B} | \bar{q}_f q_f | \mathbf{B} \rangle, \quad (2.1)$$

where m_f is the mass of the f quark flavor ($f = u, d, s$), the state $|\mathbf{B}\rangle$ is the physical state for that quark mass and normalized according to $\langle \mathbf{B}' | \mathbf{B} \rangle = (2\pi)^3 2m_B \delta^3(\vec{p}' - \vec{p})$, and $\sigma_{f\mathbf{B}}$ is the corresponding σ term. σ terms for combinations of quark masses such as m_0 , m_3 and m_8 are defined in the same way. Empirical access to σ terms is difficult in the case of baryons, being only possible for $\sigma_{\pi N} = \sigma_{(u+d)N}(\hat{m})$ via analysis of πN scattering. In the case of other σ terms it is clear that the necessary information will have to come from LQCD calculations, where tracing the baryon mass dependency with respect to quark masses is becoming increasingly accurate. The actual contribution of a given quark flavor mass to the mass of the hadron, keeping the rest of the quark flavor masses fixed, is then given by:

$$\Delta m_{\mathbf{B}}^f(m_f) = \int_0^{m_f} \frac{1}{\mu} \sigma_{f\mathbf{B}}(\mu) d\mu, \quad (2.2)$$

which in the limit of small m_f coincides with the σ term.

In this note, the focus is on the determination of $\sigma_{\pi N}$ using the Feynman-Hellmann theorem and results for baryon masses in $SU(3)$, as presented in Ref. [10], with additional brief discussions of σ terms of Δ and hyperons, and the issue of the quark mass dependence of σ terms, namely the range in m_q where the effective theory may be trusted in their description.

2.1 $\sigma_{\pi N}$

The determination of $\sigma_{\pi N}$ has a long history spanning many decades. Its extraction from the analyses of πN scattering has given values that range from 45 MeV [12, 13, 14] to 64 MeV [15, 16, 17, 18], with the larger values being from more recent analyses where their increment with respect to the old ones is understood to be a consequence of a change in the input πN scattering lengths. From a practical use point of view, $\sigma_{\pi N}$ has become very important in the studies of dark matter searches [19] in the scenarios where dark matter has scalar couplings to quarks.

$\sigma_{\pi N}$ can be expressed by the combination of σ terms:

$$\sigma_{\pi N} = \hat{\sigma} + 2 \frac{\hat{m}}{m_s} \sigma_{sN}, \quad (2.3)$$

where $\hat{\sigma} = \sqrt{3} \frac{\hat{m}}{m_8} \sigma_{8N}$. To LO in quark masses σ_{8N} is given by a combination of octet baryon masses, namely:

$$\begin{aligned} \sigma_{8N} &= \frac{1}{3} (2m_N - m_{\Sigma} - m_{\Xi}), \\ &= \frac{1}{9} \left(\frac{5N_c - 3}{2} m_N - (2N_c - 3) m_{\Sigma} - \frac{N_c + 3}{2} m_{\Xi} \right) \quad \text{for general } N_c, \end{aligned} \quad (2.4)$$

¹Although obvious, σ terms, being observable quantities, are independent of the renormalization scheme used in QCD. The expression 2.1 normally used is valid in a mass independent scheme such as $\overline{MS}$.

which leads to $\hat{\sigma} \sim 25$ MeV. Since the contribution of the term proportional to σ_{sN} , being OZI suppressed, should be expected to be small, at this lowest order in the quark masses there is a puzzle between the empirically obtained values of $\sigma_{\pi N}$ and the relation $\sigma_{\pi N} \sim \hat{\sigma}$. Either the latter is badly broken, and/or the relation 2.4 has large corrections. It will be shown that the latter is the case. It is argued that the puzzle is further emphasized by the observation that the Gell-Mann-Okubo relation ² receives small deviations, and so it would be difficult to understand why 2.4 should receive large corrections [20]. Following Ref. [5], and based on the $1/N_c$ expansion one finds that the corrections to the GMO relation are suppressed by a factor $1/N_c$ at large N_c , while the corrections to the mass relation generalized in N_c as shown in 2.4 are $\mathcal{O}(N_c)$. The deviation from the GMO relation, Δ_{GMO} , in the calculation to one-loop is independent of the NLO LECs and given solely by non-analytic finite contributions, which depend on F_π , C_{HF} and the GB masses. The same is the case for the deviations from 2.4, denoted here by $\Delta\sigma_{8N}$. Performing the analysis at generic N_c , one finds that Δ_{GMO} is indeed $\mathcal{O}(1/N_c)$ at large N_c , and in terms of the ξ power counting it is $\mathcal{O}(\xi^4)$ (an extra factor $1/N_c$ over the nominal $\mathcal{O}(\xi^3)$ of the loop corrections), while $\Delta\sigma_{8N}$ is $\mathcal{O}(\xi^2)$ with a pre-factor N_c ³. Thus they have entirely different behaviors, and on these grounds it is entirely plausible that $\Delta\sigma_{8N}$ can be as large as the resolution to the puzzle requires. It is also observed that in the physical case the ratio $\Delta\sigma_{8N}/\Delta_{GMO} \sim -14$, which is independent of F_π and C_{HF} , has only a small dependency on the LEC C_{HF} , and thus it is determined almost entirely in terms of the GB masses. Since the large corrections $\Delta\sigma_{8N}$ are due to the rather large value of m_s , it is important to check how σ_{8N} is as a function of M_K . This is shown in Figure 1, which clearly illustrates the following point: the non-analytic contributions to σ_{8N} are not large (compare σ_{8N} with the tree contribution $\sigma_N^{tree}(\mu = m_\rho)$). The corrections to the mass combination denoted here by σ_8^{rel} are very small, but they result from two large contributions, the $\sigma_N^{tree}(\mu = m_\rho)$ and a non-analytic one that largely cancels it. Thus, a large entirely non-analytic correction $\Delta\sigma_{8N}$ is the result. The figure also shows the behavior of σ_s , which has a large relative variation in the displayed interval; its size is nonetheless natural, leading in Eq. 2.3 to a small contribution by that term of the order of a few MeV. As discussed later, the σ_s terms are outside of the range of validity of the effective theory for the physical m_s values. In order to check that the effective theory is giving reasonable results, one can make use of the calculated Δ_{GMO} and check with its actual value: as shown below, this works very well; even more, the octet baryons in the loop contribute 43% of Δ_{GMO} , thus the contribution by the decuplet is crucial. One can also infer from Δ_{GMO} a value for the LO axial coupling : it is about 20% smaller than the physical one, in line with that obtained in the analysis of axial couplings [5, 7]. If one only considers the contributions by the octet baryons, which is itself $\mathcal{O}(1/N_c)$, in order to obtain the physical Δ_{GMO} the needed must be larger, conflicting with the analysis of the axial couplings [5, 7].

At this point, the effective theory can determine σ_{8N} from 2.4 and the calculated $\Delta\sigma_{8N}$. To determine $\sigma_{\pi N}$ one needs further information on the baryon masses. That information

²The GMO relation is defined by the mass combination: $3m_\Lambda + m_\Sigma - 2(m_n + m_\Xi)$, valid for all N_c .

³Note that $\sigma_{8N} = \mathcal{O}(N_c)$, while $\sigma_{sN} = \mathcal{O}(N_c^0)$

is provided by LQCD, as for instance in the analysis of octet and decuplet masses of Ref. [11], where m_s is kept approximately fixed and $\hat{m}$ is varied. A fit to the masses allows for a direct extraction of $\sigma_{\pi N}$ and also an estimate, albeit with large error, of σ_{sN} . As discussed below, the end result is that the relation $\sigma_{\pi N} \simeq \hat{\sigma}$ is approximately well satisfied. The most direct determination of σ_{8N} is thus carried out making use of the ratio $\Delta\sigma_{8N}/\Delta_{GMO}$ using a value of C_{HF} as obtained in the fit to octet and decuplet masses and correcting Δ_{GMO} by EM and $m_u - m_d$ isospin breaking effects (see Ref. [5] for details), giving $\hat{\sigma} \simeq 70$ MeV, which leads to a value for $\sigma_{\pi N}$ which is at the upper range of values obtained in previous studies.

The question is up to what extent is the determination of σ_{8N} discussed here realistic. It is clear, as emphasized below, that the σ terms associated with the strange quark at its physical mass cannot be described well by the effective theory. This implies also that the description of the hyperon masses in the physical case are somewhat outside the range where one can trust the effective theory. Thus, both the parameter free calculations of Δ_{GMO} and $\Delta\sigma_{8N}$ may not be as accurate as one would wish. There is little doubt that the analysis presented here would work reliably for a smaller m_s , for $M_K < 300 - 400$ MeV or so (see Fig. 1). The only way this can be established is via LQCD calculations with lighter m_s than the ones presently available. Such calculations would indeed provide important additional insights on the σ terms and more in general on the effectiveness of the different versions of BChPT, in particular the present one, which would be greatly welcomed.




Figure 1: σ terms as a function of M_K from baryon masses to $\mathcal{O}(\xi^3)$. σ_{8N} full red, $\sigma_{8N}^{\text{tree}}(\mu = m_\rho)$ short-dash red, σ_{8N} from the mass relation 2.4 dashed red, $10 \times \sigma_{sN}$ purple, $10 \times \hat{\sigma}_N$ green, and $10 \times \sigma_{\pi N}$ blue. Based on the analysis of Ref. [5].

2.2 Other σ terms

A similar analysis to the case of the Nucleon can be carried out for the Δ . In that case there is the following LO relation for σ_Δ^8 :

$$\sigma_{8\Delta} = \frac{N_c}{3} (m_\Delta - m_{\Sigma^*}) - \frac{5(N_c - 3)}{12} (m_\Lambda - m_\Sigma), \quad (2.5)$$

whose deviations at NLO are again calculable as in the case of the Nucleon. Since in the large N_c limit the Δ and Nucleon become degenerate, their respective σ terms must also become identical up to terms sub-leading in $1/N_c$. That regime is however reached at very large N_c (fixed m_q) for the contributions non-analytic in $\sqrt{m_q} \times N_c$.

σ terms satisfy the same tree level relations as baryon masses do. Indeed, the GMO, Equal spacing and the Gürsey-Radicati (if the LEC h_3 is neglected) mass relations, satisfied by tree contributions up to $\mathcal{O}(\xi^3)$, are automatically satisfied by the corresponding σ terms. Since the non-analytic corrections to those relations are all $1/N_c$ suppressed, the corresponding σ term relations have small deviations. There are further tree level relations satisfied by σ terms corresponding to different quark masses, in particular relating the σ terms corresponding to m_s with the $\hat{m}$ ones. The corrections to those are not $1/N_c$ suppressed and thus they receive large non-analytic corrections. As shown later, the $\sigma_{\mathbf{B}}^s$ terms show significant curvature starting at $M_K \sim 250$ MeV, indicating the range where the effective theory can be trusted with their calculation. Those additional σ terms may be of general interest in LQCD calculations and the corresponding tests of the effective theory they can provide.

3. Results using LQCD inputs

In the analysis of Ref. [10], both physical and LQCD baryon masses are considered. The LQCD baryon masses have been obtained for approximately fixed M_K , varying $m_u = m_d$ in a range from the physical limit up to $M_\pi \sim 300$ MeV [11]. Three different fits were performed, shown in the Table (1), which contains some additional results to those given in [10]. The ratio $\hat{g}_A/F_\pi$ is also a fitting parameter for the first two fits and it is consistent with the value extracted from Δ_{GMO} and also the one obtained from the analysis of axial couplings [5, 7]. The value of C_{HF} is determined most accurately by the physical $\Delta - N$ mass splitting; its value obtained solely from the LQCD results is significantly different and indication that the LQCD results do not determine accurately the hyperfine mass splittings, extrapolating to too small of a value at the physical limit. For the physical case isospin breaking was taken into account, which allows to fix the EM coefficients α and β . For the present analysis, the importance of that correction is its effect on Δ_{GMO} , whose value without EM is that obtained with the physical masses plus $\frac{4}{3}\beta$, a non-negligible effect of almost 3 MeV increase.

It is important to stress that the resulting LECs and the respective errors are natural have natural size. More accurate LQCD results and, as emphasized later, with smaller m_s would help determine how reliable is the effective theory is. More details on extracting sigma terms for the Nucleon can be found in [10]. The fit gives an estimate for σ_{sN} , which as discussed below is not credible, and should only be taken as an estimate of its magnitude for the purpose of determining $\sigma_{\pi N}$. As expected the results for the Δ 's σ terms are very similar to those of the nucleon (they also have a small imaginary part due to the width of the Δ). A summary of the present status of $\sigma_{\pi N}$ determinations is displayed in Fig. 2.

	$\frac{\tilde{g}_A}{F_\pi}$	$\frac{M_0}{N_c}$	C_{HF}	c_1	c_2	h_2	h_3	h_4	α	β
Fit	MeV $^{-1}$	MeV	MeV						MeV	MeV
1	0.0126(2)	364(1)	166(23)	-1.48(4)	0	0	0.67(9)	0.56(2)	-1.63(24)	2.16(22)
2	0.0126(3)	213(1)	179(20)	-1.49(4)	-1.02(5)	-0.018(20)	0.69(7)	0.56(2)	-1.62(24)	2.14(22)
3	0.0126*	262(30)	147(52)	-1.55(3)	-0.67(8)	0	0.64(3)	0.63(3)	-1.63*	2.14*
	$\Delta_{GMO}^{\text{phys}}$	σ_{8N}	$\Delta\sigma_{8N}$	$\hat{\sigma}_N$	$\sigma_{\pi N}$	σ_s	$\sigma_{8\Delta}$	$\Delta\sigma_{8\Delta}$	$\hat{\sigma}_\Delta$	
	MeV	MeV	MeV	MeV	MeV	MeV	MeV	MeV	MeV	
1	25.6(1.1)	-583(24)	-382(13)	70(3)(6)	—	—	-496(46)	-348(16)	59(5)(6)	
2	25.5(1.5)	-582(55)	-381(20)	70(7)(6)	69(8)(6)	-3(32)	-511(52)	-352(22)	60(10)(6)	
3	25.8*	-615(80)	-384(2)	74(1)(6)	65(15)(6)	-121(15)	-469(26)	350(27)	56(4)(6)	

Table 1: Results of fits to baryon masses [10]. Fit 1 uses only the physical octet and decuplet masses, Fit 2 uses the physical and the LQCD masses from Ref. [11] with $M_\pi \lesssim 300$ MeV, and Fit 3 uses only those LQCD masses and imposes the value of $\Delta_{GMO}^{\text{phys}}$ determined by the physical masses (corrected in the calculation by the isospin breaking effects). The renormalization scale μ and the scale Λ are taken to be equal to m_ρ . * indicates an input. A theoretical error of 6 MeV is estimated for $\hat{\sigma}$ and $\sigma_{\pi N}$.




Figure 2: From Ref. [10]; Left panel: summary of the determinations of $\sigma_{\pi N}$ from πN scattering (blue), from LQCD (red), and from this work showing the combined fit and theoretical error. Right panel: N and Δ masses from Fit 2 of Table 1: physical and LQCD masses from [21]. The squares are the results from the fit and the error bands correspond to 68% confidence interval. Note: The references given in the left panel are corresponding to the references in Ref. [10]

3.1 Dependencies on quark masses

For N and Δ the dependency of their masses on $\hat{m}$ is quite smooth up to $M_\pi \sim 300$ MeV (Fig. 2). In the case of the hyperons the dependency is less smooth the larger the magnitude of the strangeness (Fig. 3). The first indication of significant curvature appears in the σ terms as the corresponding quark mass reaches a value of about 80 MeV, or about 300 MeV for the corresponding GB masses, as illustrated by Figs. 3. This manifests itself in curvature of the baryon masses with respect to quark masses but much less pronounced,

consequence of Eq. 2.2.




Figure 3: Nucleon sigma terms with respect to quark masses $\hat{m}$ (left panel), m_s (right panel)

One can therefore estimate the range of quark masses for which the effective theory can describe baryon masses. For hadrons with a single heavy quark one can use results from HQET to determine the hadron mass as a function of the heavy quark mass [22], for which there would be a corresponding σ term. Provided a definition of the heavy quark mass, the corresponding σ term will be, up to additive corrections determined by the scale of QCD, roughly proportional to the heavy quark mass with a slope close to unity. In general, the slope is expected to scale roughly as proportional to the number of heavy quarks, and thus one can use this to give a rough estimate for the limit where the effective low energy theory ceases to describe a σ term. For small quark masses the slope of the σ term is much larger than it would be for the corresponding quark having a very large mass. The behavior of the σ terms shown in Fig. 3 illustrate the natural tendency to a reduced slope as the quark mass increases. One could therefore use the criterion that when the slope calculated in the effective low energy theory reaches a value close to the one corresponding to the large quark mass limit, the theory cannot further be trusted, representing this also the onset of its failure for describing the hadron mass itself. The analysis shown here indicates that this occurs for the relevant GB masses above 300 MeV or so. For this reason it would be very useful to have LQCD results where m_s is taken to be smaller than in present calculations, in order to assess more accurately the issue.

4. Summary

The determination of σ terms through the Feynman-Hellmann theorem has its challenges. In principle a good knowledge of baryon masses for varying quark masses would be sufficient, but that knowledge as obtained from LQCD results is still not accurate enough to deliver values for $\sigma_{\pi N}$ with a precision near that obtained from the analysis of πN scattering. Another approach using $BChPT \times 1/N_c$ in $SU(3)$ and its predictions for Δ_{GMO} and $\Delta\sigma_{8N}$ as described in this note is potentially affected by the fact that m_s is too large for the result to be considered accurate. It is however interesting that an extraction of $\sigma_{\pi N}$ using that approach and the LQCD results agree very well. A result for $\sigma_{\pi N} = 69(10)$ MeV




Figure 4: σ terms as function of quark masses. In the left panels m_s is kept fixed, and in the right panels $\hat{m}$ is kept fixed.

results from those analyses, consistent with the larger values obtained from πN scattering. It should be emphasized that a similar analysis using ordinary $BChPT$ with only the octet baryons completely fails in that respect. We also learn that the description of strangeness σ terms fails for the physical value of m_s , and thus, one would need LQCD results with reduced values of m_s to understand more precisely the range where effective theories can describe them: it looks like the for the effective theory to be able to reliably describe σ terms in $SU(3)$ would require $M_K \leq 350$ MeV.

5. Acknowledgements

The authors would like to thank José Manuel Alarcón for earlier collaboration and useful discussions. This work was supported by DOE Contract No. DE-AC05-06OR23177 under which JSA operates the Thomas Jefferson National Accelerator Facility, and by the National Science Foundation through grants PHY-1307413 and PHY-1613951.

References

- [1] E. E. Jenkins, Phys. Rev. D53 (1996) 2625–2644. [arXiv:hep-ph/9509433](https://arxiv.org/abs/hep-ph/9509433).

- [2] R. Flores-Mendieta, E. E. Jenkins, A. V. Manohar, Phys. Rev. D58 (1998) 094028. [arXiv:hep-ph/9805416](#).
- [3] R. Flores-Mendieta, M. A. Hernandez-Ruiz, C. P. Hofmann, Phys. Rev. D86 (2012) 094041. [arXiv:1210.8445](#).
- [4] A. Calle Cordon, J. L. Goity, Baryon Masses and Axial Couplings in the Combined $1/N_c$ and Chiral Expansions, Phys. Rev. D87 (1) (2013) 016019. [arXiv:1210.2364](#), [doi:10.1103/PhysRevD.87.016019](#).
- [5] I. P. Fernando, J. L. Goity, Baryon chiral perturbation theory combined with the $1/N_c$ expansion in $SU(3)$: Framework, Phys. Rev. D97 (5) (2018) 054010. [arXiv:1712.01672](#), [doi:10.1103/PhysRevD.97.054010](#).
- [6] R. Flores-Mendieta, J. L. Goity, Baryon vector current in the chiral and $1/N_c$ expansions, Phys. Rev. D90 (11) (2014) 114008. [arXiv:1407.0926](#), [doi:10.1103/PhysRevD.90.114008](#).
- [7] I. P. Fernando, J. L. Goity, these proceedings.
- [8] T. R. Hemmert, B. R. Holstein, J. Kambor, Phys.Lett. B395 (1997) 89–95. [arXiv:hep-ph/9606456](#).
- [9] T. R. Hemmert, B. R. Holstein, J. Kambor, J.Phys.G G24 (1998) 1831–1859. [arXiv:hep-ph/9712496](#).
- [10] I. P. Fernando, J. M. Alarcon, J. L. Goity, Baryon masses and σ terms in $SU(3)$ BChPT $\times 1/N_c$, Phys. Lett. B781 (2018) 719–722. [arXiv:1804.03094](#), [doi:10.1016/j.physletb.2018.04.054](#).
- [11] C. Alexandrou, V. Drach, K. Jansen, C. Kallidonis, G. Koutsou, Baryon spectrum with $N_f = 2 + 1 + 1$ twisted mass fermions, Phys. Rev. D90 (7) (2014) 074501. [arXiv:1406.4310](#), [doi:10.1103/PhysRevD.90.074501](#).
- [12] J. Gasser, H. Leutwyler, M. P. Locher, M. E. Sainio, Extracting the Pion - Nucleon Σ Term From Data, Phys. Lett. B213 (1988) 85–90. [doi:10.1016/0370-2693\(88\)91052-0](#).
- [13] J. Gasser, H. Leutwyler, M. E. Sainio, Sigma term update, Phys. Lett. B253 (1991) 252–259. [doi:10.1016/0370-2693\(91\)91393-A](#).
- [14] J. Gasser, H. Leutwyler, M. E. Sainio, Form-factor of the sigma term, Phys. Lett. B253 (1991) 260–264. [doi:10.1016/0370-2693\(91\)91394-B](#).
- [15] M. M. Pavan, I. I. Strakovsky, R. L. Workman, R. A. Arndt, The Pion nucleon Sigma term is definitely large: Results from a G.W.U. analysis of pi nucleon scattering data, PiN Newslett. 16 (2002) 110–115. [arXiv:hep-ph/0111066](#).
- [16] J. M. Alarcón, J. Martin Camalich, J. A. Oller, Phys. Rev. D85 (2012) 051503. [arXiv:1110.3797](#).
- [17] M. Hoferichter, J. Ruiz de Elvira, B. Kubis, U.-G. Meißner, Phys. Rev. Lett. 115 (2015) 092301. [arXiv:1506.04142](#).
- [18] M. Hoferichter, J. Ruiz de Elvira, B. Kubis, U.-G. Meißner, Phys. Lett. B760 (2016) 74–78. [arXiv:1602.07688](#).
- [19] J. Ellis, N. Nagata, K. A. Olive, Uncertainties in WIMP Dark Matter Scattering Revisited, Eur. Phys. J. C78 (7) (2018) 569. [arXiv:1805.09795](#), [doi:10.1140/epjc/s10052-018-6047-y](#).

- [20] H. Leutwyler, Theoretical aspects of Chiral Dynamics, PoS CD15 (2015) 022.
[arXiv:1510.07511](#), [doi:10.22323/1.253.0022](#).
- [21] C. Alexandrou, K. Hadjiyiannakou, C. Kallidonis, Axial charges of hyperons and charmed baryons using $N_f = 2 + 1 + 1$ twisted mass fermions, Phys. Rev. D94 (3) (2016) 034502.
[arXiv:1606.01650](#), [doi:10.1103/PhysRevD.94.034502](#).
- [22] N. Brambilla, J. Komijani, A. S. Kronfeld, A. Vairo, Relations between Heavy-light Meson and Quark Masses, Phys. Rev. D97 (3) (2018) 034503. [arXiv:1712.04983](#),
[doi:10.1103/PhysRevD.97.034503](#).