

$\tau \rightarrow \mu\mu\mu$ at a rate of one out of 10^{14} tau decays?

Patrick Blackstone*

*Department of Physics, Indiana University, Bloomington, IN 47408, USA and
Center for Exploration of Energy and Matter, Indiana University, Bloomington, IN 47408, USA*

Matteo Fael†

*Institut für Theoretische Teilchenphysik, Karlsruhe Institute of Technology (KIT), 76131 Karlsruhe, Germany and
Theoretische Physik I, Universität Siegen, 57068 Siegen, Germany*

Emilie Passemar‡

*Department of Physics, Indiana University, Bloomington, IN 47408, USA
Center for Exploration of Energy and Matter, Indiana University, Bloomington, IN 47408, USA and
Theory Center, Thomas Jefferson National Accelerator Facility, Newport News, VA 23606, USA
(Dated: December 13, 2019)*

We present in a full analytic form the partial widths for the lepton flavour violating decays $\mu^\pm \rightarrow e^\pm e^+ e^-$ and $\tau^\pm \rightarrow \ell^\pm \ell'^+ \ell'^-$, with $\ell, \ell' = \mu, e$, mediated by neutrino oscillations in the one-loop diagrams. Compared to the first result by Petkov in [1], obtained in the zero momentum limit $\mathcal{P} \ll m_\nu \ll M_W$, we retain full dependence on \mathcal{P} , the momenta and masses of external particles, and we determine the branching ratios in the physical limit $m_\nu \ll \mathcal{P} \ll M_W$. We show that the claim in [2] that the $\tau \rightarrow \ell\ell'\ell'$ branching ratios could be as large as 10^{-14} , as a consequence of keeping the \mathcal{P} dependence, is flawed. We find rates of order 10^{-55} , even smaller than those obtained in the zero momentum limit, as the latter prediction contains an unphysical logarithmic enhancement.

INTRODUCTION

It is widely reported by several experimental collaborations, e.g. by CMS [3], ATLAS [4], LHCb [5], BABAR [6–9] and Belle [10], that the branching ratios for the charged lepton flavour violating (CLFV) decay $\tau^\pm \rightarrow \ell^\pm \ell'^+ \ell'^-$, with $\ell, \ell' = e, \mu$, can be as large as 10^{-14} in the Standard Model extended with either a Dirac or a Majorana mass term for neutrinos. This follows from a claim by Pham in [2] that for these decays the GIM mechanism [11] produces a suppression of the form $|\sum_i U_{\ell i} U_{L i}^* \log x_i|^2$, where $x_i = m_{\nu i}^2/M_W^2$, $i = 1, 2, 3$, $m_{\nu i}$ and M_W are the masses of the three neutrinos and the W boson, and U is the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) mixing matrix [12, 13]. The result in [2] is in sharp contrast with the first evaluation by Petkov [1], which showed that these CLFV decays are instead power suppressed by $|\sum_i U_{\ell i} U_{L i}^* x_i \log x_i|^2$, so that the smallness of the ratios $m_{\nu i}/M_W$ crushes the branching fractions well below 10^{-54} , far beyond the sensitivity of any foreseeable experiment.

The calculations in [1] and [2] differ as follows. Ref. [1] employed for the evaluation of the one-loop diagrams the zero-momentum-limit (ZML) approximation, which assumes vanishing momenta and masses of the external particles while it retains the dependence on the internal masses of neutrinos and the W boson. The ZML, implicitly assumes the mass scale hierarchy

$$(\text{ZML}) \quad \mathcal{P} \ll m_{\nu i} \ll M_W,$$

where \mathcal{P} generically stands for any of the external particle momenta and masses, e.g. $\mathcal{P} \sim m_L$ or $\mathcal{P} \sim m_\ell$. This

approximation, even if far from the physical situation, allows a substantial simplification of the one-loop integrals, as in this way they depend only on x_i . Ref. [2], on the contrary, argued that once the external momentum dependence is taken into account, the GIM cancellation in $L \rightarrow \ell\ell'\ell'$, with $L = \tau$ or μ , becomes actually much milder, with a suppression only of the form $|\sum_i U_{\ell i} U_{L i}^* \log x_i|^2$, which leads to branching ratio values of the order of 10^{-14} .

If the prediction in [2] were true, it would imply, with current values for neutrino mixing angles and mass splittings [14], that the branching ratio of $\mu \rightarrow eee$ could reach 10^{-17} , for a lightest-neutrino mass of the order of 10^{-10} eV or smaller. This would be just around the corner for the MU3E experiment currently under construction at the Paul Scherrer Institute, Switzerland, which aims to reach a sensitivity of $\text{Br}(\mu \rightarrow eee) \sim 10^{-16}$ [15]. For the tau, the rates would be in the range $10^{-16} - 10^{-13}$, still several order of magnitudes smaller than the current world averages, $\text{Br}(\tau \rightarrow \ell\ell'\ell') \lesssim 10^{-8}$ [16], and the expected sensitivity of Belle II, $\text{Br}(\tau \rightarrow \ell\ell'\ell') \lesssim 10^{-10}$ [17], and the HL-LHC, $\text{Br}(\tau \rightarrow \ell\ell'\ell') \lesssim 10^{-9}$ [18–20], but more than forty order of magnitudes larger than the prediction of [1].

Therefore, the question we address in this letter is if the branching ratios of $L \rightarrow \ell\ell'\ell'$ can really change so dramatically once one assumes the physical limit (PL), i.e. the hierarchy

$$(\text{PL}) \quad m_{\nu i} \ll \mathcal{P} \ll M_W,$$

instead of the ZML, and keeps full dependence of external momenta and masses in the loop diagrams.




FIG. 1. One-loop diagrams contributing to the CLFV decay $L^- \rightarrow \ell^- \ell'^+ \ell'^-$ in the unitary gauge: the Z penguin (a,b), the photon penguin (c) and the box (d). Wave function corrections must be considered as well. In Feynman gauge there are also diagrams with Goldston bosons.

In [2] it is argued that in the Z -penguin diagram shown in Fig. 1a there are two propagators of nearly massless fermions, which give rise to a $\log x_i$ when the momentum q of the Z boson approaches $q^2 = 0$. This argument is supported by a computation of the Z -penguin as an expansion in q^2/M_W^2 taking the form $f(q^2, x_i) = f_0(x_i) + (q^2/M_W^2)f_1(x_i) + \dots$. By noticing that $f_0 \sim x_i \log x_i$ is suppressed — the term computed in [1] — while $f_1 \sim \log x_i$ is not, Ref. [2] concludes that $(q^2/M_W^2) \log x_i$ must dominate the branching ratio in the $x_i \rightarrow 0$ limit.

This conclusion is flawed. First of all, in [2] the loop integrals written in terms of Feynman parameters are computed via a simple Taylor expansion of the denominator appearing inside. Such approximation is not legitimate for arbitrary values of the Feynman parameters, so it does not lead to an expansion of the integral itself. A proper asymptotic expansion of a Feynman integral can be obtained, for instance, via the expansion-by-regions method [21, 22], in which one divides the whole integration domain into various regions and then performs different Taylor expansions in each region. Only the sum of all regions' contributions eventually yields the desired asymptotic series. In addition to that, even if such q^2/M_W^2 expansion were performed correctly, the calculation presented in [2] implicitly assumes also the hierarchy $q^2 \ll m_{\nu_i}^2$. Therefore, the series expansion $f_0(x_i) + (q^2/M_W^2)f_1(x_i) + \dots$ does not reproduce the correct $x_i \rightarrow 0$ limit at fixed values of q^2 , since this limit lies beyond the validity range of $q^2 \ll m_{\nu_i}^2$.

Recently, Ref. [23] presented a calculation of $L \rightarrow \ell \ell' \ell'$ in the PL, in which the one-loop diagrams are numerically evaluated with full dependence on \mathcal{P} . At variance with [2], they found branching ratios compatible with those in the ZML or smaller. However, the authors of Ref. [23] neglect the contribution from γ -penguins (as in Fig. 1c) and therefore their results are gauge dependent. Indeed, in processes with flavour changing neutral currents the gauge cancels entirely only in the sum of boxes, Z - and γ -penguins [24, 25]. Even if Ref. [1] retained only the logarithmic enhanced term $x_i \log x_i$ arising only from the Z penguins and the boxes, the omission of γ -penguins is not legitimate anymore as soon as one departs from this approximation. So we are still left with the doubt if the branching ratios in [23] are smaller as a consequence of

calculating in the Feynman gauge or if there is a deeper physical meaning.

In this letter we present the decay widths of $L \rightarrow \ell \ell' \ell'$ in the PL, fully analytic in M_W , m_ν and external momenta and masses. We compute them by making a systematic asymptotic expansion in \mathcal{P}/M_W and m_ν/\mathcal{P} of all Feynman diagrams by means of the expansion by regions. We will show that the neutrino mass dependence $|\sum_i x_i \log x_i|^2$ in the ZML is replaced in the PL by a much smaller enhancement $|\sum_i x_i \log(m_L^2/M_W^2)|^2$. We will give an explanation of this exchange of mass scales in the logarithm by analysing the effective operators mediating the decay once the Z and the W bosons are integrated out.

DETAILS OF THE CALCULATION

Let us consider the SM extended with neutrino mass, either a Dirac or Majorana. The flavour eigenstates of the left-handed neutrino fields $\nu_{\ell L}$ entering in the weak interactions become linear combinations of the three mass eigenstates ν_i with masses m_{ν_i} :

$$\nu_{\ell L} = \sum_{i=1}^3 U_{\ell i} \nu_i, \quad \ell = e, \mu, \tau, \quad (1)$$

where ν_{iL} is the left-handed component of ν_i and U is the PMNS matrix. The decay of a heavy lepton $L = \mu, \tau$ into three lighter charged leptons $\ell, \ell' = \mu, e$

$$L^\pm \rightarrow \ell^\pm \ell'^+ \ell'^-, \quad (2)$$

with masses m_L , m_ℓ and $m_{\ell'}$, respectively, proceeds then via three classes of one-loop diagrams shown in Fig. 1: the boxes, the Z and γ penguins. We neglect diagrams with the exchange of a Higgs boson as they are further suppressed by two extra powers of $1/M_W^2$ due the Yukawa interaction. The partial width given in [1] was obtained in the ZML. In this approximation, the one-loop integrals depend only on the ratio x_i . To leading order in x_i , the

	Branching ratio (NO)			Branching ratio (IO)		
	ZML	PL	ZML/PL	ZML	PL	ZML/PL
$\mu \rightarrow eee$	4.1×10^{-54}	2.9×10^{-55}	14	6.1×10^{-54}	4.6×10^{-55}	14
$\tau \rightarrow \mu\mu\mu$	2.0×10^{-53}	5.8×10^{-55}	34	2.0×10^{-53}	5.8×10^{-55}	34
$\tau \rightarrow \mu ee$	1.3×10^{-53}	3.8×10^{-55}	35	1.3×10^{-53}	3.8×10^{-55}	35
$\tau \rightarrow eee$	1.1×10^{-54}	3.3×10^{-56}	34	6.1×10^{-55}	1.9×10^{-56}	32
$\tau \rightarrow e\mu\mu$	7.6×10^{-55}	2.1×10^{-56}	36	4.1×10^{-55}	1.2×10^{-56}	34

TABLE I. Branching ratio for the CLFV decays $L \rightarrow \ell\ell'\ell'$ in the ZML and the PL for normal ordering (NO) and inverted ordering (IO) of neutrino masses. The ratio between the two is also reported. In the ZML we assume $m_1 = 0$ (NO) or $m_3 = 0$ (IO).

amplitudes of the three classes of diagrams are:

$$i\mathcal{A}_Z = \frac{\alpha G_F}{\sqrt{2}\pi} \sum_i U_{\ell i} U_{L i}^* x_i (3 + \log x_i) \times \left[(\bar{\ell} \gamma^\mu P_L L) (\bar{\ell}' \gamma_\mu \ell') - \frac{1}{2 \sin^2 \theta_W} (\bar{\ell} \gamma^\mu P_L L) (\bar{\ell}' \gamma_\mu P_L \ell') \right], \quad (3)$$

$$i\mathcal{A}_\gamma = -\frac{\alpha G_F}{\sqrt{2}\pi} (\bar{\ell} \gamma^\mu P_L L) (\bar{\ell}' \gamma_\mu \ell') \sum_i U_{\ell i} U_{L i}^* x_i, \quad (4)$$

$$i\mathcal{A}_{\text{Box}} = \frac{\alpha G_F}{2\sqrt{2}\pi \sin^2 \theta_W} (\bar{\ell} \gamma^\mu P_L L) (\bar{\ell}' \gamma_\mu P_L \ell') \times \sum_i U_{\ell i} U_{L i}^* x_i (1 + \log x_i), \quad (5)$$

where G_F and α are the Fermi and fine structure constants, respectively, and $\sin^2 \theta_W$ is the sine of the Weinberg angle. Retaining only the terms enhanced by $\log x_i \sim 50$, which appear in the boxes and Z penguins, one obtains the prediction for the rate in the ZML [1]:

$$\frac{\Gamma(L \rightarrow \ell\ell\ell)}{\Gamma_0} = \frac{3\alpha^2}{16\pi^2} \left| \sum_{i=2}^3 U_{\ell i} U_{L i}^* \frac{\Delta m_{i1}^2}{M_W^2} \log \frac{\Delta m_{i1}^2}{M_W^2} \right|^2, \quad (6)$$

and $\Gamma(L \rightarrow \ell\ell'\ell', \ell \neq \ell') = \frac{2}{3}\Gamma(L \rightarrow \ell\ell\ell)$, where $\Gamma_0 = G_F^2 m_L^5 / (192\pi^3)$. Eq. (6) is obtained by taking the limit $m_1 \rightarrow 0$ and assumes normal neutrino mass hierarchy, i.e. $m_1 < m_2 < m_3$, showing that the rate actually depends on the neutrino mass splittings Δm_{31}^2 and Δm_{21}^2 since the GIM-suppression term $\sum_i U_{\ell i} U_{L i}^* x_i \log x_i$ vanishes in the equal mass limit. For inverted mass hierarchy, the subscript ‘1’ must be substituted with ‘3’ and $i = 1, 2$. Eq. (6) also neglects subleading $m_{\ell, \ell'}/m_L$ corrections from phase space integration. The values of the branching ratios in the ZML with normal and inverted mass hierarchy are reported in Tab. I. Current PDG values are employed for the lepton masses, neutrino mass splittings and neutrino mixing angles [14].

Let us now describe our calculation performed in the PL. We generated the complete set of Feynman diagrams and their relative counter-terms, using FEYNARTS [26]

with a modified version of the SM file to account for neutrino masses and lepton flavour mixing. The amplitudes were reduced to one-loop tensor integrals using FORM [27], via the FORMCALC package [28], keeping the complete dependence on M_W , \mathcal{P} and $m_{\nu i}$. The setup was independently checked by a second implementation based on FEYNALC [29].

Nowadays, lengthy expressions for the tensor integrals could be obtained in principle in an analytical form with full dependence on $m_{\nu i}, M_W, m_L, m_{\ell, \ell'}$ and the invariants $s_{ij} = (p_i + p_j)^2$, with p_{1-3} the momenta of the three outgoing leptons, however their use is impractical. It is therefore more helpful to compute them as series in the small parameters \mathcal{P}^2/M_W^2 and $m_{\nu i}^2/\mathcal{P}^2$. To this end, we employed the method of expansion by regions (for an introduction see e.g. [22]). For all one-loop diagrams, we divided the integration domain into different regions and, for each region, we perform a Taylor expansion with respect to the parameters that are considered small there. Afterwards, by integrating every expanded integrand over the whole domain, and by summing the contributions from all the regions, we obtained the desired asymptotic expansion of the original one-loop diagram. The advantage of this method, compared for instance to an expansion of the full result *after* integration, is that the integrals arising in each region can be handled much more easily than the initial one, as typically they depend on just one or two mass scales.

We performed first an expansion assuming $\mathcal{P} \sim m_{\nu i} \ll M_W$, without distinguishing at this point the two scales $m_{\nu i}$ and \mathcal{P} . In a second step, the integrals arising from the first stage are further expanded in the limit $m_{\nu i} \ll \mathcal{P}$. The total amplitude is then obtained by retaining only the dependence on $m_{\nu i}$, while higher order terms further suppressed by \mathcal{P}^2/M_W^2 or $m_{\nu i}^2/\mathcal{P}^2$, or terms independent on $m_{\nu i}$, are discarded. We perform several numerical checks at different stages of the calculation as a sanity check. To this end we took advantage of MATHEMATICAS arbitrary-precision numbers and PACKAGE-X’s analytic expressions of one-loop integrals [30], available in any kinematic configuration. We verified that our approximated expressions for the tensor integrals become in-

creasingly accurate both by including higher order terms in the expansion as well as by taking the limit $M_W \rightarrow \infty$ and $m_{\nu i} \rightarrow 0$, at fixed values of \mathcal{P} .

RESULTS

The partial widths are given by integrating the approximated expression for the squared amplitude over the three-particle phase space of $L \rightarrow \ell \ell' \ell'$. These massive phase space integrals depend on two variables s_{ij} , plus two or three masses of the external particles. By employing the expansion by regions one more time, we computed the phase space integrals as series in $m_{\ell(\ell')}/m_L$, retaining only the leading terms in the final expressions for the rate. We obtain:

$$\begin{aligned} \frac{\Gamma(L \rightarrow \ell \ell \ell)}{\Gamma_0} &= \frac{3\alpha^2}{16\pi^2} \left| \sum_{i=2}^3 U_{\ell i} U_{L i}^* \frac{\Delta m_{i1}^2}{M_W^2} \right|^2 \\ &\times \left[\log^2 x_L + 2 \log x_L - \frac{1}{6} \log x_\ell + \frac{19}{18} + \frac{17}{18} \pi^2 \right. \\ &\quad \left. - \frac{1}{\sin^2 \theta_W} \left(\log x_L + \frac{11}{12} \right) + \frac{3}{8 \sin^4 \theta_W} \right], \quad (7) \end{aligned}$$

where $x_L = m_L^2/M_W^2$ and $x_\ell = m_\ell^2/M_W^2$. For $L \rightarrow \ell \ell' \ell'$ ($\ell \neq \ell'$) we have:

$$\begin{aligned} \frac{\Gamma(L \rightarrow \ell \ell' \ell')}{\Gamma_0} &= \frac{3\alpha^2}{16\pi^2} \left| \sum_{i=2}^3 U_{\ell i} U_{L i}^* \frac{\Delta m_{i1}^2}{M_W^2} \right|^2 \\ &\times \left[\frac{2}{3} \log^2 x_L + \frac{25}{18} \log x_L - \frac{1}{6} \log x_{\ell'} + \frac{55}{108} + \frac{2}{3} \pi^2 \right. \\ &\quad \left. - \frac{1}{\sin^2 \theta_W} \left(\frac{\log x_L}{2} + \frac{11}{24} + \frac{\pi^2}{18} \right) \right. \\ &\quad \left. + \frac{1}{\sin^4 \theta_W} \left(\frac{3}{16} + \frac{\pi^2}{36} \right) \right]. \quad (8) \end{aligned}$$

At variance with the result presented in [2], our expression for the rates is power suppressed by $|\sum_i U_{\ell i} U_{L i}^* x_i|^2$ and yields values for the branching ratios of the order of 10^{-55} , see Tab. I. Moreover, compared to Eq. (6) in the ZML, Eqs. (7) and (8) do not have a logarithmic enhancement $\log^2 x_i \sim 2500$. On the contrary in its place we get only $\log^2 x_\mu \sim 176$ or $\log^2 x_\tau \sim 58$, which are of comparable size with respect to other terms appearing in Eqs. (7) and (8). For this reason, the branching ratios in the PL turn out to be about one order of magnitude smaller than those in the ZML.

Note also that the presence of the singular terms $\log x_\ell$ or $\log x_{\ell'}$ is not in contradiction with the Kinoshita-Lee-Nauenberg theorem [31, 32] and the cancellation of mass singularities for inclusive observables. In fact

Eqs. (7) and (8) are valid strictly in the PL, i.e. when $m_{\nu i} \ll m_{\ell(\ell')}$. The limiting case of vanishing charged-lepton masses and non-zero neutrino masses violates the assumptions of our derivation and therefore is not a meaningful limit of our expressions.

In the case of purely Majorana masses (with no Dirac component), the neutrino weak eigenstates are still a combination of three (Majorana) fields, however the PMNS matrix takes on two new CP violating phases. These phases (1, α_{21} , and α_{31}) multiply the three columns of the Dirac mixing matrix so that $U_{\ell 1} \rightarrow 1 \cdot U_{\ell 1}$, $U_{\ell 2} \rightarrow e^{i\alpha_{21}/2} U_{\ell 2}$, and $U_{\ell 3} \rightarrow e^{i\alpha_{31}/2} U_{\ell 3}$ for all ℓ . In Eqs. (6-8) the PMNS elements show up in pairs as $U_{L i}^* U_{\ell i}$, with i always the same in the two factors, so the Majorana phases cancel.




FIG. 2. Example of two diagrams mediating $L \rightarrow \ell \ell' \ell'$ in a low energy effective field theory description.

Finally, we can understand the mechanism that converts the $x_i \log x_i$ in the ZML into a $x_i \log x_L$ in the PL by looking at the effective operators arising after integrating out the Z and the W bosons. Let us for simplicity concentrate only on the operator associated with such logarithmic enhancement and let us focus for instance on the box in Fig. 1d. Both in the ZML and in the PL, we can shrink the two W propagators to a point-like interaction and match the amplitude onto the the following dimension-six and dimension-eight operators:

$$\begin{aligned} O_6^{L\ell'} &= (\bar{\nu}_i \gamma^\mu P_L L) (\bar{\ell}' \gamma_\mu P_L \nu_j), \\ O_6^{\ell\ell'} &= (\bar{\ell} \gamma^\mu P_L \nu_i) (\bar{\nu}_j \gamma_\mu P_L \ell'), \\ O_8 &= m_{\nu i}^2 (\bar{\ell} \gamma^\mu P_L L) (\bar{\ell}' \gamma_\mu P_L \ell'). \quad (9) \end{aligned}$$

The first two operators correspond to the usual Fermi interaction mediating μ and τ leptonic decays. They contribute to $L \rightarrow \ell \ell' \ell'$ via the one-loop diagram in Fig. 2a. The third operator in (9) is necessary to renormalize the effective theory, i.e. to cancel the UV divergence from the diagram 2a. These operators' Wilson coefficients are:

$$\begin{aligned} C_6^{L\ell'} &= \frac{4G_F}{\sqrt{2}} U_{\ell' j} U_{L i}^*, & C_6^{\ell\ell'} &= \frac{4G_F}{\sqrt{2}} U_{\ell' j}^* U_{\ell i}, \\ C_8(\mu) &= \frac{G_F^2}{2\pi^2} \log \left(\frac{M_W^2}{\mu^2} \right) U_{\ell i} U_{L i}^*. \quad (10) \end{aligned}$$

We can imagine performing the matching between the SM and the effective theory at a scale $\mu = M_W$, and evolving the coefficients to a lower scale via the renormalization group. The coefficient C_8 explicitly depends on

the renormalization scale μ and this dependence reveals the difference between the PL and the ZML. In the ZML, the evolution of C_8 can proceed down to a scale $\mu \sim m_{\nu i}$, at which point we can integrate out the neutrinos and remove the operators $O_6^{L\ell'}$ and $O_6^{\ell\ell'}$ which contain the neutrino field. We are then left with an EFT with only O_8 , whose Wilson coefficient is frozen at $C_8(m_{\nu i})$, i.e. it contains a $\log(M_W^2/m_{\nu i}^2)$ (compare with Eq. (5)). On the contrary, in the PL, C_8 can run only until the scale $\mu \sim m_L$ is reached. In this case, all operators in (9) are still active at the scale m_L , however C_8 produces only a milder $\log(M_W^2/m_L^2)$ enhancement. Therefore, the ZML overestimates the values for the branching ratios as it allows an unphysical evolution of these operators between M_W and $m_{\nu i}$, while in reality the running stops at the physical intermediate scale m_L where the process happens.

CONCLUSIONS

Several experimental collaborations reported that the branching fractions of $L \rightarrow \ell\ell'\ell'$ can be as large as 10^{-14} , following the observation in [2] that the GIM cancellation is not so severe and takes the form of $|\sum_i U_{\ell i} U_{L i}^* \log x_i|^2$. In this letter we showed that this conclusion is wrong.

We calculated and presented for the first time the branching ratios in the PL by performing a series expansion of all one-loop diagrams in the small parameters \mathcal{P}/M_W and $m_{\nu i}/\mathcal{P}$. Our fully analytic expressions prove that the GIM suppression in these decays is power-like $|\sum_i U_{\ell i} U_{L i}^* x_i|^2$, similarly to what was shown long ago in [1] for the ZML, so that the claim from [2] must be rejected. We predict the branching ratios in the Standard Model including neutrino masses to be of the order of 10^{-55} . [MISSING FINALE HERE]

Acknowledgments A special thank to J. Piclum for fruitful discussions about the method of regions. We wish to thank also A. Friedland, T. Huber, and A. Signer for useful discussions. P.B. thanks also the Theoretical Particle Physics Group at the University of Siegen for the hospitality during his stay in June 2019. The work of M.F. is supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under grant 396021762 - TRR 257 “Particle Physics Phenomenology after the Higgs Discovery”. The work of P.B. and E.P. is supported in part by the U.S. Department of Energy (contract DE-AC05-06OR23177) and National Science Foundation (PHY-1714253).

* pblackst@iu.edu

† matteo.fael@kit.edu

‡ epassema@indiana.edu

- [1] S. T. Petcov, Sov. J. Nucl. Phys. **25**, 340 (1977), [Erratum: Yad. Fiz.25,1336(1977)].
- [2] X.-Y. Pham, Eur. Phys. J. **C8**, 513 (1999), arXiv:hep-ph/9810484 [hep-ph].
- [3] *Search for $\tau \rightarrow 3\mu$ decays using τ leptons produced in D and B meson decays*, Tech. Rep. CMS-PAS-BPH-17-004 (CERN, Geneva, 2019).
- [4] G. Aad *et al.* (ATLAS), Eur. Phys. J. **C76**, 232 (2016), arXiv:1601.03567 [hep-ex].
- [5] R. Aaij *et al.* (LHCb), JHEP **02**, 121 (2015), arXiv:1409.8548 [hep-ex].
- [6] B. Aubert *et al.* (BaBar), Phys. Rev. Lett. **92**, 121801 (2004), arXiv:hep-ex/0312027 [hep-ex].
- [7] B. Aubert *et al.* (BaBar), Phys. Rev. Lett. **99**, 251803 (2007), arXiv:0708.3650 [hep-ex].
- [8] B. Aubert *et al.* (BaBar), Phys. Rev. Lett. **100**, 071802 (2008), arXiv:0711.0980 [hep-ex].
- [9] J. P. Lees *et al.* (BaBar), Phys. Rev. **D81**, 111101 (2010), arXiv:1002.4550 [hep-ex].
- [10] K. Hayasaka *et al.*, Phys. Lett. **B687**, 139 (2010), arXiv:1001.3221 [hep-ex].
- [11] S. L. Glashow, J. Iliopoulos, and L. Maiani, *Meeting of the Italian School of Physics and Weak Interactions Bologna, Italy, April 26-28, 1984*, Phys. Rev. **D2**, 1285 (1970).
- [12] B. Pontecorvo, Sov. Phys. JETP **26**, 984 (1968), [Zh. Eksp. Teor. Fiz.53,1717(1967)].
- [13] Z. Maki, M. Nakagawa, and S. Sakata, Prog. Theor. Phys. **28**, 870 (1962), [34(1962)].
- [14] M. Tanabashi *et al.* (Particle Data Group), Phys. Rev. **D98**, 030001 (2018).
- [15] A. Blondel *et al.*, (2013), arXiv:1301.6113 [physics.ins-det].
- [16] Y. S. Amhis *et al.* (HFLAV), (2019), arXiv:1909.12524 [hep-ex].
- [17] W. Altmannshofer *et al.* (Belle-II), (2018), arXiv:1808.10567 [hep-ex].
- [18] A. Cerri *et al.*, (2018), arXiv:1812.07638 [hep-ph].
- [19] S. Fiorendi (CMS), *Proceedings, 17th International Conference on B-Physics at Frontier Machines (Beauty 2018): La Biodola, Elba island, Italy, May 6-11, 2018*, PoS **BEAUTY2018**, 056 (2018).
- [20] P. Azzi *et al.* (HL-LHC, HE-LHC Working Group), (2019), arXiv:1902.04070 [hep-ph].
- [21] M. Beneke and V. A. Smirnov, Nucl. Phys. **B522**, 321 (1998), arXiv:hep-ph/9711391 [hep-ph].
- [22] V. A. Smirnov, *Berlin, Germany: Springer (2002) 262 p, (Springer tracts in modern physics. 177)*, Springer Tracts Mod. Phys. **177**, 1 (2002).
- [23] G. Hernandez-Tom, G. Lopez Castro, and P. Roig, Eur. Phys. J. **C79**, 84 (2019), arXiv:1807.06050 [hep-ph].
- [24] T. Inami and C. S. Lim, Prog. Theor. Phys. **65**, 297 (1981), [Erratum: Prog. Theor. Phys.65,1772(1981)].
- [25] G. Buchalla, A. J. Buras, and M. K. Harlander, Nucl. Phys. **B349**, 1 (1991).
- [26] T. Hahn, Comput. Phys. Commun. **140**, 418 (2001), arXiv:hep-ph/0012260 [hep-ph].
- [27] J. Kuipers, T. Ueda, J. A. M. Vermaseren, and J. Vollinga, Comput. Phys. Commun. **184**, 1453 (2013), arXiv:1203.6543 [cs.SC].
- [28] T. Hahn and M. Perez-Victoria, Comput. Phys. Commun. **118**, 153 (1999), arXiv:hep-ph/9807565 [hep-ph].
- [29] V. Shtabovenko, R. Mertig, and F. Orellana, Comput. Phys. Commun. **207**, 432 (2016), arXiv:1601.01167 [hep-

- ph].
- [30] H. H. Patel, Comput. Phys. Commun. **197**, 276 (2015), arXiv:1503.01469 [hep-ph].
- [31] T. Kinoshita, J. Math. Phys. **3**, 650 (1962).
- [32] T. D. Lee and M. Nauenberg, Phys. Rev. **133**, B1549 (1964).