Searching for an Enhanced Signal of the onset of Color Transparency in Baryons with D(e, e'p)n scattering

Shujie Li¹, Carlos Yero², Jennifer Rittenhouse West¹, Clare Bennett³, Wim Cosyn^{3,5}, Douglas Higinbotham⁴, Misak Sargsian³, Holly Szumila-Vance*,⁴

- Lawrence Berkeley National Laboratory, Berkeley, CA 94720 USA
- Old Dominion University, Norfolk, VA 23529 USA
- Florida International University, Miami, FL 33199 USA
- Jefferson Lab, Newport News, VA 23601 USA
- Ghent University, B9000 Ghent, Belgium
- Correspondence: hszumila@jlab.org

Abstract: Observation of the onset of color transparency in baryons would provide a new means of studying the nuclear strong force and would be the first clear evidence of baryons transforming into a color-neutral point-like size in the nucleus as predicted by Quantum Chromodynamics. Recent C(e, e'p)results from electron-scattering at Jefferson Lab (JLab) did not observe the onset of color transparency (CT) in protons up to $Q^2 = 14.2 \text{ GeV}^2$. The traditional methods of searching for CT in (e, e'p) scattering use heavy targets favoring kinematics with already initially reduced final state interactions (FSIs) such that any CT effect that further reduces FSIs will be small. The reasoning behind this choice is the difficulty in accounting for all FSIs. D(e, e'p)n, on the other hand, has well-understood FSI contributions from double scattering with a known dependence on the kinematics and can show an increased sensitivity to hadrons in point-like configurations. Double scattering is the square of the re-scattering amplitude in which the knocked-out nucleon interacts with the spectator nucleon, a process that is suppressed in the presence of point-like configurations and is particularly well-studied for the deuteron. This suppression yields a quadratic sensitivity to CT effects and is strongly dependent on the choice of kinematics. Here, we describe a possible JLab electron-scattering experiment that utilizes these kinematics and explores the potential signal for the onset of CT with enhanced sensitivity as compared to recent experiments.

Keywords: color transparency; color confinement; QCD; hadronization





Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/).

1. Introduction

Quantum chromodynamics (QCD) is the fundamental theory of the strong interaction and the basis for all nuclear phenomena. However, the poorly understood nonperturbative dynamics of QCD and the phenomenon of confinement make the establishment of the link between QCD and nuclear dynamics one of most important tasks of modern nuclear physics. High energy and momentum transfer (i.e., hard) processes involving nuclei are the best testing grounds for understanding the link between QCD and nuclear physics. The advantage of hard processes is that scattering that takes place on an individual quark in the nucleon can be clearly identified. One outstanding puzzle for hard processes is elastic scattering from a nucleon at large Q^2 . Within the partonic model of Feynman [1], this scattering is predicted to be dominated by single quark, hard, large-angle elastic scattering as the quark momentum fraction approaches $x_B \to 1$, a process that does not affect the size of the hadron [2]. In QCD, however, the prediction is different although also dominated by the $x_B \to 1$ regime [3]. The phenomenon of color transparency (CT) [4–7] as predicted by QCD states that for exclusive processes at high momentum transfer, the cross section is dominated by a hadronic state of very small transverse size, effectively selecting it from the initial state configuration containing an infinite number of states of all sizes. Such a small size state (commonly referred as point-like

configuration (PLC)) is also color neutral when producing final (observable) hadrons. An expectation that PLCs are color neutral in hard exclusive processes constitutes the basis for the prediction of color transparency phenomena according to which, hard exclusive processes probed in nuclear medium should observe reduced final state interaction (FSI) for produced hadrons [4,8]. This prediction is complicated by the fact that PLC's are not eigenstates of the Hamiltonian, they are quantum-mechanical wave packets and therefore will expand as they propagate in the nuclear medium thereby reducing the expected CT effects [9]. To overcome expansion effects the experiments require high enough energy that Lorentz time dilation will delay such an expansion.

The color transparency effect described above for hard exclusive production of *mesons* is analogous to electric charge transparency in electromagnetic processes, for example, as observed in $\pi^0 \to e^+e^-\gamma$ decays through the reduced ionization in the medium by e^+e^- pairs at distances close to the decay point [10]. Indeed, several experiments investigating hard exclusive electroproduction of ρ and π mesons from nuclei observed signatures consistent with color transparency. Pion production measurements at JLab reported evidence for the onset of CT [11] in the process $e + A \to e' + \pi^+ + A^*$. The results of the pion electroproduction experiment showed that both the energy and A dependence of the nuclear transparency deviate from conventional nuclear physics and are consistent with models that include CT. The results indicate that the energy scale for the onset of CT in mesons is ~ 1 GeV. A CLAS experiment studied ρ -meson production from nuclei, and the results also indicated an early onset of CT in mesons [12]. An increase of the transparency with Q^2 for both C and Fe was observed. The rise in transparency was found to be consistent with predictions of CT by models [13–15] which had also accounted for the increase in transparency for pion electroproduction. The π and ρ electroproduction data set the energy range to be a few GeV for the onset of CT in mesons.

The uniqueness of QCD, as encoded in the $SU(3)_C$ symmetry obeyed by strong force interactions, also predicts the formation of color singlet 3-quark qqq PLCs. As a result, in addition to mesons, one expects to observe CT effects also for hard exclusive electroproduction of baryons. Thus, an observation of CT in the baryonic sector will indicate a unique interplay of $SU(3)_C$ symmetry and nuclear forces probed in the nuclear medium. The onset of CT is favored to be observed at lower energy in mesons than baryons since only two quarks must come close together, and therefore the quark-antiquark pair is more likely to form a PLC [16] than a three quark baryon. The QCD subfield holographic light-front QCD predicts the onset of CT at 14 GeV² for the proton [3]. The kinematic regime favorable for observation of CT effects in the baryonic sector is as yet unknown.

The experimental observation of the signal of the onset of CT in baryons remains ambiguous. The first attempt measured large angle A(p,2p) scattering at Brookhaven National Lab (BNL) [17–20]. These experiments measured the transparency as the quasi-elastic cross section from the nuclear target to the free pp elastic cross section. The measurements were taken with perpendicular kinematics which, in electron scattering, is generally dominated by nucleon re-scattering [21,22]. The results of these experiments indicated a rise in the transparency for outgoing proton momenta of 6–9.5 GeV/c consistent with models of CT. However, the transparency was observed to decrease at higher momenta between 9.5–14.4 GeV/c. This decrease is inconsistent with CT alone. Possible explanations for the transparency decrease include an elastic energy dependent cross section due to nuclear filtering from the Landshoff mechanism [23,24] or excitation of charm resonances beyond the charm production threshold [25]. The BNL experiments have the added complication that the incident proton suffers a reduction of flux in medium (these are initial state interactions, ISIs) that must be included in any transparency calculation.

In the (e, e'p) reaction, however, only the final state proton suffers a reduction of flux and needs to be considered in the measurement. The first experiments using an electron beam

to measure CT were at SLAC [26,27] followed by experiments at JLab [28,29]. In high Q^2 quasielastic (e,e'p) scattering from nuclei, the electron scatters from a single proton, which has some associated Fermi motion [30]. In the plane wave impulse approximation (PWIA), the proton is ejected without final state interactions with the residual A-1 nucleons. The measured A(e,e'p) cross-section would deviate from the PWIA prediction in the presence of FSI, where the proton can scatter both elastically and inelastically from the surrounding nucleons as it exits the nucleus. The ratio of the measured cross section to the PWIA calculation defines the nuclear transparency T (see [31]). In complete CT, the FSI vanish, and the nuclear transparency plateaus at T=1. This is in contrast to the conventional picture for the nuclear transparency in A(e,e'p) processes in which the proton's initial momentum is small. There, the transparency will be less than 1 due to FSIs and other correlations and follows the same energy dependence as the NN cross section, which is relatively constant for momenta above the few GeV range.

None of the earlier A(e,e'p) experiments observed the onset of proton CT. However, the onset of CT depends both on momentum and energy transfers, affecting the "squeezing" and "freezing" of PLCs [32]. Since A(e,e'p) scattering measurements are carried out at $x_B=1$ kinematics, they have a high Lorentz factor characterized by lower energy transfers compared to momentum transfers. It is therefore possible that the JLab 6 GeV era experiments were unable to satisfy the energy dependence of the freezing requirement, even if the momentum transfer was sufficient to create a PLC. The most recent result on this topic was published with the C(e,e'p) data in Hall C at JLab as part of the 12 GeV program [31]. With the beam energy almost doubled as compared with previous JLab measurements, this experiment eliminated the possibility that the increase in transparency observed at BNL was due to the higher incident proton momentum than was attainable during the 6 GeV era.

Results from the recent CT experiment [31] do not indicate a rise in CT up to $Q^2 = 14.2 \, \text{GeV/c}^2$ within a 6% uncertainty. It should be noted that these results searched for the onset of CT in kinematics in which the initial momentum of the struck proton was small and the FSI effect was dominated by screening effects due to the interference of PWIA and single rescattering amplitudes. Such kinematics for medium nuclei are characterized by small amplitudes of FSI and the rescattering process takes place over the full radius of the nucleus. As a result the transparency T has a diminished sensitivity to CT effects, both due to possibility that PLC was not fully formed at the available Q^2 and/or that the PLC could undergo a full expansion during propagation in the nuclear medium after ep scattering. Thus the lack of the CT effects in the hard exclusive production of the proton from nuclei indicates that the effects are much smaller than initially expected, either due to the difficulty in production of small size PLCs at the available Q^2 or due to a fast expansion of a 3-quark PLC in nuclear medium.

In this paper we discuss how to enhance a potential signal by looking for the onset of CT using deuterium breakup and measuring changes in the yields for kinematics of known double scattering and screening regimes of FSIs as a function of increasing momentum transfer. We will show that the experiment under consideration is highly sensitive to final state rescattering of a PLC and that the distance of propagation of the PLC can be suppressed by the external kinematics of the recoil nucleon.

2. The role of D(e, e'p)n in CT studies

The deuteron is the simplest target nucleus and its dynamics in collisions are well-described using the generalized eikonal approximation (GEA) [33,34]. With a deuteron target, the kinematics in exclusive processes can be precisely chosen such that the inter-nucleon distances of the struck and spectator nucleon lead to well-controlled FSIs [35]. In the case of high Q^2 , with the onset of the CT regime, these reactions enable observations of the formation of the PLC before expansion and control its expansion in the FSI process.

The proton knockout reaction in deuterium breakup D(e, e'p)n is described by

$$e + D \rightarrow e' + p + n, \tag{1}$$

where the final state proton carries almost all of the momentum transferred to the deuteron and can be identified as a struck nucleon. In kinematics in which $Q^2 \geq 1~{\rm GeV^2}$, two main scattering amplitudes describe the dynamics of the process (Fig. 1). For small initial momenta of the struck proton ($\leq 200~{\rm MeV/c}$), the re-scattering amplitude as described in Fig. 1(b) is small due to the large inter-nucleon separation. Therefore, the deuteron cross section's deviation from PWIA is determined mainly by Glauber screening effects (*i.e.*, the interference between PWIA and FSI amplitudes). As a result, the transparency as defined by the ratio of the full cross section to the PWIA cross section is less than unity (see the 100 and 200 MeV/c curves in Fig.2). This effect is relatively small, thus deuterium is traditionally not a good target for CT measurements for the case of small initial momenta.




Figure 1. Electron scattering interaction probing deuterium with a) breakup and b) re-scattering effects in the initial deuterium rest frame [36].

However, if the initial momentum is selected to be above 300 MeV/c, the FSI amplitude (Fig. 2(b)) in near perpendicular kinematics (i.e. the recoil neutron angle is nearly perpendicular to the momentum transfer) significantly exceeds the PWIA contribution. As a result the transparency as described above will exceed unity (see the 400, 500 and 600 MeV/c curves in Fig.2). Note that the experimental missing momentum in the deuteron is equivalent to the recoil neutron momentum, p_r , which is fixed by kinematics. This demonstrates that the experimental kinematics can easily be tuned such that FSIs dominate the signal. In this situation, the measured cross sections are very sensitive to any change in FSI effects such as the onset of color transparency.

According to our discussion for perpendicular kinematics, CT will increase the full cross section at small recoil neutron momenta ($p_r \le 200 \text{ MeV/c}$) due to the decrease of Glauber screening effects. At the same time CT will decrease the full cross section at $p_r > 400 \text{ MeV/c}$ due to the decrease of double scattering contribution. Thus, the experimentally measured D(e,e'p)n cross section ratio of high to low recoiling neutron momentum at perpendicular kinematics can be used to track the size of FSI effects with higher sensitivity than in traditional measurements of CT effects in eelp processes on medium to heavy nuclei. Indeed, in this case CT makes FSI effects smaller, thereby decreasing the large p_r cross section (numerator) and increasing the small p_r cross section (denominator). As Q^2 increases, the onset of CT can be detected through a decrease of such ratios as compared to PWIA calculations. A possible way of performing such a high sensitivity CT measurement with the deuteron at JLab is discussed in Sec. 5.




Figure 2. The calculated cross section ratio with respect to the PWIA is shown for various neutron recoil angles θ_{nq} (i.e. angle is with respect to \vec{q}) at different values of the recoil neutron momenta for $Q^2 = 14 \, (\text{GeV}/c)^2$. Solid curves are calculated with FSI+PWIA [37]. Dashed curves with the same color and marker are the same calculation including a CT effect of $\Delta M^2 = 2 \, \text{GeV}^2$.

2.1. Recent D(e, e'p)n studies at high Q^2

The first studies of D(e,e'p)n at high Q^2 (> 1 (GeV/c)²) were carried out at JLab experimental Halls A [38] and B [39]. Both experiments observed a strong angular dependence of the D(e,e'p)n cross section with neutron recoil angle, θ_{nq} , peaking at $\theta_{nq} \sim 70^{\circ}$ in agreement with the GEA predictions [40]. Due to statistical limitations in Hall B, however, it was necessary to integrate over a wide angular range (θ_{nq}) to determine the deuteron momentum distributions. As a result, for recoil momenta > 300 MeV/c, these were dominated by FSIs, meson-exchange currents (MECs) and bound-nucleon excitations into a resonance state (isobar currents or ICs).

The Hall A high-resolution spectrometers, on the other hand, allowed for a measurement of the recoil momentum dependence of the D(e, e'p)n cross sections for fixed bins in θ_{nq} . Different theoretical models (i.e., Paris [41] and CD-Bonn-based [42] calculations) were tested for recoil momenta between $\sim 300-600\,\mathrm{MeV/}c$. The results from [38] also revealed a kinematic window at $\theta_{nq}\sim 35-45^\circ$ where the FSIs are significantly reduced and approximately independent of the recoil nucleon momentum. An extension of the D(e, e'p)n cross-section measurements beyond $p_r\sim 500\,\mathrm{MeV/}c$ is possible and may potentially reveal a transition between nucleonic and quark-gluon degrees of freedom.

Recent studies of deuterium electro-disintegration [43] carried out in JLab Hall C used the kinematic window ($\theta_{nq} \sim 35-45^{\circ}$) found in [38] to extend the D(e,e'p)n cross section measurements to very high recoil momenta ($p_r \sim 500\text{-}1000 \text{ MeV/}c$) and large Q^2 ($\sim 4-5 \text{ (GeV/}c)^2$). The results demonstrated that it is possible to select bins in θ_{nq} that minimize re-scattering effects and enhance the PWIA allowing for a direct probe of the least well-known part of the NN interaction at the sub-Fermi (<1 fm) distance scale, where nucleons begin to significantly overlap.

The extracted momentum distributions from [43] were compared to the theoretical momentum distributions based on either the Paris [41] (calculated by J.M. Laget [44]), AV18 [45] or CD-Bonn [42] (calculated by M. Sargsian [46]) potentials. For recoil momenta below 400

MeV/c, all theoretical calculations agree with data, however, at larger recoil momenta, the data was best reproduced by the calculations using the CD-Bonn [42] potential up to recoil momenta 700 MeV/c. Beyond this momentum range, current theoretical calculations start to break down and are unable to describe data.

Nevertheless, the progress made in exploring the short-range part of the NN interaction using deuterium has demonstrated that, unlike in heavier nuclei, FSIs are well-understood from both an experimental and theoretical perspective. It has also been demonstrated, particularly in [43], that it is possible to select specific kinematics of θ_{nq} to either enhance or suppress the effects of FSI. This last point is crucial in CT studies using the deuteron, as will be described in Sec. 4.

2.2. PLC expansion effects

An important aspect of CT is that the PLC will expand as it moves through the nucleus because it is not an eigenstate of the Hamiltonian. The expansion time τ (or equivalently, coherence length l_c over which the hadron travels without evolving out of a PLC) is unknown, but there are several estimation methods in the literature. An early wavefunction expansion method gives $\tau \approx (E_h/m_h)\tau_0$, where E_h is the lab energy and $\tau_0 \approx 1$ fm is the characteristic hadron rest frame time [47], which shows that an increase in E_h allows a PLC to remain small for a longer time, thus increasing the suppression of FSIs. Holographic light front QCD (HLFQCD) techniques give a coherence length of $l_c \sim 2-3$ fm [48], more than sufficient to observe CT of the proton in deuterium, but we note that there are a few differences in analysis from the recent work of [3]. The naive parton model in which PLC constituents separate at the speed of light gives a transverse hadron size of $x_t \sim \tau \sim (E_h/m_h)^{-1}z$, where z is the longitudinal coordinate in the lab frame and E_h/m_h is the time dilation factor as before. This gives a coherence length estimate of $l_c \approx E_h/m_h$ fm for the PLC [9]. The quantum diffusion model is inspired by perturbative QCD (pQCD) and gives a coherence length $l_c \simeq \langle 1/(E_n - E_h) \rangle \simeq$ $2p_h\langle 1/(m_n^2-m_h^2)\rangle$, where m_n^2 is the invariant mass squared of the intermediate excited state nof the hadron [9]. The uncertainty principle gives a maximum coherence length, $l_c \sim \frac{1}{\Lambda M} \frac{p_h}{m_t}$, where ΔM is a characteristic excitation energy [49]. For small values of m_h , this matches the quantum diffusion method, $l_c \approx \frac{2p_h}{\Delta M^2}$ with $\Delta M^2 = m_n^2 - m_h^2$ and gives a complete loss of coherence for distances $l > l_c \sim 0.3 - 0.5$ fm $\times p_h$, with p_h measured in GeV/c. The smaller size of deuterium assists in minimizing expansion effects by minimizing PLC travel length prior to collision site exit.

3. Constraints on the color diffusion model from ${}^{12}C(e,e'p)$

The flat Q^2 -dependence of the recent Hall C CT measurement [31] is in agreement with standard Glauber calculations not including any CT effects. Calculations including CT effects using the color diffusion model [9,35] with standard parameters ($\Delta M^2 = 1 \text{ GeV}^2$) overshoot the data by a large amount for the highest Q^2 values [31,50]. It is possible, however, to modify the parameters in the color diffusion model to see which values would still be compatible with the measured Hall C data. These values could then be used to determine the optimal choice of kinematics to observe the onset of CT in deuterium (see Sec. 5, we find perpendicular kinematics to be the best choice).

In Glauber calculations using the quantum diffusion model, the total cross section parameter entering the Glauber profile function is replaced with a position dependent one. The cross section evolves from a reduced value (reflecting the small-sized color transparent configuration)




Figure 3. Transparencies resulting from RMSGA calculations with CT effects included in the quantum diffusion model. The dependence on the ΔM^2 parameter (x-axis) and ratio $\mathcal{R} = \frac{\sigma_{\rm eff}(z=0)}{\sigma^{\rm tot}}$ at the interaction point (y-axis) is shown. The four panels show the four measured Q^2 bins of the recent JLab experiment. Red dashed lines on the plots delineate the region (above the red line) compatible with the recent JLab data [31] at the 1σ level. White horizontal lines denote the $\mathcal{R}(\alpha=1)$ standard value.

to its normal value along a coherence length l_c , where the z coordinate is along the momentum of the ejected particle with z=0 at the point of interaction with the probe:

$$\sigma^{\text{tot}} \longrightarrow \sigma^{\text{eff}}(z) = \sigma^{\text{tot}} \left\{ \left[\frac{z}{l_c} + \frac{\langle n^2 k_t^2 \rangle}{Q^2} \left(1 - \frac{z}{l_c} \right) \right] \theta(l_c - z) + \theta(z - l_c) \right\} . \tag{2}$$

Here, $l_c=2p/\Delta M^2$, with p the momentum of the knocked out proton. For the parameters in $\sigma^{\rm eff}$ a standard assumption was $\Delta M^2=1.0~{\rm GeV}^2$, n=3 reflects the elementary constituents in the proton, and $k_t=0.35~{\rm GeV}/c$ the average transverse momentum of a quark inside a hadron [9]. One sees that the effective cross section is reduced at the interaction point z=0 to $\sigma_{\rm eff}(z=0)=\frac{\langle n^2k_t^2\rangle}{Q^2}\sigma^{\rm tot}$ and then rises linearly over the coherence length l_c to its standard value $\sigma^{\rm tot}$.

In order to map the quantum diffusion model parameter values that would still agree with the recent Hall C data, we perform calculations within the relativistic multiple scattering Glauber approximation (RMSGA) [50–52] with modified parameters: we take modified ΔM^2 values, which results in a modified coherence length, and we introduce a new modification parameter α such that

$$\mathcal{R} = \alpha \frac{\langle n^2 k_t^2 \rangle}{Q^2} \,, \tag{3}$$

and

$$\sigma^{\text{eff}}(z=0) = \mathcal{R}\sigma^{\text{tot}} = \alpha \frac{\langle n^2 k_t^2 \rangle}{Q^2} \sigma^{\text{tot}}$$
(4)

is now the modified cross section at the interaction point.

Figure 3 shows which parameters values would still be compatible with the recent Hall C data. It is clear that the highest Q^2 data provides the most stringent constraints and that coherence lengths corresponding with a ΔM^2 of 3 GeV² or more are required.

4. Probing proton CT in double scattering kinematics

The recent proton results of [31] reinvigorated the physics community and have prompted significant ongoing discussion as well as a re-examination of CT models. It is possible to investigate the effects of CT in the deuteron where the FSI and kinematics are well-understood, as described in Sec. 2. Double scattering accesses inter-nucleon distances on the order of 1-2 fm. Access to distances of this magnitude enables observations of any PLCs and would help constrain the PLC expansion rate which may be larger than expected due to the lack of recent experimental observation [49]. The larger the momentum of the spectator nucleon, the smaller the inter-nucleon distance and thus the shorter the distance between the production and re-scattering vertices leading to higher FSI effects. While the idea to explore CT in re-scattering (perpendicular) kinematics was proposed prior to the 12 GeV upgrade of JLab [49], updated predictions from Sargsian include the cross section from the recent JLab Hall C D(e, e'p)nmeasurement [43] and implementation of CT effects in the color diffusion model using the new constraint of $\Delta M^2 > 3 \text{ GeV}^2$ implied from the recent C(e, e'p) CT results. These new predictions indicate that there is parameter space to observe the signal of CT at $Q^2 > 8$ (GeV/c)² as shown in Fig. 4. It is important to note that the new calculations only change ΔM^2 and do not modify the ratio R. Here, the cross section ratio on the vertical axis is




Figure 4. Predictions of cross section ratios $R_D^{theory} = \frac{\sigma(p_r = 400 \text{MeV}/c)}{\sigma(p_r = 200 \text{MeV}/c)}$ in the D(e, e'p)n reaction at $\alpha_n \approx 1$ for various hypotheses of ΔM^2 [9] that are compatible with the recent Hall C results [31]. Upper: Predicted cross section ratios include the effect of CT with ΔM^2 between 2 and 3 GeV². Lower: the change of cross section ratio with various ΔM^2 values comparing to "no CT". Calculations performed in the GEA framework [40,46] using the AV18 deuterium wave function.

$$R_D^{theory}(Q^2) = \frac{\sigma(p_r = 400 \text{MeV}/c; Q^2) \downarrow}{\sigma(p_r = 200 \text{MeV}/c, Q^2) \uparrow'}$$
(5)

the ratio between high ($p_r = 400 \text{ MeV}/c$) and low ($p_r = 200 \text{ MeV}/c$) recoiling neutron momenta in the D(e, e'p)n reaction at perpendicular kinematics, i.e. the neutron recoil angle

 θ_{nq} is between 60 to 90° such that FSI effects deviate most from PWIA calculation as shown in Fig.2. More specifically, in this calculation we require the light cone momentum fraction of the nucleus carried by the recoil nucleon $\alpha_n \approx 1$, with

$$\alpha_n = \frac{(E_n - p_n \cos \theta_{nq})}{m_n} \tag{6}$$

expressed in the deuterium rest frame in terms of the final state spectator nucleon energy E_n , momentum p_n , mass m_n , and angle with the virtual photon θ_{nq} . The high p_r region is dominated by double scattering, the large increase in the cross section from PWIA prediction is shown in data from [39]. As Q^2 increases, we expect to see an increased probability of scattering from a PLC (and hence experiencing less FSI effects) so that the observed cross section decreases at high p_r , and increases at low p_r . As a result, the high-to-low ratio decreases as compared with the traditional Glauber calculations in the GEA framework [40].

As shown in Fig. 4, an increasing Q^2 -dependence of the FSIs when compared to "no CT" predictions made using the GEA could indicate a regime for the onset of CT. This allows us to experimentally explore proton CT in kinematics that were elusive or inaccessible in previous (e, e'p) experiments. Furthermore, comparing cross section ratios between data and theory relies less on the model dependence than comparing the absolute cross section to calculations directly, since some assumptions in the theory are canceled in the ratio.

5. A possible near-term Hall C experiment

With near term planned upgrades to the JLab beamline, it will be possible for Hall C to receive an incident 11 GeV electron beam energy. Hall C is uniquely capable of measuring high luminosity and high precision and has already been used in both color transparency and deuterium electro-disintegration experiments. This 11 GeV beam energy, together with the full momentum capabilities of the Hall C spectrometers, can extend the measurable Q^2 up to 17 $(\text{GeV}/c)^2$. Here, we discuss the possibility to measure the onset of CT in D(e,e'p)n by detecting the electron and proton in coincidence in Hall C from electron scattering on a deuterium target. Hall C is comprised of two spectrometer arms: the High Momentum Spectrometer (HMS) and the Super High Momentum Spectrometer (SHMS). The spectrometer pair is capable of detecting particles with momenta up to 7 GeV/c with a precision on the order of 0.1% (HMS), and 11 GeV/c with a precision on the order of 0.05% (SHMS). Note that the spectrometer momentum and angular range, not the beam energy, set the limit on our Q^2 coverage. Both spectrometers are equipped with standard particle detectors including time-of-flight scintillators for fast timing and triggering, drift chambers for precise particle tracking, and Cherenkov detectors and electromagnetic calorimeters for detailed particle identification.

In such an experiment, for each Q^2 setting, the HMS would detect the scattered electrons at the quasi-elastic peak, while the struck protons would be detected in the SHMS with high momentum resolution ($\sim 10~{\rm MeV/c}$). The SHMS momentum acceptance could be centered at $p_r = 200~{\rm MeV/c}$ which would enable coverage for both the low ($50-150~{\rm MeV/c}$) and high ($300-600~{\rm MeV/c}$) p_r regions simultaneously due to its large momentum acceptance. The experimental observable is the ratio of the measured (e,e'p) cross sections from those two momentum regions:

$$R_D^{exp}(Q^2) = \frac{\sigma(300 \le p_r \le 600 \text{ MeV}/c; Q^2) \downarrow}{\sigma(50 \le p_r \le 150 \text{ MeV}/c; Q^2) \uparrow}$$
(7)

As discussed previously, R_D^{exp} is expected to decrease with increasing Q^2 in the presence of PLCs. We also require the neutron recoil angle θ_{nq} of 60 to 90° to maximize FSI effects in both p_r regions so that the expected change in the ratio is maximized in the presence of CT. With

the standard Hall C spectrometer settings, the proposed measurement can overlap with the previous C(e, e'p) experiment in Q^2 requiring minimal beam time.

The missing mass is an essential quantity to reconstruct for all measured kinematics. Kinematics and rate studies done with the Hall C SIMC Monte-Carlo simulation package [53,54] show the reconstructed missing mass of the neutron with and without radiative effects as shown in Fig. 5. The estimated detectable rates rely on a cut on the reconstructed neutron mass.




Figure 5. The missing mass from the Hall C simulation software, SIMC, with (red) and without (blue) radiative effects at $Q^2 = 14 \, (\text{GeV}/c)^2$.

With a 25cm liquid deuterium target, this proposed experiment can measure the cross section ratio R_D at Q^2 from 8 to 15 $(\text{GeV}/c)^2$ in about 3 months of beam time with an approximate 3% statistical uncertainty at each Q^2 setting. While a higher beam energy would not increase the Q^2 coverage due to the spectrometer limitations, increasing the beam energy by 0.5 GeV would increase the rates by \sim 30% and could significantly reduce beam time and/or improve statistical precision.

6. Summary

We propose a realistic near-term experimental effort to search for the onset signal of CT using a deuterium target at JLab that could be accomplished with approximately 3 months of 11 GeV electron beam time. The experiment would probe a Q^2 range from 8–15 GeV 2 and would have an increased sensitivity as compared to previous electron-scattering experiments, allowing it to conclusively observe or rule out a signal for the onset of CT in the proton for this momentum range. The deuteron is advantageous for such a measurement because the CT onset signal is expected to be enhanced by measuring knockout reactions in kinematical regions already dominated by high FSIs and is described well in terms of the generalized eikonal approximation at high momentum. Double-scattering effects have a unique enhancement to the cross section ratio that is well-studied experimentally and has a clear dependence on the recoiling neutron angle. JLab Hall C is the first choice to perform such a measurement, but other experimental setups could be explored with upgrades to detector acceptances, luminosities and electron beam energy.

Author Contributions: All authors have contributed to this manuscript and have read and agreed to the published version of the manuscript.

Funding: This research was funded in part by Department of Energy grant number DE-AC05-06OR23177 under which the Jefferson Science Associates operates the Thomas Jefferson National Accelerator Facility.

SL is supported by Department of Energy, Office of Nuclear Physics, under contract number DE-AC02-05CH11231. CY is supported by the National Science Foundation MPS-Ascend Postdoctoral Research Fellowships under award number 2137604. JRW is supported by the LDRD programs of LBNL, the EIC Center at Jefferson Lab and by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under contract number DE-AC02-05CH11231. WC is supported by the National Science Foundation under Award No. 2111442. CB acknowledges support by Department of Energy, Office of Nuclear Physics under contract DE-SC0022007. MS's work is supported by the U.S. DOE Office of Nuclear Physics grant DE-FG02-01ER41172.

References

- 1. Bjorken, J.D.; Paschos, E.A. Inelastic Electron Proton and gamma Proton Scattering, and the Structure of the Nucleon. *Phys. Rev.* **1969**, 185, 1975–1982. doi:10.1103/PhysRev.185.1975.
- 2. Drell, S.D.; Yan, T.M. Connection of Elastic Electromagnetic Nucleon Form Factors at Large *Q*² and Deep Inelastic Structure Functions Near Threshold. *Phys. Rev. Lett.* **1970**, 24, 181–186. doi:10.1103/PhysRevLett.24.181.
- 3. Brodsky, S.J.; de Teramond, G.F. Onset of Color Transparency in Holographic Light-Front QCD. MDPI Physics 2022, 4, 633–646, [arXiv:hep-ph/2202.13283]. doi:10.3390/physics4020042.
- 4. Brodsky, S.J.; Mueller, A.H. Using Nuclei to Probe Hadronization in QCD. *Phys. Lett. B* **1988**, *206*, 685–690. doi:10.1016/0370-2693(88)90719-8.
- 5. Brodsky, S.J.; Frankfurt, L.; Gunion, J.F.; Mueller, A.H.; Strikman, M. Diffractive leptoproduction of vector mesons in QCD. *Phys. Rev. D* **1994**, *50*, 3134–3144, [hep-ph/9402283]. doi:10.1103/PhysRevD.50.3134.
- 6. Dutta, D.; Hafidi, K.; Strikman, M. Color Transparency: past, present and future. *Prog. Part. Nucl. Phys.* 2013, 69, 1–27, [arXiv:nucl-th/1211.2826]. doi:10.1016/j.ppnp.2012.11.001.
- 7. Brodsky, S.J. Supersymmetric and Other Novel Features of Hadron Physics from Light-Front Holography. 24th Workshop on What Comes Beyond the Standard Models?, 2021, [arXiv:hep-ph/2112.02453].
- 8. Mueller, A.H. COLOR TRANSPARENCY AND NUCLEAR SHADOWING. Workshop on Nuclear and Particle Physics on the Light Cone, 1988.
- 9. Farrar, G.R.; Liu, H.; Frankfurt, L.L.; Strikman, M.I. Transparency in Nuclear Quasiexclusive Processes with Large Momentum Transfer. *Phys. Rev. Lett.* **1988**, *61*, 686–689. doi:10.1103/PhysRevLett.61.686.
- 10. Perkins, D. Phil. Mag. 1955, 46, 1146.
- 11. Clasie, B.; et al. Measurement of Nuclear Transparency for the $A(e, e'\pi^+)$ Reaction. *Phys. Rev. Lett.* **2007**, 99, 242502, [arXiv:nuclex/0707.1481]. doi:10.1103/PhysRevLett.99.242502.
- 12. El Fassi, L.; et al. Evidence for the onset of color transparency in ρ^0 electroproduction off nuclei. *Phys. Lett. B* **2012**, *712*, 326–330, [arXiv:nucl-ex/1201.2735]. doi:10.1016/j.physletb.2012.05.019.
- 13. Frankfurt, L.; Miller, G.A.; Strikman, M. Color Transparency in Semi-Inclusive Electroproduction of rho Mesons. *Phys. Rev. C* **2008**, 78, 015208, [arXiv:nucl-th/0803.4012]. doi:10.1103/PhysRevC.78.015208.
- 14. Gallmeister, K.; Kaskulov, M.; Mosel, U. Color transparency in hadronic attenuation of ρ^0 mesons. *Phys. Rev. C* **2011**, *83*, 015201, [arXiv:hep-ph/1007.1141]. doi:10.1103/PhysRevC.83.015201.
- 15. Cosyn, W.; Ryckebusch, J. Nuclear ρ meson transparency in a relativistic Glauber model. *Phys. Rev. C* **2013**, *87*, 064608, [arXiv:nucl-th/1301.1904]. doi:10.1103/PhysRevC.87.064608.
- 16. Blaettel, B.; Baym, G.; Frankfurt, L.L.; Strikman, M. How transparent are hadrons to pions? *Phys. Rev. Lett.* **1993**, *70*, 896–899. doi:10.1103/PhysRevLett.70.896.
- 17. Carroll, A.S.; et al. Nuclear Transparency to Large Angle pp Elastic Scattering. Phys. Rev. Lett. 1988, 61, 1698–1701. doi:10.1103/PhysRevLett.61.1698.
- 18. Mardor, I.; et al. Nuclear transparency in large momentum transfer quasielastic scattering. *Phys. Rev. Lett.* **1998**, *81*, 5085–5088. doi:10.1103/PhysRevLett.81.5085.
- 19. Leksanov, A.; et al. Energy dependence of nuclear transparency in C(p, 2p) scattering. *Phys. Rev. Lett.* **2001**, 87, 212301, [arXiv:hep-ex/hep-ex/0104039]. doi:10.1103/PhysRevLett.87.212301.
- 20. Aclander, J.; et al. Nuclear transparency in $90^{\circ}_{\text{c.m.}}$ quasielastic A(p,2p) reactions. *Phys. Rev. C* **2004**, 70, 015208, [arXiv:nucl-ex/nucl-ex/0405025]. doi:10.1103/PhysRevC.70.015208.
- 21. Barbieri, C.; Lapikas, L. Effects of rescattering in (e,e-prime p) reactions within a semiclassical model. *Phys. Rev. C* **2004**, 70, 054612, [nucl-th/0409032]. doi:10.1103/PhysRevC.70.054612.
- 22. Rohe, D.; et al. Nuclear transparency from quasielastic C-12 (e, e-prime p). *Phys. Rev. C* **2005**, 72, 054602, [nucl-ex/0506007]. doi:10.1103/PhysRevC.72.054602.
- 23. Kundu, B.; Samuelsson, J.; Jain, P.; Ralston, J.P. Perturbative color transparency in electroproduction experiments. *Phys. Rev. D* **2000**, 62, 113009, [hep-ph/9812506]. doi:10.1103/PhysRevD.62.113009.

- 24. Ralston, J.P.; Pire, B. Quantum chromotransparency. Phys. Rev. Lett. 1990, 65, 2343–2346. doi:10.1103/PhysRevLett.65.2343.
- 25. Brodsky, S.J.; Schmidt, I.; de Téramond, G.F. Nuclear-bound quarkonium. Phys. Rev. Lett. 1990, 64, 1011–1014. doi:10.1103/PhysRevLett.64.1011.
- 26. Makins, N.; et al. Momentum transfer dependence of nuclear transparency from the quasielastic ¹²C(e,e'p) reaction. *Phys. Rev. Lett.* **1994**, 72, 1986–1989. doi:10.1103/PhysRevLett.72.1986.
- 27. O'Neill, T.G.; et al. *A*-dependence of nuclear transparency in quasielastic A(e, e'p) at high Q^2 . *Phys. Lett. B* **1995**, 351, 87–92, [arXiv:hep-ph/hep-ph/9408260]. doi:10.1016/0370-2693(95)00362-O.
- 28. Abbott, D.; et al. Quasifree (*e*, *e*'*p*) reactions and proton propagation in nuclei. *Phys. Rev. Lett.* **1998**, *80*, 5072–5076. doi:10.1103/PhysRevLett.80.5072.
- 29. Garrow, K.; et al. Nuclear transparency from quasielastic A(e,e'p) reactions up to $Q^2 = 8.1 (\text{GeV}/c)^2$. *Phys. Rev. C* **2002**, *66*, 044613, [arXiv:hep-ex/hep-ex/0109027]. doi:10.1103/PhysRevC.66.044613.
- 30. Frullani, S.; Mougey, J. Single-Particle Properties of Nuclei through (e, e'p) Reactions. Advances in Nuclear Physics 1984, 14.
- 31. Bhetuwal, D.; Matter, J.; Szumila-Vance, H.; Kabir, M.L.; Dutta, D.; Ent, R.; et al. Ruling out Color Transparency in Quasielastic 12 C(e, e'p) up to Q^2 of 14.2 (GeV/c) 2 . *Phys. Rev. Lett.* **2021**, 126, 082301. doi:10.1103/PhysRevLett.126.082301.
- 32. Frankfurt, L.; Miller, G.A.; Strikman, M. Precocious dominance of point like configurations in hadronic form-factors. *Nucl. Phys. A* **1993**, 555, 752–764. doi:10.1016/0375-9474(93)90504-Q.
- 33. Capel, P. The eikonal model of reactions involving exotic nuclei; Roy Glauber's legacy in today's nuclear physics. *SciPost Phys. Proc.* **2020**, *3*, 017, [arXiv:nucl-th/1911.11446]. doi:10.21468/SciPostPhysProc.3.017.
- 34. Sargsian, M.M.; Abrahamyan, T.V.; Strikman, M.I.; Frankfurt, L.L. Exclusive electrodisintegration of He-3 at high Q**2. I. Generalized eikonal approximation. *Phys. Rev. C* **2005**, *71*, 044614, [nucl-th/0406020]. doi:10.1103/PhysRevC.71.044614.
- 35. Frankfurt, L.L.; Greenberg, W.R.; Miller, G.A.; Sargsian, M.M.; Strikman, M.I. Color transparency effects in electron deuteron interactions at intermediate Q**2. Z. Phys. A 1995, 352, 97–113, [nucl-th/9501009]. doi:10.1007/BF01292764.
- 36. Yero, C. Cross Section Measurements of Deuteron Electro-Disintegration at Very High Recoil Momenta and Large 4-Momentum Transfers (*Q*²). PhD thesis, Florida Intl. U., Florida Intl. U., 2020, [arXiv:nucl-ex/2009.11343]. doi:10.25148/etd.FIDC009154.
- 37. Sargsian, M.M.; Abrahamyan, T.V.; Strikman, M.I.; Frankfurt, L.L. Exclusive electro-disintegration of ³He at high Q2. II. Decay function formalism. *Phys. Rev. C* **2005**, *71*, 044615, [arXiv:nucl-th/0501018]. doi:10.1103/PhysRevC.71.044615.
- 38. Boeglin, W.U.; et al. Probing the high momentum component of the deuteron at high *Q*². *Phys. Rev. Lett.* **2011**, 107, 262501, [arXiv:nucl-ex/1106.0275]. doi:10.1103/PhysRevLett.107.262501.
- 39. Egiyan, K.S.; et al. Experimental Study of Exclusive ${}^{2}H(e,e'p)n$ Reaction Mechanisms at High Q^{2} . *Phys. Rev. Lett.* **2007**, 98, 262502. doi:10.1103/PhysRevLett.98.262502.
- Frankfurt, L.L.; Sargsian, M.M.; Strikman, M.I. Feynman graphs and generalized eikonal approach to high energy knock-out processes. Phys. Rev. C 1997, 56, 1124.
- 41. Lacombe, M.; Loiseau, B.; Richard, J.M.; Mau, R.V.; Côté, J.; Pirès, P.; de Tourreil, R. Parametrization of the Paris N-N potential.
- Phys. Rev. C 1980, 21, 861–873. doi:10.1103/PhysRevC.21.861.
 Machleidt, R. High-precision, charge-dependent Bonn nucleon-nucleon potential. Phys. Rev. C 2001, 63, 024001. doi:10.1103/PhysRevC.63.02400
- 43. Yero, C.; et al. Probing the Deuteron at Very Large Internal Momenta. Phys. Rev. Lett. 2020, 125, 262501.
- 44. Laget, J. The electro-disintegration of few body systems revisited. Physics Letters B 2005, 609, 49-56.
- 45. Wiringa, R.B.; Stoks, V.G.J.; Schiavilla, R. Accurate nucleon-nucleon potential with charge-independence breaking. *Phys. Rev. C* **1995**, 51, 38–51. doi:10.1103/PhysRevC.51.38.
- 46. Sargsian, M.M. Large Q² Electrodisintegration of the Deuteron in Virtual Nucleon Approximation. *Phys. Rev. C* **2010**, *82*, 014612, [arXiv:nucl-th/0910.2016]. doi:10.1103/PhysRevC.82.014612.
- 47. Frankfurt, L.; Greenberg, W.R.; Miller, G.A.; Strikman, M. Sum rule description of color transparency. *Phys. Rev. C* **1992**, 46, 2547–2553, [nucl-th/9211002]. doi:10.1103/PhysRevC.46.2547.
- 48. Caplow-Munro, O.; Miller, G.A. Color transparency and the proton form factor: Evidence for the Feynman mechanism. *Phys. Rev. C* **2021**, *104*, L012201, [arXiv:nucl-th/2104.11168]. doi:10.1103/PhysRevC.104.L012201.
- 49. Sargsian, M.M.; Arrington, J.; Bertozzi, W.; Boeglin, W.; Carlson, C.E.; Day, D.B.; Frankfurt, L.L.; Egiyan, K.; Ent, R.; Gilad, S.; et al. Hadrons in the nuclear medium. *Journal of Physics G: Nuclear and Particle Physics* **2003**, 29, R1–R45. doi:10.1088/0954-3899/29/3/201.
- 50. Cosyn, W.; Ryckebusch, J. Nuclear C(e, e'p) Transparencies in a Relativistic Glauber Model. *MDPI Physics* **2022**, *4*, 672–676, [arXiv:nucl-th/2206.03374]. doi:10.3390/physics4020045.
- 51. Ryckebusch, J.; Debruyne, D.; Lava, P.; Janssen, S.; Van Overmeire, B.; Van Cauteren, T. Relativistic formulation of Glauber theory for A(e,e'p) reactions. *Nucl. Phys.* **2003**, *A728*, 226–250, [arXiv:nucl-th/nucl-th/0305066]. doi:10.1016/j.nuclphysa.2003.08.022.
- 52. Cosyn, W.; Martinez, M.C.; Ryckebusch, J.; Van Overmeire, B. Nuclear transparencies from photoinduced pion production. *Phys. Rev.* C 2006, 74, 062201, [arXiv:nucl-th/nucl-th/0701029]. doi:10.1103/PhysRevC.74.062201.
- 53. SIMC Monte Carlo. Available at https://hallcweb.jlab.org/wiki/index.php/SIMC_Monte_Carlo.
- 54. Dutta, D.; et al. Separated spectral functions for the quasifree C-12 (e, e-prime p) reaction. *Phys. Rev. C* 2000, 61, 061602. doi:10.1103/PhysRevC.61.061602.