Target mass corrections in lepton-nucleus DIS: theory and applications to nuclear PDFs

R. Ruiz \bigcirc , ^{*a*} K.F. Muzakka \bigcirc , ^{*b,c*} C. Leger, ^{*d*} P. Risse \bigcirc , ^{*b*} A. Accardi \bigcirc , ^{*e,f*} P. Duwentäster \bigcirc , ^{*b,g,h*} T.J. Hobbs \bigcirc , ^{*i*} T. Ježo \bigcirc , ^{*b*} C. Keppel \bigcirc , ^{*e*} M. Klasen \bigcirc , ^{*b,j*} K. Kovařík \bigcirc , ^{*b*} A. Kusina \bigcirc , ^{*a*} J.G. Morfín \bigcirc , ^{*k*} F.I. Olness \bigcirc , ^{*l*} J.F. Owens, \bigcirc , ^{*m*} I. Schienbein \bigcirc , ^{*d*} J.Y. Yu. \bigcirc ^{*d*}

^aInstitute of Nuclear Physics Polish Academy of Sciences, PL-31342 Krakow, Poland

CNRS/IN2P3, 53 avenue des Martyrs, 38026 Grenoble, France

ABSTRACT: Motivated by the wide range of kinematics covered by current and planned deep-inelastic scattering (DIS) facilities, we revisit the formalism, practical implementation, and numerical impact of target mass corrections (TMCs) for DIS on unpolarized nuclear targets. An important aspect is that we only use nuclear and later partonic degrees of freedom, carefully avoiding a picture of the nucleus in terms of nucleons. After establishing that formulae used for individual nucleon targets (p, n), derived in the Operator Product Expansion (OPE) formalism, are indeed applicable to nuclear targets, we rewrite expressions for nuclear TMCs in terms of re-scaled (or averaged) kinematic variables. As a consequence, we find a representation for nuclear TMCs that is approximately independent of the nuclear target. We go on to construct a single-parameter fit for all nuclear targets that is in good numerical agreement with full computations of TMCs. We discuss in detail qualitative and quantitative differences between nuclear TMCs built in the OPE and the parton model formalisms, as well as give numerical predictions for current and future facilities.

KEYWORDS: DIS, Structure Functions, Target Mass Corrections, OPE, nuclear PDFs

^bInstitut für Theoretische Physik, Westfälische Wilhelms-Universität Münster,

Wilhelm-Klemm-Straße 9, D-48149 Münster, Germany

^cInstitut für Energie- und Klimaforschung, Forschungszentrum Jülich GmbH, 52425 Jülich, Germany

^dLaboratoire de Physique Subatomique et de Cosmologie, Université Grenoble-Alpes,

^e Jefferson Lab, Newport News, VA 23606, U.S.A.

^f Hampton University, Hampton, VA 23668, USA

^g University of Jyväskylä, Department of Physics, P.O. Box 35, FI-40014 University of Jyväskylä, Finland

^hHelsinki Institute of Physics, P.O. Box 64, FI-00014 University of Helsinki, Finland

ⁱHigh Energy Physics Division, Argonne National Laboratory, Argonne, Illinois 60439, USA

^jSchool of Physics, The University of New South Wales, Sydney NSW 2052, Australia

^k Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA

¹Department of Physics, Southern Methodist University, Dallas, TX 75275-0175, U.S.A.

^mDepartment of Physics, Florida State University, Tallahassee, Florida 32306-4350, USA

Contents

1	Intr	roduction	3		
2	Kin	lematics of lepton-nucleus DIS	6		
	2.1	Definitions of kinematic variables	6		
	2.2	Leptonic and hadronic tensors in DIS from experimentally observable kinematics	7		
3	Nuclear structure functions in the OPE		10		
	3.1	Light-cone dominance of nuclear DIS	10		
	3.2	Structure functions in the OPE	12		
	3.3	Master formula for structure functions with TMCs in ℓA DIS	16		
4	\mathbf{Res}	scaling	19		
	4.1	Nuclear and Nucleon Kinematics	20		
	4.2	Rescaled Structure Functions	21		
	4.3	Impact of Quark Masses	22		
5	Par	ton model	23		
	5.1	Nuclear DGLAP Evolution	24		
5 6	5.2	Relation to the OPE	25		
	5.3	Rescaling	25		
	5.4	Kinematic \boldsymbol{W} Cut	28		
	5.5	nPDFs for $x_N \ge 1$	30		
	5.6	Threshold problem and higher twist contributions	32		
6	$\mathrm{Th}\epsilon$	e parton model with quark and hadron masses	33		
	6.1	Helicity Formulation of the Parton Model	33		
	6.2	.2 Helicity Decomposition			
	6.3	B Boost Invariant Polarizations			
	6.4	Relation to Master Formulas			
	6.5	Relation of Parton Model to OPE	36		
7	Nu	merical results	37		
	7.1	Proton Structure Functions with TMC	39		
7	7.2	Nuclear Structure Functions with TMCs	40		
	7.3	Parameterizing $F_i^{\text{TMC}}/F_i^{\text{Leading-TMC}}$	48		
	7.4	Reduced Cross Sections with TMCs for Nuclei	51		
8	Cor	nclusions	53		
	Appe	endix A Nuclear structure functions with TMCs in the OPE	55		
	Ap	opendix A.1 Preliminaries and the inclusive DIS cross section formula	55		
	Ap	opendix A.2 The DIS matrix element and inclusive hadronic tensor	56		
	Ap	opendix A.3 Nuclear structure functions from the OPE I: organization	60		
	Ap	ppendix A.4 Nuclear structure functions from the OPE II: structure-function mixing	72		

Appendix B Derivation of the Full/Leading TMC Parameterization

84

1. Introduction

Deep-inelastic scattering (DIS) of high-energy leptons off hadrons is a key process for studying the structure of hadrons in terms of their quark and gluon degrees of freedom [1-4]. For both nucleons [5-10] and nuclei [11-28], present data from electron and neutrino DIS form the backbone of modern determinations of parton distribution functions (PDFs), which describe and encode the internal structure of hadrons. This process will again be at the forefront of the future Electron-Ion Collider (EIC) [29, 30], where DIS off nucleons and various nuclear targets will be studied with high precision over a wide kinematic range. Similar endeavors are also in discussion for the proposed Forward Physics Facility [31]. Additionally, high-precision fixed-target measurements of JLab can provide complementary information [32]. Beyond this, DIS also plays an important role at current and future accelerator-based neutrino oscillation facilities such as DUNE and the Short-Baseline Neutrino Program at Fermilab [33–35], where experiments operate at somewhat lower energies compared to the EIC, and where the (quasi)elastic and resonant processes must also be taken into account. (Here, precise control over the contributions from each of these processes is essential, and duality-based arguments for the scaling of DIS cross sections at higher energies into the resonance region can play an important role [36, 37].) Due to the relative weakness of the neutrino-nucleon effective interaction, neutrino experiments typically use nuclear targets to enhance event rates. Consequently, charged lepton-nucleus and neutrino-nucleus interactions need to be precisely understood, particularly the differences from *nucleon* interactions, to reach the ambitious physics goals of future experiments.

Along these lines, there are two major theoretical approaches to lepton-*nucleon* DIS that are standard material in textbooks and review articles; both have been employed to include the effects of target nucleons having nonzero masses. One approach is based on the operator product expansion (OPE) [38–40]. The other is the QCD-improved parton model rigorously derived from first principles of QCD in the context of the collinear factorization theorems [41–48]. The OPE was first employed to derive target mass corrections (TMCs) to structure functions in DIS at leading order (LO) in QCD in the seminal paper by Georgi and Politzer in 1976 [49, 50]. In the same year Barbieri, et al. [51, 52] derived TMCs at LO including effects arising from non-zero quark masses. Later, the original work of Georgi and Politzer was extended to next-to-leading order (NLO) in QCD for the structure functions $W_{1,2}$ in the context of local duality in electro-production and using off-shell regularization [53]. More recently, Kretzer and Reno presented TMCs for charged current (CC) and weak neutral current (NC) neutrino-nucleon DIS, including NLO QCD corrections and heavy quark mass effects using modern conventions [54].

In the context of the parton model, Ellis, Furmanski, and Petronzio derived TMCs at LO in QCD in the framework of a non-collinear version of the parton model that includes the effects of partonic transverse momentum (k_T) [55]. Agreement with the OPE-based results [51, 52] was found at leading power, thereby demonstrating the equivalence of the OPE at LO in QCD to a non-collinear parton approach, where the incident parton is on-shell but not collinear with its parent nucleon. We note also that TMC prescriptions based on the OPE and the factorization approach of Ellis, Furmanski, and Petronzio were similarly compared for semi-inclusive processes in Ref. [56]. In the collinear parton model ($k_T = 0$), TMCs have been accounted for in the Aivazis-Collins-Olness-Tung (ACOT) formalism [57, 58], which is rigorously based on the factorization theorem including heavy quark masses [46]. Furthermore, the full QCD framework for the evaluation of tau-neutrino deep-inelastic charged current cross sections, including NLO corrections, charm production, tau mass threshold, and target mass effects in the collinear approximation, has also been presented [59].

An earlier review [60] of TMCs for nucleons in unpolarized DIS culminated in a set of so-called master formulas [54], which are now updated and presented below for *nuclei* in Eq. (4.15). This set of equations is quite remarkable, organizing the rather complicated expressions in the aforementioned OPE-based papers into a simple, easy-to-use, and modular form, valid at all orders in perturbation QCD while still taking into account quark masses. In principle, TMCs derived for DIS off nucleons [49, 50, 54, 60, 61] apply also to DIS off nuclear targets since the OPE is (in principle) independent of target states. However, previous discussions [60] do not address the subtle distinctions between a nucleon and nuclear target: the fact that established notation does not consistently distinguish between nucleons and nuclei in their respective kinematics, the fact that the spin of a generic nucleus can be different from 1/2, and the fact that the master equations in Eq. (4.15) are expressed in terms of "averaged nucleon mass" has led to the question, why nucleon-like expressions could possibly be valid for nuclear targets?

With an abundance of recent nuclear data from both electron and neutrino beams, and with the EIC and 3rd-generation long-baseline oscillations experiments on the horizon, there is a compelling need to rigorously revisit the derivation of TMCs to structure functions in DIS with a particular focus on the *nuclear* case. One of the main goals for this article is to properly demonstrate the validity of the TMC master equations for the case of massive nuclear targets within the framework of the OPE. Compared to past works on the subject [54, 60], here we present the derivation of TMCs from the OPE in much greater detail. We hope that this material will be useful for students and researchers looking for a modern, in-depth discussion of the many technicalities that are frequently omitted in the literature.

In the following, we discuss and contrast the results for TMCs obtained in both the OPE and the collinear parton model; throughout this review, we consider them on equal footing. As discussed above, the collinear parton model is rigorously based on factorization theorems. These provide field-theoretic definitions of PDFs and make statements about the error of the factorization approximation, which is generally inversely proportional to a positive power of a hard scale of the process.¹ It is generally believed that collinear factorization remains valid in lepton-nucleus and proton-nucleus collisions, possibly with nuclear-enhanced higher twist terms [62]. However, the literature on factorization in the nuclear case is sparse and we consider this a working assumption. Moreover, the non-collinear parton model (where the TMCs were shown to be equivalent to the OPE results at leading power) is not covered by the factorization proofs cited above. While the application of the OPE to DIS is widely accepted, proofs of the OPE have only been presented for simple scalar models [63, 64]. The question as to whether this result can be extended to QCD in general remains open to the best of our knowledge. Exploring these more theoretical questions is interesting and relevant but beyond the scope of this article. Here, we take the collinear parton model and the OPE for granted, and explore extensively what happens when we transition from nucleons to nuclei.

¹Such power-suppressed terms in DIS structure functions are generally called "higher twist" terms. In the context of the OPE we define the twist $\tau = d - s$ of a field operator $\hat{\mathcal{O}}$ in the canonical fashion, i.e., the dimension d of $\hat{\mathcal{O}}$ (in the usual power-counting sense) minus its spin s (the number of un-contracted Lorentz indices).

$\ell_1(k_1) \ell_1(k_2) \\ \bigvee V(q)$				
	$\overset{X(p_X)}{\blacksquare}$			
Kinematic variable	Description			
$\nu_A = \frac{q \cdot p_A}{M_A} \stackrel{lab}{=} E_{\ell_1} - E_{\ell_2}$	Lepton energy loss in the laboratory (nucleon rest) frame			
$y_A = \frac{q \cdot p_A}{k_1 \cdot p_A} \stackrel{lab}{=} \frac{\nu_A}{E_{\ell_1}}$	Inelasticity $y_A \in [0, 1]$			
$Q^2 = -q^2 > 0$	Boson squared momentum transfer			
$x_A = \frac{Q^2}{2p_A \cdot q} = \frac{Q^2}{2M_A \nu_A}$	Bjorken x_A with $x_A \in [0, 1]$			
$W^{2} = (p_{A} + q)^{2} = M_{A}^{2} + Q^{2} \frac{1 - x_{A}}{x_{A}}$	Mass squared of the recoil system			
$s = (k_1 + p_A)^2 = \frac{Q^2}{x_A y_A} + M_A^2 + m_\ell^2$	Center of Mass System (CMS) energy squared			

Figure 1.1: We consider the basic charged current $(V=W^{\pm})$ or neutral current $(V=\{\gamma, Z\})$ lepton-nucleus DIS process $\ell_1(k_1) + A(p_A) \rightarrow \ell_2(k_2) + X(p_X)$ where the lepton can be a charged lepton (electron, muon) or a neutrino ν_e, ν_μ, ν_τ . The 4-momentum of the exchange boson is denoted $q = k_1 - k_2$, and $p_A^2 = M_A^2$.

The starting point of our analysis is to consider the full nucleus as our target and apply only general symmetry principles, e.g., Lorentz invariance, in deriving nuclear structure functions and their TMCs. This means that until Sec. 4 there is no reference to (or dependence on) nucleon degrees of freedom. Furthermore, until Sec. 5 there is no reference to (or dependence on) partonic degrees of freedom.² To do this, we first outline in Sec. 2 key kinematic relations and definitions in DIS of a lepton off a nucleus, $\ell_1(k_1) + A(p_A) \rightarrow \ell_2(k_2) + X(p_X)$. As depicted in Fig. 1.1, A is a nuclear target with mass number A, ℓ denotes either a charged lepton or neutrino, and X represents all final-state hadrons. In Sec. 3, we discuss precisely the criteria for light-cone dominance in nuclear DIS, and then present in Sec. 3.3 a formula for nuclear TMCs. The result is analogous to the nucleon case [60], but expressed in terms of the nuclear scaling variable x_A and the mass of the nucleus M_A .

In Sec. 4, we go on to perform a rescaling in order to express our formula for nuclear TMCs to the more familiar averaged nucleon quantities $x_N = Ax_A$ and $M_N = M_A/A$. This is the key step that

 $^{^{2}}$ Explicitly, until Sec. 5 we consider operators and matrix elements derived from quark, antiquark, and gluon fields but do not identify these as partons, nor identify structure functions as combinations of PDFs, i.e., the parton model.

allows us to compare structure functions across different nuclei, including the proton, in a meaningful way. As an important consequence of rescaling, we obtain an alternative formula for nuclear TMCs that is universally applicable to all nuclei. In Sec. 5 we introduce the QCD-improved parton model, both before and after rescaling. We show that rescaling at the hadronic level and in the parton model are consistent and we comprehensively discuss the relationship between nuclear structure functions and nuclear PDFs (nPDFs). We believe that the discussions in Secs. 4 and 5, while relatively straight forward, are original and have never been presented in full detail. It is key for a proper theoretical definition of nuclear structure functions and nPDFs as they are intuitively used in the literature.

In Sec. 6, we derive nuclear TMCs in the ACOT light cone formalism, which has been used in past nCTEQ analyses [16, 17, 19–21, 65, 66], and compare them to the OPE results. Having provided rigorous theoretical definitions for the physical observables, we perform numerical studies in Sec. 7. This includes a comparison to a selection of data, as well as cross section predictions for current and future DIS experiments. We provide a parameterization that is accurate at the sub-percent level and can be used with any underlying set of nuclear structure functions available in the massless parton model to obtain the full TMCs for any nucleus in a simple way. We hope that this parameterization will be useful for the community working on hadron structure. Finally, in Sec. 8 we highlight the key observations of this analysis and conclude.

A detailed derivation of TMCs at twist $\tau = 2$ for nuclear structure functions using the OPE is provided in Appendix A. Finally, technical details on our parameterization of the TMCs have been relegated to Appendix B.

2. Kinematics of lepton-nucleus DIS

Before proceeding with an in-depth treatment of nuclear structure functions, we layout in this section the definitions, notation, and baseline assumptions used throughout this work. (However, assumptions related to light-cone dominance are discussed in Sec. 3.1.) While much of the following is standard material for modern textbooks, many conventions are used in the literature. Therefore, this section serves (a) to fix the notation and conventions that we use, and (b) to make the work both selfconsistent and self-contained. We start with definitions of kinematic variables in Sec. 2.1 and move onto the definitions of leptonic and hadronic tensors for DIS in Sec. 2.2.

2.1. Definitions of kinematic variables

Throughout this work we consider the basic charged-current (CC) or neutral-current (NC) leptonnucleus DIS process involving a high-energy lepton $\ell_1(k_1)$ scattering off an unpolarized nuclear target $A(p_A)$ of mass M_A , with an outgoing lepton $\ell_2(k_2)$ in association with the inclusive hadronic remnant $X(p_X)$ of mass $W = \sqrt{p_X^2}$. As sketched in Fig. 1.1, this is given by the hadron-level expression

$$\ell_1(k_1) + A(p_A) \to \ell_2(k_2) + X(p_X)$$
 (2.1)

We consider that the incoming lepton can be a light charged lepton $\{e^{\pm}, \mu^{\pm}\}$, a neutrino $\{\nu_e, \nu_\mu, \nu_\tau\}$, or an antineutrino $\{\overline{\nu_e}, \overline{\nu_\mu}, \overline{\nu_\tau}\}$ in the Standard Model. For clarity, we work in the limit that all incoming leptons are massless but allow that the outgoing lepton may instead be a τ^{\pm} lepton. We denote the atomic and mass numbers of nucleus A by (Z, A), respectively. $X(p_X)$ represents the fragmentation of A. It is an *n*-body final state with net quantum numbers corresponding to a colorless state carrying the same spin statistic, QED charge, and weak isospin charge as A, modulo differences between ℓ_1 and ℓ_2 . The momentum of X can be parameterized by $p_X = \sum_{i=1}^n p_{X_i}$. We work at lowest order in the electroweak theory, in the so-called "one-boson-exchange" approximation. Under this assumption, the leptonic system $(\ell_2\ell_1)$ probes the hadronic system via the exchange of a single, electroweak boson V ($V \in \{\gamma, Z\}$ or $V = W^{\pm}$) with time-like momentum $q = k_1 - k_2 = p_X - p_A$. The kinematic invariants and scaling variables of DIS are given and defined by

$$k_1^2 = m_{\ell_1}^2, \qquad k_2^2 = m_{\ell_2}^2, \qquad p_A^2 = M_A^2 > 0, \qquad Q^2 = -q^2 > 0, \qquad s = (k_1 + p_A)^2,$$
 (2.2a)

$$x_A = \frac{Q^2}{2p_A \cdot q}, \qquad \nu_A = \frac{p_A \cdot q}{M_A}, \qquad y_A = \frac{p_A \cdot q}{p_A \cdot k}, \tag{2.2b}$$

$$W^{2} = p_{X}^{2} = (p_{A} + q)^{2} = M_{A}^{2} - Q^{2} + 2\nu_{A}M_{A} = M_{A}^{2} + Q^{2}\frac{1 - x_{A}}{x_{A}} \ge M_{A}^{2}.$$
 (2.2c)

We generally neglect the lepton masses $\{m_{\ell_1}, m_{\ell_2}\}$ except for the tau-lepton. The inelasticity variable (y_A) and the Bjorken scaling variable (x_A) take the values $x_A, y_A \in [0, 1]$, with $x_A = 1$ corresponding to the elastic limit where $W^2 = M_A^2$.

For later use with TMCs to structure functions (Secs. 3.3 and 4), we also define the quantity

$$r_A = \sqrt{1 + 4x_A^2 M_A^2/Q^2} = 1 + \frac{2x_A^2 M_A^2}{Q^2} + \mathcal{O}\left(\frac{4x_A^4 M_A^4}{Q^4}\right), \qquad (2.3)$$

as well as the so-called Nachtmann variable [67]

$$\xi_A = \frac{2}{1+r_A} x_A \equiv R_M x_A$$
, where $R_M = \frac{2}{1+r_A}$. (2.4)

 R_M is the target-mass dependent factor relating the Bjorken scaling variable x_A to the Nachtmann scaling variable ξ_A . Note that in the limit that $(M_A/Q) \to 0$ or that $x_A \to 0$, we have $r_A \to 1$, $R_M \to 1$, and $\xi_A \to x_A$ since $\xi_A = x_A(1 - x_A^2 M_A^2/Q^2) + \mathcal{O}(x_A^4 M_A^4/Q^4)$.

The difference between ξ_A and x_A is illustrated in Fig. 2.1, where ξ_A is plotted as a function of x_A for a selection of exchange-boson virtualities Q with $M_A = M_{\text{proton}}$. Fig. 2.1(a) displays the full $\{\xi_A, x\}$ plane. Fig. 2.1(b) highlights the large x region, where the difference between ξ_A and x is more pronounced. Qualitatively, x_A and ξ_A are largely indistinguishable for $x_A \leq 0.4-0.5$ over a large range of Q and M_A . Quantitatively, at around $x_A = 0.8$, one has approximately $\xi_A \approx 0.63$ (0.71) [0.76] for Q = 1.3 (2) [3] GeV. For the same Q values, one has approximately $\xi_A \approx 0.76$ (0.78) [0.84] at around $x_A = 0.9$.

2.2. Leptonic and hadronic tensors in DIS from experimentally observable kinematics

Deeply inelastic scattering is a powerful, elucidating probe of the internal structure of the proton and nuclei, i.e., hadrons. The ability stems from the fact that under strong but general assumptions one can write the cross section for any sufficiently inclusive DIS process $\ell_1 + A \xrightarrow{V} \ell_2 + X$ as a combination of: (a) leptonic and hadronic kinematics, which can be measured, and (b) hadronic structure functions, which parameterize the internal dynamics of A. This discussion is based solely on kinematics and symmetries, including Lorentz symmetry. The only approximation of consequence that is made when constructing structure functions from experimentally observable kinematics is the one where DIS is mediated by the exchange of only one electroweak boson. (Relaxing this has been explored elsewhere [68–71].) Apart from this assumption, expressions derived from kinematics are completely general: they take into account all twist contributions, at all orders in the strong coupling constant, and all hadron- and quark- mass effects. And in particular, the parton model does not need to be



Figure 2.1: (a) The Nachtmann scaling variable $\xi_A(x_A, Q)$ as a function of the Bjorken scaling variable x_A for a selection of Q values [GeV], as indicated in the legend, with $M_A = M_{\text{proton}}$. (b) Same as (a) but for large x_A .

invoked.

Schematically, the differential DIS cross section $d\sigma$ is proportional to the product of a leptonic tensor $L^{\mu\nu}(k_1, k_2)$, which is the square of the $\ell_1 \rightarrow \ell_2 V$ splitting matrix element (the matrix element of the leptonic current), and a hadronic tensor $\tilde{W}^A_{\mu\nu}(p_A, q)$, which is proportional to the square of the $AV \rightarrow X$ matrix element (the matrix element of the hadronic current). $d\sigma$ can be written as

$$d\sigma \propto \tilde{W}^A_{\mu\nu} L^{\mu\nu}$$
 , (2.5)

and is given more completely by Eq. (A.6) in Appendix A.1. The tilde (~) notation will be made clear in Sec. 4 when we perform a rescaling of nuclear momentum to average nucleon momentum.

The leptonic tensor can be constructed from Feynman diagrams with standard technology. For the case of massless charged leptons exchanging a photon with the nucleus A, the spin-summed leptonic tensor for massless leptons is

$$\sum_{\{\lambda\}} L^{\mu\nu} \Big|_{\text{QED}} = 4e^2 \Big\{ k_1^{\mu} k_2^{\nu} + k_1^{\nu} k_2^{\mu} - (k_1 \cdot k_2) g^{\mu\nu} \Big\} \quad .$$
 (2.6)

Here, $\{\lambda\}$ represents the sum over spins of external particles; spin-averaging is not yet performed. Analogous expressions for the exchange of other bosons can be found in Appendix A.2.

In practice, the tensor $\tilde{W}^A_{\mu\nu}(p_A, q)$, which can only be a function of external momenta p_A and q for unpolarized targets, is decomposed into a sum over tensor-valued coefficient functions multiplied by dimensionless, scalar-valued functions \tilde{W}_i . These \tilde{W}_i are known in the literature as structure functions. Lorentz symmetry and Hermiticity dictate that only certain combinations of p_A and q are allowed in the coefficient functions. For the case of an unpolarized (spin-averaged) nuclear target, the "hadronic" tensor $\tilde{W}^A_{\mu\nu}(p_A, q)$, in terms of the square of the $AV \to X$ current J(z) in coordinate space, is given by

$$\tilde{W}^{A}_{\mu\nu}(p_{A},q) \equiv \frac{1}{4\pi} \oint d^{4}z \ e^{iq \cdot z} \langle A(p_{A}) | J_{\mu}(z) | X(p_{X}) \rangle \langle X(p_{X}) | J_{\nu}(0) | A(p_{A}) \rangle$$
(2.7)

$$= -g_{\mu\nu}\tilde{W}_{1} + \frac{p_{A\mu}p_{A\nu}}{M_{A}^{2}}\tilde{W}_{2} - i\epsilon_{\mu\nu\rho\sigma}\frac{p_{A}q}{M_{A}^{2}}\tilde{W}_{3} + \frac{q_{\mu}q_{\nu}}{M_{A}^{2}}\tilde{W}_{4} + \frac{p_{A\mu}q_{\nu} + p_{A\nu}q_{\mu}}{M_{A}^{2}}\tilde{W}_{5} + \frac{p_{A\mu}q_{\nu} - p_{A\nu}q_{\mu}}{M_{A}^{2}}\tilde{W}_{6} .$$
(2.8)

In Eq. (2.7), the normalization factor $1/(4\pi)$ is conventional and the sum and integral (\slashed{f}) run over all discrete and continuous configurations of X, implying that $\tilde{W}^A_{\mu\nu}(p_A,q)$ is inclusive with respect to X (see also Appendix A.2). We note that, although the targets considered in this work are nuclei, we refer to quantities such as that given in Eq. (2.7) as a hadronic tensors, in keeping with convention. For a polarized target, the decomposition into structure functions takes on a more complicated structure, see e.g. [29, 61, 72–75]. In Sec. 3.2 and Appendix A.5, we review the connection between the \tilde{W}_i and the internal structure of A.

Modern notation calls for using the structure functions \tilde{F}_i rather than \tilde{W}_i . The mapping between the two sets of dimensionless structure functions is given by

$$\{\tilde{F}_1, \ \tilde{F}_2, \ \tilde{F}_3, \ \tilde{F}_4, \ \tilde{F}_{5,6}\}$$

$$= \left\{ \tilde{W}_1, \ \frac{Q^2}{2x_A M_A^2} \tilde{W}_2, \ \frac{Q^2}{x_A M_A^2} \tilde{W}_3, \ \frac{Q^2}{2M_A^2} \tilde{W}_4, \ \frac{Q^2}{2x_A M_A^2} \tilde{W}_{5,6} \right\} .$$

$$(2.9)$$

The purpose of using \tilde{F}_i is to factor out known dependence on Q^2 and make more manifest the phenomena of scaling, i.e., that \tilde{F}_i depend only on x_A , a dimensionless quantity, up to small, logarithmic QCD corrections. In the discussion that follows, we focus on \tilde{F}_1, \tilde{F}_2 , and \tilde{F}_3 . However, we include a detailed discussion of \tilde{F}_4 and \tilde{F}_5 in the Appendix. The structure functions \tilde{F}_4 and \tilde{F}_5 enter into differential cross sections, but are suppressed by a factor $\mathcal{O}(\frac{m_\ell^2}{M_A E_\ell})$, where m_ℓ^2 is the lepton mass squared, M_A is the hadronic mass, and E_ℓ is the energy of one of the external leptons [54]. This suppression is a consequence of contracting the symmetric leptonic tensor $L^{\mu\nu}$ with q_μ (or q_ν), which subsequently vanishes due to the conservation of weak currents by massless leptons. Notably, finite lepton-mass effects could be measured in ν_{τ} -DIS, such as at the SHIP, FASER, or SND@LHC detectors at CERN [31, 76–78]. As for \tilde{F}_6 , which signifies charge-parity violation, the coefficient vanishes when contracted with Eq. (2.6), hence it does not contribute to the cross-section.

Considerations for Spin 1 and Greater

Contrary to a nucleon target with spin-1/2, nuclei can have spin-1 or greater. The case of a spin-1 nuclear targets in NC DIS with charged leptons has been discussed since the 80s[79]. At leading twist, i.e., twist $\tau = 2$, the additional effects of scattering on a polarized spin-1 target reside in a single new structure function $\tilde{b}_1(x)$. This structure function effectively measures the extent to which a target nucleus deviates from a trivial bound state of protons and neutrons. For the deuteron, it is expected that $\tilde{b}_1 \approx 0$, but for other nuclei one could have $\tilde{b}_1 \sim \mathcal{O}(\tilde{F}_1)$.

More generally, using gauge invariance and P- and T-invariance for the spin-1 case, the hadronic tensor can be expressed in terms of eight independent structure functions, $\{\tilde{F}_1, \tilde{F}_2, \tilde{b}_{1,2,3,4}, \tilde{g}_1, \tilde{g}_2\}$. Similar results are found for the hadronic tensor of a (space-like) virtual photon target [80]. The functions \tilde{F}_1 , \tilde{F}_2 , \tilde{g}_1 , and \tilde{g}_2 are analogous to the scaling structure functions of a spin-1/2 target.

 \tilde{F}_1 and \tilde{F}_2 can be measured in the DIS of unpolarized leptons off an unpolarized target whereas the measurement of \tilde{g}_1 and \tilde{g}_2 requires polarized lepton beams and a polarized target. The four structure functions $\tilde{b}_{1,2,3,4}$ are new in the spin-1 case. $\tilde{b}_{1,2}$ are quantities that appear at leading twist, while $\tilde{b}_{3,4}$ appear at higher twist. $\tilde{b}_{1,2}$ are connected via $\tilde{b}_2 = 2x\tilde{b}_1$ in a manner analogous to the Callan-Gross relation, and will receive corrections beyond lowest order in QCD. The structure functions $\tilde{b}_{1,2,3,4}$ can be measured using an unpolarized lepton beam but require a polarized spin-1 target. The TMCs for a (spin-1) deuterium target have been calculated [81] as well as those for a virtual photon target [82].

In this work we are studying unpolarized DIS off an unpolarized nucleus with any spin greater or equal to 1/2. For such circumstances, the hadronic tensor has the same decomposition into structure functions as for a spin-1/2 target. For this reason we will focus only on the \tilde{F}_i structure functions in the remainder of this text; the treatment of additional structure functions, such as \tilde{b}_i , can be dealt with in a similar manner, but we shall not address them explicitly.

3. Nuclear structure functions in the OPE

In this section, we discuss nuclear structure functions and their TMCs in the context of the OPE. We start in Sec. 3.1 with a precise stipulation of the criteria for light-cone dominance involving nuclear targets, whose masses can readily exceed $\mathcal{O}(50 - 100)$ GeV. In Sec. 3.2, we sketch the construction of nuclear structure functions and their TMCs in the OPE; a fuller derivation is given in Appendix A. We then present in Sec. 3.3 the main formula for nuclear TMCs in terms of nuclear quantities x_A and M_A .

3.1. Light-cone dominance of nuclear DIS

A crucial step in deriving structure functions for nucleons and nuclei involves employing the OPE to expand the product of currents that comprise the hadronic tensor of Eq. (2.7). This expansion, however, requires that one is in the limit of light-cone dominance, i.e., $z^2 \sim 0$, where z is the Fourier conjugate of the DIS momentum q. While this is well-established in the case of a nucleon target,³ demonstrating light-cone dominance for nuclear targets requires care due to the relative sizes of $Q^2 = -q^2 > 0$ and M_A^2 .

In many constructions of PDFs and related quantities in DIS, light cone-dominance is described as corresponding to the momentum configuration

$$Q^2 \sim |p_N \cdot q| \gg M_N^2 \,, \tag{3.1}$$

where p_N and M_N are respectively the four-momentum and mass of a single nucleon. Under this condition, naively replacing p_N and M_N by the nuclear momentum p_A and nuclear mass M_A would imply (incorrectly) that light-cone dominance is only satisfied when

$$Q^2 \sim |p_A \cdot q| \gg M_A^2 \,. \tag{3.2}$$

This suggests increasingly large Q^2 are needed for increasing A, and downplays the validity of both pQCD and the parton model. For example: Eq. (3.2) implies that $Q \gg 50$ GeV, which is more than

 $^{^{3}}$ A demonstration of the light cone dominance of DIS can be found, for example, in the textbook by Muta [63, 64], see (2nd edition) pages 228-229 and 262.

half the Z boson mass, is needed to describe DIS data for ⁵⁶Fe, or that $Q \gg 180$ GeV, which is beyond the top quark mass, is needed for ¹⁹⁷Au.

To resolve this, we apply the arguments of Refs. [63, 64] for light-cone dominance in DIS with a nucleon to the case of an arbitrary *nucleus*. Our goal is to identify the dominant contribution to the Fourier integral in Eq. (2.7) in the deeply inelastic limit:

$$Q^2 \to \infty, \quad \nu_A \to \infty, \quad \text{such that} \quad \frac{Q^2}{\nu_A} = 2M_A x_A \quad \text{is fixed.}$$
(3.3)

Note that in terms of averaged nuclear quantities, $Q^2/\nu_A = Q^2/\nu_N = 2M_A x_A = 2M_N x_N$. That is to say, a bigger nuclear mass M_A is compensated by a smaller Bjorken variable x_A such that the fixed Q^2/ν_A is independent of the atomic number.

Following Refs. [63, 64], we examine $\tilde{W}^{A}_{\mu\nu}(p_{A},q)$ in Eq. (2.7), and identify the integration regions that give rise to the dominant contributions. In the DIS limit of Eq. (3.3), as $|q \cdot z| \to \infty$ the exponential of the Fourier integral oscillates without bound and thus makes a vanishing contribution to the integral. Therefore, we need only to consider the region with finite $|q \cdot z|$ in the deeply inelastic regime.

In the target's rest frame, $\nu_A = q_0$. Defining the quantity $\mathbf{z} = \mathbf{q} \cdot \mathbf{z}/|\mathbf{q}|$, we have

$$q \cdot z = q_0 z_0 - \vec{q} \cdot \vec{z} = \nu_A \left(z_0 - \frac{|\vec{q}|}{\nu_A} \, i \right) = \nu_A \left(z_0 - \sqrt{1 + Q^2 / \nu_A^2} \, i \right) \,. \tag{3.4}$$

We note again that this equation is independent of A.

In the DIS limit, $Q^2/\nu_A^2 = (2M_A x_A)/\nu_A$ is small and we can expand the square root:

$$q \cdot z = \nu_A(z_0 - i) - M_A x_A i + \mathcal{O}\left(\frac{M_A^2 i}{\nu_A}\right).$$
(3.5)

Here, the target mass M_A appears but only in combination with x_A and always satisfies $M_A x_A = M_N x_N$. In order to keep $|q \cdot z|$ finite in the deeply inelastic limit, each term on the right hand side of Eq. (3.5) must separately be finite. (Being separately infinite requires the scaling $(z_0 - \imath) \sim (\imath/\nu_A) \rightarrow \infty$, which cannot be consistently satisfied.) Since $M_A x_A$ is fixed, \imath itself must be finite. Therefore, for some constants c > 0 and d > 0, one has

$$|z_0 - \mathbf{z}| < c/\nu_A \quad \text{and} \quad |\mathbf{z}| < d/(x_A M_A). \tag{3.6}$$

The first inequality implies that $|z_0| < |z| + c/\nu_A$. After squaring and using $z^2 = \vec{z}^2 - z_{\perp}^2 < \vec{z}^2$, where z_{\perp} is the component of z orthogonal to \vec{q} , we obtain the inequality chain

$$z_0^2 < (|\mathbf{z}| + c/\nu_A)^2 = \mathbf{z}^2 + \frac{2c|\mathbf{z}|}{\nu_A} + \mathcal{O}\left(\frac{1}{\nu_A^2}\right) < \mathbf{z}^2 + \frac{2cd}{(x_A M_A \nu_A)} + \mathcal{O}\left(\frac{1}{\nu_A^2}\right).$$
(3.7)

To obtain the rightmost bound, we used the second inequality in Eq. (3.6). Using the rightmost equality in Eq (3.3), we obtain the final result:

$$z^{2} = z_{0}^{2} - \vec{z}^{2} < \frac{2cd}{(x_{A}M_{A}\nu_{A})} + \mathcal{O}\left(\frac{1}{\nu_{A}^{2}}\right) = \frac{4cd}{Q^{2}} + \mathcal{O}\left(\frac{1}{\nu_{A}^{2}}\right).$$
(3.8)

We therefore find that the dominant region remains $0 \le z^2 \le (2cd/Q^2)$, and is independent of

the target. This assumes, of course, that no extra A dependence is hidden in the constants c and d. Such an assumption, however, is safe upon inspecting the above derivation. For example: suppose there were separate (minimal) c_A and d_A for each A that satisfied the inequality of Eq. (3.6). Since there are finite many A, we can simply take the largest c_A and d_A , and then define $c = \max\{c_A\}$ and $d = \max\{d_A\}$. This implies that there is a large enough c and d such that Eq. (3.6) holds independent of A.

In light of Eq. (3.8), the requirement of Eq. (3.2) is obviously too harsh. Instead, we advocate that Eq. (3.1) should be understood as requiring $Q^2 \gtrsim$ a few GeV², independent of the hadronic target, in order to justify the application of pQCD and the parton model. More precisely, given a non-perturbative (NP) scale $\Lambda_{\rm NP} \lesssim 1$ GeV, which parameterizes the onset of hadronization, light-cone dominance in DIS for a nucleus A occurs when

$$Q^2 \sim |p_A \cdot q| \gg \Lambda_{\rm NP}^2. \tag{3.9}$$

Under this criterion, Eq. (3.1) is automatically satisfied since nucleon masses scale as $M_N \sim \Lambda_{\rm NP}$. At smaller Q^2 , but still satisfying $Q^2 \gtrsim \Lambda_{\rm NP}^2$, perturbative techniques and the parton model can still be employed as evident by the phenomenon of "precocious scaling" [49, 50, 53]. This denotes the onset of Bjorken scaling at moderate energies despite $\alpha_s(\mu_r = Q) \leq \mathcal{O}(1)$ being sizable, and follows from the fact that light quark masses m_q are comparable to the scale of the QCD Landau pole, $\Lambda_{\rm QCD} \sim \mathcal{O}(0.1)$ GeV.

Such a separation of scales implies the inequality

$$Q^2 \gtrsim \Lambda_{\rm NP}^2 \gg \Lambda_{\rm QCD}^2 \sim m_q^2$$
, (3.10)

where we take m_q to be the light-quark constituent mass at soft quark momenta. This demonstrates that, up to corrections of order $\mathcal{O}(\Lambda_{\rm QCD}^2/Q^2)$, light quarks can be approximated as free and massless partons, even for low Q^2 . However, when such Q^2 is probing a nuclear target A at large x_A , it is paramount to stress that $x_A^2 M_A^2/Q^2$ is not guaranteed to be small or below unity. In this case, TMCs are important.

3.2. Structure functions in the OPE

In this section, we briefly outline the key elements of the OPE required to obtain expressions for structure functions with TMCs at leading twist τ . Formally, TMCs are $\mathcal{O}(M_A^2/Q^2)$ corrections to structure functions that are first generated by the *same* operators that define the structure functions themselves at leading power. For this reason they are sometimes called "kinematical" power corrections [55, 83]. Such power corrections can be isolated from "dynamical" power corrections, which are proportional to the QCD coupling constant, by setting the coupling constant to zero. We generally follow the early literature of Refs. [49, 50, 53, 67, 84, 85], which shed light on the underlying dynamics of hadrons, but note that TMCs can also be derived using diagrammatic methods and finite parton k_T [55, 83]. While we only outline the derivation here, a fuller derivation of nuclear structure functions and their TMCs at leading power in the OPE is provided in Appendix A.

Constructing TMCs for structure functions in the OPE provides a systematic organization of short and long distance physics that does not manifestly rely on the perturbativity of QCD. (This means that it is possible to obtain all-orders results.) Inspired by the program of Refs. [49, 50], the construction of structure functions follows by first recognizing that the inclusive hadronic tensor describing $\gamma^* A \to X$



Figure 3.1: (Left) A contour (circle) along the radius of convergence of the time-ordered matrix element $\tilde{T}^A_{\mu\nu}(p_A,q)$ in the complex $(1/x_A)$ plane for $\gamma^*A \rightarrow \gamma^*A$ scattering. Branch cuts (sawtooth lines) are along the real axis at $x_A^{-1} < 1$ and $x_A^{-1} > 1$ lines. (Right) A deformation of the contour along the branch cuts with vanishing arcs.

scattering in the deeply inelastic limit,⁴

$$\tilde{W}^{A}_{\mu\nu}(p_{A},q) = \frac{1}{4\pi} \int d^{4}z \; e^{iq \cdot z} \langle A | J^{\dagger}_{XA\mu}(z) \; J_{XA\nu}(0) | A \rangle \;, \tag{3.11}$$

which is equivalent to the expression in Eq. (2.7) of Sec. 2.2, and the time-ordered amplitude for virtual Compton scattering process $\gamma^* A \rightarrow \gamma^* A$ in the short-distance limit

$$\tilde{T}^{A}_{\mu\nu}(p_{A},q) \equiv \int d^{4}z \ e^{iq\cdot z} \left\langle A | \mathcal{T} J^{\dagger}_{A\mu}(z) \ J_{A\nu}(0) | A \right\rangle,$$

$$= -g_{\mu\nu} \Delta \tilde{T}^{A}_{1} + \frac{p_{A\mu}p_{A\nu}}{M^{2}_{A}} \Delta \tilde{T}^{A}_{2} - i\epsilon_{\mu\nu\alpha\beta} \frac{p^{\alpha}_{A}q^{\beta}}{M^{2}_{A}} \Delta \tilde{T}^{A}_{3}$$
(3.12)

$$+\frac{q_{\mu}q_{\nu}}{M_{A}^{2}}\Delta\tilde{T}_{4}^{A} + \frac{\left(p_{A\mu}q_{\nu} \pm p_{A\nu}q_{\mu}\right)}{M_{A}^{2}}\Delta\tilde{T}_{5,6}^{A} , \qquad (3.13)$$

are related by the following dispersion relationship [40, 85]

$$\tilde{T}^{A}_{\mu\nu}(p_A,q)\Big|_{(1/x_A)-i\varepsilon}^{(1/x_A)+i\varepsilon} = 4\pi \ \tilde{W}^{A}_{\mu\nu}(p_A,q), \quad \text{for} \quad x_A > 0 \ , \tag{3.14a}$$

$$\tilde{T}^{A}_{\mu\nu}(p_A,q)\Big|_{(1/x_A)+i\varepsilon}^{(1/x_A)-i\varepsilon} = 4\pi \left[\tilde{W}^{A}_{\mu\nu}(p_A,-q)\right]^{\dagger}, \quad \text{for} \quad x_A < 0 \ . \tag{3.14b}$$

In Eq. (3.13), $\tilde{T}^{A}_{\mu\nu}(p_A,q)$ is decomposed into coefficients of Lorentz structures in the same manner as $\tilde{W}^{A}_{\mu\nu}(p_A,q)$ in Eq. (2.8). As $\tilde{T}^{A}_{\mu\nu}(p_A,q)$ is time ordered, the first (second) relation in Eq. (3.14) is for DIS with (anti)particles. We assume here and below that the hadronic currents J^{μ}_{A} and J^{ν}_{A} are always renormalized objects in QCD.

The distinction between the short-distance limit and the DIS limit is important: In the DIS limit, $(Q^2/M_A^2) \rightarrow \infty$ while $x_A = (Q^2/2p_A \cdot q)$ is fixed. In the short-distance limit, $(Q^2/M_A^2) \rightarrow \infty$ while x_A/Q is fixed, meaning that x_A grows with (Q/M_A) ; equivalently, the quantity $(1/x_A)$ is small with

⁴The subscripts XA on the vector currents J_{μ}, J_{ν} are simply labels to denote the hadronic current that takes A to X. The current is independent of states A, X but depends on whether the current is electromagnetic or weak.

increasing (Q/M_A) in the short-distance limit. Formally, $\tilde{T}^A_{\mu\nu}(p_A,q)$ and $\tilde{W}^A_{\mu\nu}(p_A,q)$ are defined in two different limits but can be related through analytic continuation. When analytically continued to complex values of x_A , $\tilde{T}^A_{\mu\nu}$ has desirable analytic properties for $|(1/x_A)| < 1$ (the short-distance limit), whereas $\tilde{W}^A_{\mu\nu}$ is meaningful for $|(1/x_A)| > 1$ (the DIS limit). Hence the discontinuities (branch cuts) of $\tilde{T}^A_{\mu\nu}$ correspond to the kinematics of DIS. Therefore, one can take $\tilde{T}^A_{\mu\nu}(p_A,q)$, decompose it into a contour integral over x_A^{-1} using Cauchy's integral formula, and deform the contour around the discontinuities at $x_A^{-1} < 1$ and $x_A^{-1} > 1$ as shown in Fig. 3.1. (The sawtooth lines along $x_A^{-1} > 1$ are the branch cuts of $\tilde{T}^A_{\mu\nu}$.) Taking the difference of $T^A_{\mu\nu}$ when evaluated just below and above the branch cut, i.e., at the points $x_A^{-1} \pm i\varepsilon$ for $x_A^{-1} > 1$, gives $\tilde{W}^A_{\mu\nu}(p_A,q)$. (For a fuller derivation, see the steps before Eq. (A.45) in Appendix A.3.)

An important consequence of Eq. (3.14) is establishing a link between individual $\Delta \tilde{T}_i^A$ in Eq. (3.13) and the Mellin moments of the structure functions \tilde{F}_i^A defined in Eq. (2.9). Following the steps below Eq. (A.49) in Appendix A.3, one can write for the case⁵ of \tilde{F}_2 [85]

$$\left(\frac{Q^2}{2x_A M_A^2}\right) \Delta \tilde{T}_2^A(Q^2, x_A) = -4i \sum_{N=0}^{\infty} \tilde{F}_2^{AN}(Q^2) x_A^{-N} .$$
(3.15a)

Here, $\tilde{F}_i^{AN}(Q^2)$ is the N^{th} Mellin moment of the structure function $\tilde{F}_i^A(Q^2, x_A)$. The normalizations of the Mellin transformation and its inverse are set by the definitions

$$F^{N} = \int_{0}^{1} dz \ z^{N-1} \ F(z) \quad \text{with} \quad F(z) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} dN \ z^{-N} \ F^{N} \ , \tag{3.16}$$

where c has to be to the right of any poles of F^N in the complex N-plane. As a brief comment, the Mellin transform is an integral over the positive reals, while the inverse transform is an integral over the complex plane. We therefore assume that structure functions can be analytically continued. (In practice, Mellin and inverse-Mellin transforms are implemented in the many computer codes that perform the DGLAP evolution equations in Mellin space, see e.g. Ref. [86].)

The next step in deriving TMCs in the OPE formalism is to take the operator-product expansion of $\tilde{T}^A_{\mu\nu}$ itself in the short-distance limit. After considerable reorganization (see Appendix A.3), one obtains expressions (see Appendix A.4) for $\Delta \tilde{T}^A_i(Q^2, x_A)$ at leading power in the OPE. More specifically, one constructs a power series with respect to powers of x^{-1}_A , and therefore obtains expressions for $\tilde{F}^{AN}_i(Q^2)$, at leading power.

In more detail: the OPE makes use of basic symmetries, e.g., Lorentz invariance and unitarity, to expand hadronic matrix elements into a complete set of local operators \mathcal{O} [87]. As described in Sec. 3.1, this hinges on field operators being along the light cone, i.e., when $z^2 \rightarrow 0$. For forward Compton scattering, the OPE of $\tilde{T}^A_{\mu\nu}$ is given by [40]:

$$\lim_{z^2 \to 0} T^A_{\mu\nu}(p_A, q) \stackrel{\text{OPE}}{=} -2i \sum_{\iota, n} c^{\tau=2,\iota}_{\mu\nu\mu_1\dots\mu_n}(q) \langle A | \mathcal{O}^{\mu_1\dots\mu_n}_{\iota,\tau=2} | A \rangle$$

+ power corrections at $\mathcal{O}(\tau > 2)$ (3.17)

⁵See Appendix A.3 for the treatment of all six \tilde{F}_i^A .

$$= -2i \sum_{k=k_{\min}}^{\infty} \left[-g^{\mu\nu} q_{\mu_1} q_{\mu_2} C_1^{2k} + g^{\mu}_{\mu_1} g^{\nu}_{\mu_2} Q^2 C_2^{2k} - i \epsilon^{\mu\nu\alpha\beta} g_{\alpha\mu_1} q_{\beta} q_{\mu_2} C_3^{2k} \right. \\ \left. + \frac{q^{\mu} q^{\nu}}{Q^2} q_{\mu_1} q_{\mu_2} C_4^{2k} + \left(g^{\mu}_{\mu_1} q^{\nu} q_{\mu_2} \pm g^{\nu}_{\mu_1} q^{\mu} q_{\mu_2} \right) C_{5,6}^{2k} \right] \cdot \left\{ \begin{aligned} 1; & k = 1 \\ \prod_{m=3}^{2k} q_{\mu_m}; & k > 1 \end{aligned} \right\} \\ \left. \times \frac{2^{2k}}{Q^{4k}} A_{\tau=2}^{2k} \tilde{\Pi}^{\mu_1 \dots \mu_{2k}} + \mathcal{O}(\tau > 2). \end{aligned}$$
(3.18)

In Eq. (3.18), the summation over n is changed to n = 2k. The starting point for the summation over k depends on the particular Wilson coefficient; specifically, $k_{\min} = 2$ for C_2^{2k} . while $k_{\min} = 1$ for all other C_{ι}^{2k} 's. In Eq. (3.17), the $\mathcal{O}_{\iota,\tau}^{\mu_1...\mu_n}$ are composite local operators, consisting of quark fields, QCD covariant derivatives, and/or gluon field strengths. Operators with n un-contracted Lorentz indices carry a spin of n. Operators and their coefficients are organized according to their twist $\tau = d - n$, where d is the dimensionality of $\mathcal{O}_{\iota,\tau}^{\mu_1...\mu_n}$ in the standard sense of dimension power counting in an effective field theory. In this study, we focus exclusively on the OPE at leading twist, i.e., $\tau = 2$. We neglect power corrections at $\tau > 2$ but attempt to keep track of their presence. The index ι catalogs all operators \mathcal{O} at a fixed spin and twist. All numerical prefactors are sequestered into effective Wilson coefficients $c_{\mu\nu\mu_1...\mu_n}^{\tau,\iota}(q)$, which are only functions of the external momentum q.

The Wilson coefficients⁶ C_{ι}^{2k} , which are the same for all possible nuclei, can be identified and matched to quantities calculated in perturbative QCD. The quantity $A_{\tau=2}^{2k}$ is the scalar-valued coefficient of the expectation value $\langle A | \mathcal{O}_{\iota,\tau=2}^{\mu_1...\mu_{2k}} | A \rangle = A_{\tau=2}^{2k} \times \tilde{\Pi}^{\mu_1...\mu_{2k}}$. $A_{\tau=2}^{2k}$ is sometimes labeled the reduced matrix element of $\langle A | \mathcal{O}_{\iota,\tau=2}^{\mu_1...\mu_{2k}} | A \rangle$. Importantly, the Lorentz structure of this expectation value $(\tilde{\Pi}^{\mu_1...\mu_{2k}})$ can only be a function of p_A as the q dependence is sequestered elsewhere. Consequentially, $\tilde{\Pi}^{\mu_1...\mu_{2k}}$ can be organized according to permutations and powers of momenta $p_A^{\mu_m}$ and metrics $g^{\mu_m\mu_n}$, and scales as [49, 50]

$$\tilde{\Pi}^{\mu_1...\mu_{2k}} = \sum_{j=0}^k (-1)^j \; \frac{(2k-j)!}{2^j (2k)!} \; \underbrace{\{g...g\}}_{j \; g^{\mu_n \mu_m'_s}} \; \underbrace{\{p_A...p_A\}}_{(2k-2j) \; p_A^{\mu_n'_s}} \; (p_A^2)^j \; + \text{ power corrections} \tag{3.19}$$

$$\sim \underbrace{\{p_A...p_A\}}_{2k \ p_A^{\mu_m}} + \underbrace{\{p_A...p_A\}}_{(2k-2) \ p_A^{\mu_m}} \underbrace{\{g...g\}}_{1 \ g^{\mu_m\mu_n}} (p_A^2)^1 + \underbrace{\{p_A...p_A\}}_{(2k-4) \ p_A^{\mu_m}} \underbrace{\{g...g\}}_{2 \ g^{\mu_m\mu_n}} (p_A^2)^2 + \dots$$
(3.20)

Focusing on Eq. (3.20), one sees that the momentum factor $\Pi^{\mu_1...\mu_{2k}}$ scales as a tower of momentum and mass terms. For a fixed index k, the series ranges from the product of 2k instances of the nucleus' momentum $p_A^{\mu_m}$ and no instance of the nucleus' mass $p_A^2 = M_A^2$ (as shown in the leftmost term), to no instances of $p_A^{\mu_m}$ but one instance of $(p_A^2)^k = (M_A^2)^k$. The brackets $\{...\}$ represent the collection of all possible permutations of Lorentz indices. The dimension of each term is [mass]^{2k}. Again, we neglect higher-order power corrections.

Whether one retains $(M_A^2)^j$ terms or neglects them outright is the distinction between obtaining structure functions with TMCs or obtaining structure functions without TMCs (which we shall later label as "No-TMC"). For example: after contracting all Lorentz indices in Eq. (3.18), one can neglect all factors of (M_A^2/Q^2) . This is equivalent to truncating the summation in Eq. (3.19) at j = 0. Doing

⁶In principle, the Wilson coefficients C_{ι}^{2k} also carry a species index f, which runs over the gluon and quark flavors. In this case, C_{ι}^{2k} should be replaced by $\sum_{f=u,g,\ldots} C_{if}^{2k}$, but this dependence is neglected here for simplicity.

so and comparing the result to Eq. (3.15) recovers (see Appendix A.5 for details):

$$\tilde{F}_{i}^{AN} = \int_{0}^{1} dy \ y^{N-1} \ \tilde{F}_{i}^{A}(y,Q^{2}) = C_{i}^{N}(Q^{2})A_{\tau=2}^{N} + \text{ power corrections, for } i = 1,3,4,5, \quad (3.21a)$$

$$\tilde{F}_{2}^{A(N-1)} = \int_{0}^{1} dy \ y^{N-2} \ \tilde{F}_{2}^{A}(y,Q^{2}) = C_{2}^{N}(Q^{2})A_{\tau=2}^{N} + \text{ power corrections }.$$
(3.21b)

Non-trivially, Eq. (3.21) states that one can identify the product of the Wilson coefficient and reduced hadronic matrix element, $(C_{\iota}^{2k} A_{\tau=2}^{2k})$, as integer Mellin moments of structure functions, up to power corrections. (We note that footnote 6 is still applicable.)

If one does not truncate the momentum factor $\Pi^{\mu_1\dots\mu_{2k}}$ in Eq. (3.19) at j = 0 and instead retains the sums over all j, and hence retains the sum over powers of $p_A^2 = M_A^2$, then one obtains expressions that are analogous to Eq. (3.21). For example: one can identify the Mellin transformation of the structure function F_2 with a nonzero target mass as the product $(C_{\iota=2}^{2k}A_{\tau=2}^{2k})$ with a coefficient proportional to $(M_A^2/Q^2)^j$:

$$\int_{0}^{1} dx_A \, x_A^{N-2} \tilde{F}_2^{A, \text{TMC}}(x_A, Q^2) = \sum_{j=0}^{\infty} \left(\frac{M_A^2}{Q^2}\right)^j \frac{(N+j)!}{j! \ (N-2)!} \ \frac{C_2^{N+2j} A_{\tau=2}^{N+2j}}{(N+2j)(N+2j-1)} \ . \tag{3.22}$$

Similar expressions can be found in Appendix A.5 for the other structure functions. Intuitively, the right hand side of Eq. (3.22) states that structure functions with TMCs can be thought simply as the product in moment space of a structure function for a massless target, i.e., the $(C_i A_{\tau=2})$ factor, and a kinematical factor, i.e., everything else. For this reason, the TMCs under discussion are sometimes called "purely kinematical" [55, 83, 88]. It is not obvious but, with the use of generating functions, the inverse-Mellin transform of Eq. (3.22) has a closed form [49, 50, 81]. That is to say, one obtains expressions for structure functions with TMCs in terms of structure functions for a massless target in x-space. These results are summarized in the following section for F_1^A, F_2^A , and F_3^A , and in Appendix A.5 for all F_i .

3.3. Master formula for structure functions with TMCs in ℓA DIS

Following the procedure outlined in Sec. 3.2, we obtain a set of master formulas for twist-2 target mass corrections to structure functions for nuclei that are similar to those in Eq. (23) in Ref. [60] for nucleons. Using the notation of Refs. [54, 60], the general master formula for target mass-corrected structure functions $F_i^{A,\text{TMC}}$ for $j = 1, \ldots, 6$ reads:

$$\tilde{F}_{j}^{A,\text{TMC}}(x_{A},Q^{2}) = \sum_{i=1}^{6} A_{j}^{i} \tilde{F}_{i}^{A,(0)}(\xi_{A},Q^{2}) + B_{j}^{i} \tilde{h}_{i}^{A}(\xi_{A},Q^{2}) + C_{j} \tilde{g}_{2}^{A}(\xi_{A},Q^{2}).$$
(3.23)

On the left-hand side, the $F_j^{A,\text{TMC}}$ take as arguments the Bjorken scaling variable x_A and scale Q; on the right-hand side, individual terms are given in terms of structure functions for massless nuclei $\tilde{F}_i^{A,(0)}$, which take as arguments the Nachtmann variable ξ_A and scale Q. The coefficients A_j^i , B_j^i , C_j are derived in Appendix A, and are the same that are given in Tables I, II, III in [54] for $j = 1, \ldots, 5$. Suppressing the Q^2 dependence for brevity, one finds at twist τ = 2 the following:

$$\tilde{F}_{1}^{A,\text{TMC}}(x_{A}) = \left(\frac{x_{A}}{\xi_{A}r_{A}}\right)\tilde{F}_{1}^{A,(0)}(\xi_{A}) + \left(\frac{M_{A}^{2}x_{A}^{2}}{Q^{2}r_{A}^{2}}\right)\tilde{h}_{2}^{A}(\xi_{A}) + \left(\frac{2M_{A}^{4}x_{A}^{3}}{Q^{4}r_{A}^{3}}\right)\tilde{g}_{2}^{A}(\xi_{A}), \qquad (3.24a)$$

$$\tilde{F}_{2}^{A,\text{TMC}}(x_{A}) = \left(\frac{x_{A}^{2}}{\xi_{A}^{2}r_{A}^{3}}\right)\tilde{F}_{2}^{A,(0)}(\xi_{A}) + \left(\frac{6M_{A}^{2}x_{A}^{3}}{Q^{2}r_{A}^{4}}\right)\tilde{h}_{2}^{A}(\xi_{A}) + \left(\frac{12M_{A}^{4}x_{A}^{4}}{Q^{4}r_{A}^{5}}\right)\tilde{g}_{2}^{A}(\xi_{A}), \qquad (3.24b)$$

$$\tilde{F}_{3}^{A,\text{TMC}}(x_{A}) = \left(\frac{x_{A}}{\xi_{A}r_{A}^{2}}\right)\tilde{F}_{3}^{A,(0)}(\xi_{A}) + \left(\frac{2M_{A}^{2}x_{A}^{2}}{Q^{2}r_{A}^{3}}\right)\tilde{h}_{3}^{A}(\xi_{A}), \qquad (3.24c)$$

$$\tilde{F}_{4}^{A,\text{TMC}}(x_{A}) = \left(\frac{x_{A}}{\xi_{A}r}\right) \tilde{F}_{4}^{A,(0)}(\xi_{A}) \Big|_{\text{No TMCs}} - \left(\frac{2M_{A}^{2}x_{A}^{2}}{Q^{2}r^{2}}\right) \tilde{F}_{5}^{A,(0)}(\xi_{A}) + \left(\frac{M_{A}^{4}x_{A}^{3}}{Q^{4}r^{3}}\right) \tilde{F}_{2}^{A,(0)}(\xi_{A})
+ \left(\frac{M_{A}^{2}x_{A}^{2}}{Q^{2}r^{3}}\right) \tilde{h}_{5}^{A}(\xi_{A}) - \left(\frac{2M_{A}^{4}x_{A}^{4}}{Q^{4}r^{4}}\right) \left(2 - \xi^{2}M_{A}^{2}/Q^{2}\right) \tilde{h}_{2}^{A}(\xi_{A})
+ \left(\frac{2M_{A}^{4}x_{A}^{3}}{Q^{4}r^{5}}\right) \left(1 - 2x_{A}^{2}M_{A}^{2}/Q^{2}\right) \tilde{g}_{2}^{A}(\xi_{A}) ,$$
(3.24d)

$$\tilde{F}_{5}^{A,\text{TMC}}(x_{A}) = \left(\frac{x_{A}}{\xi_{A}r^{2}}\right) \tilde{F}_{5}^{A,(0)}(\xi_{A}) - \left(\frac{M_{A}^{2}x_{A}^{2}}{Q^{2}r^{3}\xi_{A}}\right) \tilde{F}_{2}^{A,(0)}(\xi_{A}) + \left(\frac{M_{A}^{2}x_{A}^{2}}{Q^{2}r^{3}}\right) \tilde{h}_{5}^{A}(\xi_{A}) - \left(\frac{2M_{A}^{2}x_{A}^{2}}{Q^{2}r^{4}}\right) \left(1 - x\xi M_{A}^{2}/Q^{2}\right) \tilde{h}_{2}^{A}(\xi_{A}) + \left(\frac{6M_{A}^{4}x_{A}^{3}}{Q^{4}r^{5}}\right) \tilde{g}_{2}^{A}(\xi_{A}) , \qquad (3.24e)$$

$$\tilde{F}_{6}^{A,\text{TMC}}(x_{A}) = \left(\frac{x_{A}}{\xi_{A}r_{A}^{2}}\right) \tilde{F}_{6}^{A,(0)}(\xi_{A}) + \left(\frac{2M_{A}^{2}x_{A}^{2}}{Q^{2}r^{3}}\right) \tilde{h}_{6}(\xi_{A}) .$$
(3.24f)

Here, the functions $\tilde{h}_i^A(\xi_A, Q^2)$ and $\tilde{g}_2^A(\xi_A, Q^2)$ are given by the integrals

$$\tilde{h}_{2}^{A}(\xi_{A},Q^{2}) = \int_{\xi_{A}}^{1} du_{A} \; \frac{\tilde{F}_{2}^{A,(0)}(u_{A},Q^{2})}{u_{A}^{2}}, \tag{3.25a}$$

$$\tilde{h}_{3}^{A}(\xi_{A},Q^{2}) = \int_{\xi_{A}}^{1} du_{A} \; \frac{\tilde{F}_{3}^{A,(0)}(u_{A},Q^{2})}{u_{A}}, \tag{3.25b}$$

$$\tilde{h}_{5}^{A}(\xi_{A},Q^{2}) = \int_{\xi_{A}}^{1} du_{A} \ \frac{2\tilde{F}_{5}^{A,(0)}(u_{A},Q^{2})}{u_{A}}, \tag{3.25c}$$

$$\tilde{h}_{6}^{A}(\xi_{A},Q^{2}) = \int_{\xi_{A}}^{1} du_{A} \; \frac{\tilde{F}_{6}^{A,(0)}(u_{A},Q^{2})}{u_{A}}, \tag{3.25d}$$

$$\tilde{g}_2^A(\xi_A, Q^2) = \int_{\xi_A}^1 du_A \ \tilde{h}_2^A(u_A, Q^2).$$
(3.25e)

The equation Eq. (3.23) does not assume or imply any Callan-Gross relation between F_1 and F_2 .

Therefore, one can compute the longitudinal structure function and obtain:

$$\tilde{F}_{L}^{A,\text{TMC}}(x_{A}) = r_{A}^{2}\tilde{F}_{2}^{A,\text{TMC}}(x_{A}) - 2x_{A}\tilde{F}_{1}^{A,\text{TMC}}(x_{A})
= \frac{x_{A}^{2}}{\xi_{A}^{2}r_{A}} [\tilde{F}_{2}^{A,(0)}(\xi_{A}) - 2\xi_{A}\tilde{F}_{1}^{A,(0)}(\xi_{A})] + \frac{4M_{A}^{2}x_{A}^{3}}{Q^{2}r_{A}^{2}}\tilde{h}_{2}^{A}(\xi_{A}) + \frac{8M_{A}^{4}x_{A}^{4}}{Q^{4}r_{A}^{3}}\tilde{g}_{2}^{A}(\xi_{A})
= \frac{x_{A}^{2}}{\xi_{A}^{2}r_{A}}\tilde{F}_{L}^{A,(0)}(\xi_{A}) + \frac{4M_{A}^{2}x_{A}^{3}}{Q^{2}r_{A}^{2}}\tilde{h}_{2}^{A}(\xi_{A}) + \frac{8M_{A}^{4}x_{A}^{4}}{Q^{4}r_{A}^{3}}\tilde{g}_{2}^{A}(\xi_{A}) .$$
(3.26)

This general result is non-zero and thus violates the Callan-Gross relation, as it should. The leading term $\tilde{F}_L^{A,(0)}$ will be non-zero for finite quark masses, and the sub-leading terms $\tilde{h}_i^A, \tilde{g}_2^A$ contribute for finite hadron mass M_A .

Note that effects from heavy quark masses are taken into account in the $\tilde{F}_i^{A,(0)}$ as outlined in Sec. 4 and Ref. [60]. This is not done by generalizing the Nachtmann variable ξ_A directly in Eq. (3.23).

Remarks

We note that the formula for TMCs in Eq. (3.23) was derived without invoking perturbative QCD. In principle, one can start from matrix elements that describe collinear parton splitting at some fixed order in QCD, construct structure functions, compare to $\tilde{T}_{\mu\nu}^{A}$ in Eq. (3.18), and obtain expressions for C_{ι}^{2k} to some power in the strong coupling constant α_s . However, by constructing structure functions in the OPE, the C_{ι}^{2k} in Eq. (3.18) are defined at any order α_s . Ultimately, this implies that all the α_s dependence of $\tilde{F}_i^{A,\text{TMC}}$ is contained in $\tilde{F}_i^{A,(0)}$. This also means that the coefficients A_j^i , B_j^i and C_j , and the variable ξ_A are the same irrespective to the order (LO, NLO, NNLO, ...) at which the structure functions $\tilde{F}_i^{A,(0)}$ are considered.

In the above expressions for F_i^{TMC} , we emphasize that it is (x_A, Q^2) and not (ξ_A, Q^2) that is the correct point in phase space to evaluate structure functions. While on the surface it may appear strange to have the left-hand side of Eq. (3.23) to be a function of x_A while the right-hand side be a function of ξ_A , this difference arises naturally in the calculation. In particular, final-state kinematics in ℓA DIS are constrained by momentum conservation, with $\delta^4(q + P_A - P_X) \sim \delta(x_A - \xi_A)$. Thus, we can write schematically

$$F_i^{\text{TMC}}(x_A, Q^2) \sim F_i^{(0)}(x_A, Q^2) \,\delta(x_A - \xi_A) \sim F_i^{(0)}(\xi_A, Q^2). \tag{3.27}$$

Note that it would be incorrect to write $F_i^{\text{TMC}}(\xi_A, Q^2) \sim F_i^{(0)}(\xi_A, Q^2)$, as the Mellin and inverse-Mellin transformations in Sec. 3.2 are defined with respect to x_A , not ξ_A .

Another feature of Eq. (3.23) is that h_2 and g_2 appear in the formulas for both F_1^{TMC} and F_2^{TMC} . This follows directly from the forms of $\tilde{T}_{\mu\nu}^A$ and the momentum factor $\tilde{\Pi}$ in Eq. (3.18), in the massive target limit. For example: the terms proportional to C_1^{2k} and to C_2^{2k} both contribute to $\Delta \tilde{T}^{A,1}$, which multiplies the Lorentz structure $-g_{\mu\nu}$ in Eq. (3.18). As shown in Eq. (3.20), $\tilde{\Pi}$ in the massive-target limit is proportional to terms that (a) consist exclusively of $p_A^{\mu_n}$ factors as well as to terms that (b) include factors of momentum $p_A^{\mu_n}$, metric $g^{\mu_m\mu_n}$, and mass M_A . Among the many possible contractions in Eq. (3.18) are the "(a)" terms contracting with the C_1^{2k} term and the "(b)" terms contracting with the C_2^{2k} term. Both sets of contractions generate a term proportional to $-g_{\mu\nu}$. However, the "(b)" set of contractions only enters $\Delta \tilde{T}^{A,1}$ as a TMC, i.e., it vanishes in the $(M_A^2/Q^2) \rightarrow 0$ limit. Arguably, one can think of C_1^{2k} feeding into F_1^{TMC} as a helicity-conserving contribution, which survives in the massless limit, whereas the contribution from C_2^{2k} is a helicity-inverting contribution, which vanishes in the massless limit.

Nucleus A	Nucleon N				
$M_A = A M_N$	$M_N = M_A/A$				
$p_A = A p_N$	$p_N = p_A/A$				
$x_A = \frac{Q^2}{2p_A \cdot q} \equiv x_N / A$	$x_N = \frac{Q^2}{2p_N \cdot q} \equiv A x_A$				
$x_A \in [0,1]$	$x_N \in [0, A]$				
$W_A^2 = (p_A + q)^2$	$W_N^2 = (p_N + q)^2$				
$\nu_A = (q \cdot p_A)/M_A \equiv \nu_N$	$\nu_N = (q \cdot p_N)/M_N \equiv \nu_A$				
$y_A = \nu_A / E \equiv y_N$	$y_N = \nu_N / E \equiv y_A$				
Nachtmann Variable & Hadronic Mass					
$r_A = \sqrt{1 + \frac{4x_A^2 M_A^2}{Q^2}} \equiv r_N$	$r_N = \sqrt{1 + \frac{4x_N^2 M_N^2}{Q^2}} \equiv r_A$				
$\xi_A = R_M x_A \equiv \xi_N / A$	$\xi_N = R_M x_N \equiv A \xi_A$				
$\xi_A \in [0,1]$	$\xi_N \in [0, A]$				
Since $r_A = r_N \equiv r$, then $R_M = \frac{2}{1 + r_{A,N}}$					
Also, $\xi_A / x_A = \xi_N / x_N = R_M = \frac{2}{1+r}$					
We define $\varepsilon = (xM/Q)$.					

Summary table of key relations:

Table 4.1: We summarize the key relations for a nucleus (A) and nucleon (N). We also find it convenient to define $\varepsilon = (xM/Q)$; we omit the subscripts on ε for brevity as $(x_AM_A/Q) = (x_NM_N/Q)$. We caution that W_A and W_N are **not** simply related, *c.f.*, Sec. 5.4. Note, the target mass modifies the scaling variable via $\xi_A = R_M x_A$. Additionally, we introduce the shorthand notation $r = r_A = r_N$.

4. Rescaling

In this section we introduce a rescaling between the nuclear and averaged nucleon kinematics. Many of the key relations are summarized in Table 4.1. We typically identify the nuclear variables with an "A" subscript (x_A, ξ_A, r_A) , and those of the nucleon with an "N" subscript (x_N, ξ_N, r_N) . Similarly, the original nuclear structure functions are identified with a tilde (\tilde{W}, \tilde{F}) , while those re-scaled to the kinematics of an averaged nucleon are without (W, F).

4.1. Nuclear and Nucleon Kinematics

As can be seen from Eqs. (3.24) and (3.26), the M_A^{2j} terms are always accompanied by x_A^{2j} factors in *x*-space. Furthermore, the structure functions $\tilde{F}_2^{A,(0)}(x_A)$ have their main support at $x_A = x_N/A \sim 1/A$. Therefore, we have effectively

$$\frac{M_A^{2j} x_A^{2j}}{Q^{2j}} = \frac{\left(M_N^{2j} A^{2j}\right) x_A^{2j}}{Q^{2j}} = \frac{M_N^{2j} x_N^{2j}}{Q^{2j}} \sim \frac{M_N^{2j}}{Q^{2j}}, \qquad (4.1)$$

where $M_N = M_A/A$ is the (average) nucleon mass such that the terms with j > 0 in Eq. (3.20) are suppressed for $Q^2 > M_N^2$ independent of the nuclear target.

We introduce the *average nucleon momentum* $p_N \coloneqq p_A/A$ inside the nucleus and define a nucleon scaling variable

$$x_N = \frac{Q^2}{2p_N \cdot q} = A x_A , \quad \text{where} \quad x_N \in [0, A] , \qquad (4.2)$$

in contrast to the original Bjorken variable

$$x_A = \frac{Q^2}{2p_A \cdot q} = x_N / A$$
, where $x_A \in [0, 1]$. (4.3)

Note that the original x_A variable can be constructed directly from the external momenta of the particles whereas the "averaged quantities" x_N (and p_N) are not directly observable for nuclear targets with A > 1.

Using $M_A = AM_N$, we find for the Nachtmann variable

$$\xi_A = \frac{2x_A}{1 + \sqrt{1 + 4x_A^2 M_A^2/Q^2}} = \frac{1}{A}\xi_N \tag{4.4}$$

with

$$\xi_N = \frac{2x_N}{1 + \sqrt{1 + 4x_N^2 M_N^2/Q^2}} \,. \tag{4.5}$$

Conversely, the quantity R_M , which relates the Bjorken x variable to the scaling variable ξ via $\xi = R_M x$, is the same for both the nucleus and nucleon case:

$$R_M = \frac{2}{1+r_A} = \frac{2}{1+r_N} \ . \tag{4.6}$$

This is because the quantity r_A appearing in the master formula is also the same:

$$r_A = \sqrt{1 + 4x_A^2 M_A^2/Q^2} = \sqrt{1 + 4x_N^2 M_N^2/Q^2} = r_N \,. \tag{4.7}$$

Hence, we have $\xi_A = R_M x_A$ and $\xi_N = R_M x_N$, where R_M takes the same value in both equations.

⁷Note that we have introduced the nucleon scaling variable $x_N = Ax_A$; this will be discussed in detail throughout this section. Although x_N can in principle extend to $x_N = A$, the dominant range of the kinematics is $0 \le x_N \le 1$. Discussion on the $x_N > 1$ region are given in Sec. 5.5.

4.2. Rescaled Structure Functions

Turning to the structure functions, we define a rescaled hadronic tensor via the relation

$$AW^{A}_{\mu\nu}(p_{N},q) \coloneqq \tilde{W}^{A}_{\mu\nu}(p_{A},q) , \qquad (4.8)$$

expressed in terms of rescaled structure functions:

$$F_2^A(x_N, Q^2) \coloneqq \tilde{F}_2^A(x_A, Q^2) , \qquad x_N F_{1,3}^A(x_N, Q^2) \coloneqq x_A \tilde{F}_{1,3}^A(x_A, Q^2) .$$
(4.9)

The general relations in Eq. (4.9) hold also for $\tilde{F}_{4,5,6}$, both with and without TMCs, and for any value of the twist. As we show below, they are consistent with the master formula for the target-mass-corrected structure functions. Hence, the pattern of the master equations will be consistent between the nucleus and nucleon.

Indeed, one easily finds from Eq. (3.25):

$$\tilde{h}_{2}^{A}(\xi_{A}) = \int_{\xi_{A}}^{1} du_{A} \frac{\tilde{F}_{2}^{A(0)}(u_{A})}{u_{A}^{2}} = A \int_{\xi_{N}}^{A} du_{N} \frac{F_{2}^{A(0)}(u_{N})}{u_{N}^{2}} =: Ah_{2}^{A}(\xi_{N}), \qquad (4.10)$$

$$\tilde{h}_{3}^{A}(\xi_{A}) = \int_{\xi_{A}}^{1} du_{A} \frac{\tilde{F}_{3}^{A(0)}(u_{A})}{u_{A}} = A \int_{\xi_{N}}^{A} du_{N} \frac{F_{3}^{A(0)}(u_{N})}{u_{N}} =: Ah_{3}^{A}(\xi_{N}), \qquad (4.11)$$

$$\tilde{h}_{5}^{A}(\xi_{A}) = \int_{\xi_{A}}^{1} du_{A} \frac{2\tilde{F}_{5}^{A(0)}(u_{A})}{u_{A}} = A \int_{\xi_{N}}^{A} du_{N} \frac{2F_{5}^{A(0)}(u_{N})}{u_{N}} =: Ah_{5}^{A}(\xi_{N}), \qquad (4.12)$$

$$\tilde{h}_{6}^{A}(\xi_{A}) = \int_{\xi_{A}}^{1} du_{A} \frac{\tilde{F}_{6}^{A(0)}(u_{A})}{u_{A}} = A \int_{\xi_{N}}^{A} du_{N} \frac{F_{6}^{A(0)}(u_{N})}{u_{N}} =: Ah_{6}^{A}(\xi_{N}), \qquad (4.13)$$

$$\tilde{g}_{2}^{A}(\xi_{A}) = \int_{\xi_{A}}^{1} du_{A} \tilde{h}_{2}^{A}(u_{A}) = \frac{1}{A} \int_{\xi_{N}}^{A} du_{N} A h_{2}^{A}(u_{N}) =: g_{2}^{A}(\xi_{N}).$$
(4.14)

With these expressions, and using $x_N = Ax_A$, $\xi_N = A\xi_A$, $r_A = r_N$, $M_A = AM_N$, we obtain the master

TMC formula for the rescaled nuclear structure functions:

$$F_1^{A,\text{TMC}}(x_N, Q^2) = \left(\frac{x_N}{\xi_N r_N}\right) F_1^{A,(0)}(\xi_N, Q^2) + \left(\frac{M_N^2 x_N^2}{Q^2 r_N^2}\right) h_2^A(\xi_N, Q^2) + \left(\frac{2M_N^4 x_N^3}{Q^4 r_N^3}\right) g_2^A(\xi_N, Q^2) ,$$
(4.15a)

$$F_2^{A,\text{TMC}}(x_N, Q^2) = \left(\frac{x_N^2}{\xi_N^2 r_N^3}\right) F_2^{A,(0)}(\xi_N, Q^2) + \left(\frac{6M_N^2 x_N^3}{Q^2 r_N^4}\right) h_2^A(\xi_N, Q^2) + \left(\frac{12M_N^4 x_N^4}{Q^4 r_N^5}\right) g_2^A(\xi_N, Q^2) ,$$
(4.15b)

$$F_3^{A,\text{TMC}}(x_N, Q^2) = \left(\frac{x_N}{\xi_N r_N^2}\right) F_3^{A,(0)}(\xi_N, Q^2) + \left(\frac{2M_N^2 x_N^2}{Q^2 r_N^3}\right) h_3^A(\xi_N, Q^2) , \qquad (4.15c)$$

$$F_{4}^{A,\text{TMC}}(x_{A},Q^{2}) = \left(\frac{x_{A}}{\xi_{A}r}\right) F_{4}^{A,(0)}(\xi_{A},Q^{2}) - \left(\frac{2M_{A}^{2}x_{A}^{2}}{Q^{2}r^{2}}\right) F_{5}^{A,(0)}(\xi_{A},Q^{2}) + \left(\frac{M_{A}^{4}x_{A}^{3}}{Q^{4}r^{3}}\right) F_{2}^{A,(0)}(\xi_{A},Q^{2}) + \left(\frac{M_{A}^{2}x_{A}^{2}}{Q^{2}r^{3}}\right) h_{5}^{A}(\xi_{A},Q^{2}) - \left(\frac{2M_{A}^{4}x_{A}^{4}}{Q^{4}r^{4}}\right) \left(2 - \xi^{2}M_{A}^{2}/Q^{2}\right) h_{2}^{A}(\xi_{A},Q^{2}) + \left(\frac{2M_{A}^{4}x_{A}^{3}}{Q^{4}r^{5}}\right) \left(1 - 2x_{A}^{2}M_{A}^{2}/Q^{2}\right) g_{2}^{A}(\xi_{A},Q^{2}) , \qquad (4.15d)$$

$$F_{5}^{A,\text{TMC}}(x_{A},Q^{2}) = \left(\frac{x_{A}}{\xi_{A}r^{2}}\right) F_{5}^{A,(0)}(\xi_{A},Q^{2}) - \left(\frac{M_{A}^{2}x_{A}^{2}}{Q^{2}r^{3}\xi_{A}}\right) F_{2}^{A,(0)}(\xi_{A},Q^{2}) + \left(\frac{M_{A}^{2}x_{A}^{2}}{Q^{2}r^{3}}\right) h_{5}^{A}(\xi_{A},Q^{2}) - \left(\frac{2M_{A}^{2}x_{A}^{2}}{Q^{2}r^{4}}\right) \left(1 - x\xi M_{A}^{2}/Q^{2}\right) h_{2}^{A}(\xi_{A},Q^{2}) + \left(\frac{6M_{A}^{4}x_{A}^{3}}{Q^{4}r^{5}}\right) g_{2}^{A}(\xi_{A},Q^{2}) , \qquad (4.15e)$$

$$F_6^{A,\text{TMC}}(x_A, Q^2) = \left(\frac{x_A}{\xi_A r_A^2}\right) F_6^{A,(0)}(\xi_A, Q^2) + \left(\frac{2M_A^2 x_A^2}{Q^2 r^3}\right) h_6(\xi_A, Q^2) .$$
(4.15f)

The master equations take the same form as before, but are entirely expressed in terms of averaged nucleon variables. This is one of the main results of this rescaling demonstration.

4.3. Impact of Quark Masses

Notably, we stress that all of the above derivations have made no mention of the parton model. This can be understood from the fact that the quark and gluon sub-structure of nuclei enters only implicitly in intermediate steps of the derivation of the master formula since the hadronic currents appearing in the hadronic tensor are composed of quark and gluon fields and we have relied on the validity of the OPE for the products of these currents. Thus, due to the nature of the factorization, all detail related to quark masses is contained entirely inside the $F_i^{A,(0)}$ structure functions, and the forms of the master formulas in Eq. (4.15) are unchanged.

The treatment of the quark masses was presented in detail in Ref. [60], but it is instructive to briefly mention how the quark masses effect the rescaling of the kinematic variables as summarized in Table 4.2. Just as the hadron mass modifies the scaling variable via the relation $\xi = R_M x$, the quark masses further modify the scaling variable via the relation $\bar{\xi} = R_{ij}\xi = R_{ij}R_M\xi$. The R_M factor depends on the hadron mass, and the R_{ij} factor depends on the incoming and outgoing quark masses $\{m_i, m_j\}$, respectively. Note the hadronic and partonic correction factors factorize, and are applied multiplicatively. Additionally, observe that in the limit the initial-state quark mass vanishes $m_i = 0$, we have the familiar "slow-rescaling" limit $R_{ij} = (1 + m_j^2/Q^2)$.

Partonic Masses				
$\bar{\xi}_A = R_{ij}\xi_A = R_{ij}R_M x_A$		$\bar{\xi}_N = R_{ij}\xi_N = R_{ij}R_M x_N$		
For $q_i + V \rightarrow q_j$				
$R_{ij} = \frac{1}{2Q^2} [Q^2 - m_i^2 + m_j^2 + \Delta(Q^2, m_i^2, m_j^2)]$				
$\Delta(a, b, c) = \sqrt{a^2 + b^2 + c^2 - 2(ab + bc + ca)}$				
For $g + V \rightarrow q_j + q_k$				
$R_{jk} = 1 + \frac{(m_j + m_k)^2}{Q^2}$				

Table 4.2: Summary table of partonic mass relations. Note, the target mass modifies the scaling variable via $\xi_A = R_M x_A$. This is further modified by the parton masses $\overline{\xi}_A = R_{ij}\xi_A = R_{ij}R_M x_A$; hence these multiplicative corrections factorize between the hadron and parton masses [60]. In the limit the initial-state parton mass $m_i \to 0$, we obtain the "slow rescaling" result: $R_{ij} = (1 + m_j^2/Q^2)$.

5. Parton model

We now turn to a discussion of structure functions in the context of the parton model originally proposed by Feynman in 1969 [89] and applied to electron-proton DIS by Bjorken and Paschos [90]. In the years following these works, partons were identified with quarks and gluons, and the heuristic parton model including QCD effects was rigorously derived from first principles of QCD in the context of factorization theorems [41–48]. In this section we summarize the main elements of the conventional "massless" QCD-improved parton model, also known as Zero Mass Variable Flavor Number Scheme (ZM-VFNS), where quark-mass effects in the hard scattering cross sections and target-mass effects are neglected. There is an extensive body of literature on computations with heavy partons; Collins extended the factorization proof [42] to the case of heavy partons in Ref. [45], and implementations can be found in Refs. [58, 91–94]. A discussion of the parton model including such mass effects will be given in Sec. 6.

In the QCD-improved parton model the nuclear structure functions $(\tilde{\mathcal{F}}_k^A)$ are given as convolutions of target-independent, short-distance, Wilson coefficients $(C_{k,i})$ with universal nuclear parton distribution functions (\tilde{f}_i^A) :

$$\tilde{\mathcal{F}}_{k}^{A}(x_{A},Q^{2}) = \int_{x_{A}}^{1} \frac{dy_{A}}{y_{A}} \tilde{f}_{i}^{A}(y_{A},Q^{2}) C_{k,i}(x_{A}/y_{A}) + \dots$$
(5.1)

where the ellipses represent higher twist contributions $\tilde{\mathcal{F}}_{k}^{A,\tau\geq 4}(x_{A},Q^{2})$, and a sum over i = q, g species is understood. As in the previous section, we use the tilde notation to identify the nuclear PDFs

 $\tilde{f}_i^A(x_A, Q^2)$, and the rescaled PDFs are without the tilde $f_i^A(x_N, Q^2)$. Here, we use the shorthand

$$\tilde{\mathcal{F}}_{k}^{A}(x_{A}) = \left[\tilde{F}_{1}(x_{A}), \tilde{F}_{2}(x_{A})/x_{A}, \tilde{F}_{3}(x_{A})\right] \qquad (k = 1, 2, 3).$$
(5.2)

The nuclear PDFs are defined as Fourier transforms of matrix elements of local, twist-2 operators (\mathcal{O}) composed of quark and gluon fields:

$$\tilde{f}_i^A(x_A, Q^2) \sim \langle A(p_A) | \mathcal{O}_i | A(p_A) \rangle,$$
(5.3)

where $x_A \in [0,1]$ and the factorization and renormalization scales have been identified with the exchange boson virtuality $\mu_F = \mu_R = Q$.

For a spin j target there are 2(2j + 1) independent helicity-dependent quark distributions $\tilde{q}_{+}^{m}(x)$ and $\tilde{q}_{-}^{m}(x)$ with $-j \leq m \leq j$. Here $\tilde{q}_{+}^{m}(x)$ ($\tilde{q}_{-}^{m}(x)$) describes the number density for finding a quark with momentum fraction x and spin m parallel (anti-parallel) to the the z-axis of a hadron moving with infinite momentum along the z-axis. Due to parity invariance of the strong interaction, we have a relation between the up (parallel) and down (anti-parallel) spin ($\tilde{q}_{+}^{m} = \tilde{q}_{-}^{-m}$) such that there remain only 2j + 1 independent distributions. However, this symmetry is broken once weak interaction effects are taken into account.

In this paper we focus on unpolarized DIS. However, the application to polarized DIS are relatively straightforward. For a spin-j target, the unpolarized quark densities are related to the helicity-dependent distributions as follows:

$$\tilde{q}_i^A(x_A, Q^2) \coloneqq \frac{1}{2j+1} \sum_{m=-j}^j (\tilde{q}_+^m + \tilde{q}_-^m).$$
(5.4)

For a spin-1/2 target (and using $\tilde{q}_{+}^{1/2} = \tilde{q}_{-}^{-1/2}$), this reduces to the well-known expression:

$$\tilde{q}_i^A(x_A, Q^2) \coloneqq \tilde{q}_+^{1/2} + \tilde{q}_-^{1/2}.$$
 (5.5)

Similar definitions can be written down for the unpolarized gluon distribution.

5.1. Nuclear DGLAP Evolution

The unpolarized nuclear PDFs (for any target spin) satisfy the usual Dokshitzer-Gribov-Lipatov-Altarelli-Parisi (DGLAP) evolution equations [95–97]

$$\frac{d\tilde{f}_i^A(x_A, Q^2)}{d\ln Q^2} = \frac{\alpha_s(Q^2)}{2\pi} \int_{x_A}^1 \frac{dy_A}{y_A} P_{ij}(y_A) \tilde{f}_j^A(x_A/y_A, Q^2)$$
(5.6)

$$\equiv \frac{\alpha_s(Q^2)}{2\pi} \int_{x_A}^1 \frac{dy_A}{y_A} P_{ij}(x_A/y_A) \,\tilde{f}_j^A(y_A, Q^2) \quad . \tag{5.7}$$

Furthermore, the following sum rules due to charge, baryon number, and momentum conservation are satisfied:

$$\int_{0}^{1} dx_A \ \tilde{u}_v^A(x_A, Q^2) = 2Z + N \,, \tag{5.8a}$$

$$\int_{0}^{1} dx_A \, \tilde{d}_v^A(x_A, Q^2) = Z + 2N \,, \tag{5.8b}$$

$$\int_{0}^{1} dx_A \ x_A \sum_{i} \tilde{f}_i^A(x_A, Q^2) = 1 \,, \tag{5.8c}$$

where Z is the electric charge of the nucleus with baryon number A = Z + N.

5.2. Relation to the OPE

A Mellin transformation can turn a convolution integral $(A \otimes B)(x) = \int_x^1 dy/y A(y)B(x/y)$ into an ordinary product of Mellin moments:

$$\int_0^1 dx x^{n-1} (A \otimes B)(x) \equiv (A \otimes B)_n = A_n \cdot B_n , \qquad (5.9)$$

where

$$A_n \equiv \int_0^1 dx x^{n-1} A(x)$$
 and $B_n \equiv \int_0^1 dx x^{n-1} B(x)$. (5.10)

Applying this to structure functions we obtain:

$$\int_0^1 dx_A x_A^{n-1} \tilde{\mathcal{F}}_k^A(x_A, Q^2) = \sum_n \tilde{f}_i^{A,n}(Q^2) C_{k,i}^n .$$
(5.11)

where on the RHS the Mellin transformation has yielded a simple sum of Mellin moments. (Note, we indicate the Mellin moment by the "n" superscript, as the tilde-notation has already been used for the pre-scaled PDFs as we discuss in the following section.)

The relation between the structure functions in the parton model and the OPE can now be easily seen by comparing the Mellin-moment of (5.1) with the j=0-term in (3.22):

$$\tilde{\mathcal{F}}_{k}^{(0),n} = C_{k,i}^{n} \tilde{A}_{i}^{n} = C_{k,i}^{n} \tilde{f}_{i}^{A,n}(Q^{2}) , \quad (k = 1, 2, 3), \quad (5.12)$$

where a sum over i = q, g is understood. From this equation we see that the matrix elements \tilde{A}^n of Eq. (3.22) are simply the Mellin moments of the parton distribution functions:

$$\tilde{A}_{i}^{n}(Q^{2}) = \tilde{f}_{i}^{A,n}(Q^{2}) \equiv \int_{0}^{1} dx_{A} x_{A}^{n-1} \tilde{f}_{i}(x_{A}, Q^{2}) .$$
(5.13)

Similarly, the Wilson coefficients $C_{k,i}^n$ of (5.12) are directly identified with those of Eq. (3.22).

5.3. Rescaling

The parton distributions $\tilde{f}_i^A(x_A)$ are number densities defined on the interval $x_A \in [0,1]$ such that $\tilde{f}_i^A(x_A)dx_A$ can be interpreted as the number of partons "i" carrying a fraction of the parent hadron momentum in the interval $[x_A, x_A + dx_A]$. We define rescaled parton densities $f_i^A(x_N, Q^2)$ in the variable $x_N \in [0, A]$ with $x_N = Ax_A$ by imposing that the number of partons remains unchanged in

the corresponding momentum intervals:

$$f_i^A(x_N, Q^2)dx_N \coloneqq \tilde{f}_i^A(x_A, Q^2)dx_A .$$

$$(5.14)$$

This equation may appear a bit odd as $x_N \in [0, A]$ on the LHS, while $x_A \in [0, 1]$ on the RHS; we will provide additional clarification in the following discussion. Additionally, it should be noted that the rescaled nuclear PDF $f_i^A(x_N, Q^2)$ are the ones usually used in the literature as $f_i^A(x_N, Q^2)$ can be meaningfully compared across different nuclei, in contrast to $\tilde{f}_i^A(x_A, Q^2)$.

It can be easily verified that the rescaling defined here at the level of the PDFs is consistent with the rescaling introduced above at the hadronic level.

Correspondingly, the rescaled objects satisfy analogous DGLAP evolution equations:

$$\frac{df_i^A(x_N, Q^2)}{d\ln Q^2} = \frac{\alpha_s(Q^2)}{2\pi} \int_{x_N}^A \frac{dy_N}{y_N} P_{ij}\left(\frac{x_N}{y_N}\right) f_j^A(y_N, Q^2).$$
(5.15)

Furthermore, the sum rules take the following form after the rescaling:

$$\int_{0}^{A} dx_{N} \ u_{v}^{A}(x_{N}, Q^{2}) = 2Z + N, \qquad (5.16a)$$

$$\int_{0}^{A} dx_N \ d_v^A(x_N, Q^2) = Z + 2N, \qquad (5.16b)$$

$$\int_{0}^{A} dx_{N} \ x_{N} \sum_{i} f_{i}^{A}(x_{N}, Q^{2}) = A.$$
(5.16c)

Note the above integrations extend to A since $x_N \in [0, A]$.

While the rescaled nuclear PDFs are formally defined on the interval $x_N \in [0, A]$, it is important to note how the bulk of the PDFs are distributed. For a proton with three valence quarks, we generally expect the valence PDFs to be peaked in the region of $x_A = x_N \sim 1/3$. In a similar manner, for nuclei of mass A, we generally expect the valence PDFs to be peaked in the region of $x_A \sim 1/(3A)$, or equivalently $x_N \sim 1/3$, which is independent of A and thus facilitates a meaningful comparison between different nuclei. Therefore, even though x_N can in principle span the range [0, A], we expect the dominant support of the PDFs to be in the region $x_A \leq 1/A$, or $x_N \leq 1$. For x_N to exceed unity, one parton alone would essentially need to acquire more momentum than an average nucleon; this is highly unlikely. Thus, it is common to assume the probability for $x_N > 1$ to be very small, and global analyses of nPDFs typically implement the condition that $f_i^A(x_N, Q^2) = 0$ for $x_N > 1$. This induces a very small error, and we discuss the possibility of $x_N > 1$ further in Sec. 5.5.

Setting $f_i^A(x_N, Q_0^2) = 0$ for $x_N \ge 1$ at the initial scale Q_0 yields an important computational benefit since the DGLAP evolution framework consistently leads to $f_i^A(x_N, Q^2) = 0$ for $x_N \ge 1$ at all larger scales Q. Consequently, the advantage of setting $f_i^A(x_N \ge 1, Q^2) = 0$ is that the same evolution equations can be used for all nuclei, including nucleons:

$$\frac{df_i^A(x_N, Q^2)}{d\ln Q^2} = \begin{cases} \frac{\alpha_s(Q^2)}{2\pi} \int_{x_N}^1 \frac{dy_N}{y_N} P_{ij}\left(\frac{x_N}{y_N}\right) f_j^A(y_N, Q^2) &: 0 < x_N \le 1 ,\\ 0 &: 1 < x_N \le A . \end{cases}$$
(5.17)

With the above framework for the nuclear PDFs, we can perform the DGLAP evolution on the interval $x \in [0, 1]$ in a manner similar to the proton evolution. However, the sum rules of Eq. (5.16a), (5.16b), and (5.16c) differ from the proton case. Thus, it is common (but not necessary) to further



Figure 5.1: We display the nCTEQ15 nPDFs [17] as a function of x at Q = 10 GeV for selected nuclear A values as indicated in the legend. These are the scaled nPDFs of Eq. (5.18), and this feature is most evident when comparing the xu(x,Q) and xd(x,Q) of the proton (A = 1).

decompose the nuclear PDF f_i^A as

$$f_i^A(x,Q) = \frac{Z}{A} f_i^{p/A}(x,Q) + \frac{A-Z}{A} f_i^{n/A}(x,Q) , \qquad (5.18)$$

where $f_i^{p/A}$ and $f_i^{n/A}$ represent effective "bound proton and neutron" PDFs for finding a parton "*i*" inside a nucleon. The "bound proton" PDF $f_i^{p/A}$ satisfies identical sum rules as the (free) proton PDF, so this quantity can be computed using standard DGLAP programs. The "bound neutron" PDF is commonly obtained from the "bound proton" one by assuming isospin symmetry.⁸

⁸While isospin symmetry is typically used to relate $f_i^{n/A}$ to $f_i^{p/A}$, in principle one could add isospin violating contributions. The recent Marathon data [98] suggests there may be isospin violations in the very large x region [99].

For comparison of nPDFs across different A nuclei, it is typically the scaled nPDF $f_i^{(A,Z)}(x,Q)$ of Eq. (5.18) which can be meaningfully compared, as we illustrate in Fig. 5.1. This quantity is defined on $x \in [0,1]$ and normalized so that the integrated momentum is unity. Conversely, although the effective "bound proton and neutron" PDFs of Eq. (5.18) are convenient for computing the DGLAP evolution, these are *not* physically measurable objects. Caution is required when comparing them.

In particular, if we try to invert the relations of Eq. (5.18) for the up and down quarks, we find:

$$u^{p/A} = \frac{Z}{2Z - A} u^A - \frac{A - Z}{2Z - A} d^A \quad \text{and} \quad d^{p/A} = \frac{Z}{2Z - A} d^A - \frac{A - Z}{2Z - A} u^A \quad .$$
(5.19)

This relation has the expected limits that for a proton with A = p and $\{A, Z\} = \{1, 1\}$ we find $\{u^{p/A}=u^{A=p}; d^{p/A}=d^{A=p}\}$. Also, for a neutron with A = n and $\{A, Z\} = \{1, 0\}$ we find $\{d^{p/A}=u^{A=n}; u^{p/A}=d^{A=n}\}$. However, for an isoscalar target (A = 2Z), Eq. (5.19) is singular because in the isoscalar limit it is impossible to differentiate the u^A and d^A distributions and the decomposition into $u^{p/A}$ and $d^{p/A}$ becomes arbitrary. Since the majority of the nuclear data is taken on nuclear targets which lie along the nuclear stability line in the $\{A, Z\}$ plane where $A \sim 2Z$ (except for very heavy nuclei), our ability to separately determine the nuclear u^A and d^A distributions is limited.

5.4. Kinematic W Cut

Determinations of PDFs are performed by global analyses using a wide array of data sets that can be described by the parton model within the factorization framework. It is therefore important to exclude data in the kinematic region that are not reliably predicted within this model.

In lepton-nucleon scattering, there are different types of scattering scenarios depending on the value of the hadronic invariant mass, $W^2 = (p+q)^2$, and the value of the virtuality of the exchanged boson Q [100]. We can divide these cases into:

- 1. elastic scattering $(W = M_N)$,
- 2. shallowly inelastic scattering (SIS) $(W \leq 2 \text{ GeV})$,
- 3. soft DIS (W > 2 GeV, Q < 1 GeV), and
- 4. DIS $(W > 2 \text{ GeV and } Q \ge 1 \text{ GeV})$.

Among these scenarios, the DIS and soft DIS (dominated by non-resonant pion production) cases are the most reliably described by the parton model. Therefore, a W cut is traditionally imposed on DIS data to extract PDFs.

While imposing a W cut is straightforward for the case of lepton-proton scattering, it is more subtle for lepton-nucleus scattering when we compute in terms of the rescaled variables. For $\ell_1 + A \rightarrow \ell_2 + X$ scattering, we have:

$$W_A^2 = (p_A + q)^2 = M_A^2 + Q^2 \frac{1 - x_A}{x_A} = A^2 M_N^2 + Q^2 \frac{1 - x_N/A}{x_N/A} \quad .$$
 (5.20)

The average W_A per nucleon, $W_{aver} = W_A/A$, is then given by

$$W_{\rm aver}^2 = \left(\frac{W_A}{A}\right)^2 = M_N^2 + Q^2 \frac{A - x_N}{A^2 x_N} \quad .$$
 (5.21)



Figure 5.2: We display the neutron structure function F_2^n as measured by the BONuS data [101] at selected Q^2 values. The open (filled) circles represent data for a beam energy of E = 4.223 (5.262) GeV. This is compared with ABKM PDF predictions [94] including higher twist effects and target mass corrections. We observe the resonance structure at low W^2 values. Figure taken from Ref. [101].

Additionally, we observe that W^2_{aver} and W^2_N are not the same:

$$W_N^2 = (p_N + q)^2 = M_N^2 + Q^2 \frac{1 - x_N}{x_N} \quad .$$
 (5.22)

Note, that for $x_N \in [0, A]$, we find W_{aver}^2 is always positive while W_N^2 can be negative for $x_N > 1$. On the other hand, W_N is independent of A and can be used to compare different nuclei as long as x_N is restricted to values below unity.

If we impose a lower bound on W_A , this translates to the well-known upper bound on x_A of

$$0 \le x_A \le \left[1 + \frac{W_A^2 - M_A^2}{Q^2}\right]^{-1} \le 1.$$
(5.23)

In terms of the rescaled variable $x_N = Ax_A \in [0, A]$ this translates to

$$0 \le x_N \le A \left[1 + \frac{W_A^2 - M_A^2}{Q^2} \right]^{-1} \le A.$$
(5.24)

Consequently, when we impose a W kinematic cut for the nuclear case (A > 1), we need to be cognizant of these issues. One could try to limit the resonance region with a mass cut (m_{cut}) by setting $W_N^2 = M_N^2 + m_{\text{cut}}^2/A^2$. For A > 1, the resulting high-x cut turns out to be more relaxed than the usual W cut. Thus, the use of traditional W cut here is much safer and hence, despite its pathological nature at x > 1, this cut can still be used. While cutting on W_A or (W_A/A) is more natural, the value of W_N , which is A dependent, needs further investigations.



Figure 5.3: We display a characteristic nuclear correction ratio F_2^{Fe}/F_2^D for iron over deuterium. We observe the shadowing region $(x \leq 0.1)$, the anti-shadowing region $(x \sim 0.2)$, the EMC region $(x \sim 0.5)$, and the Fermi region $(x \geq 0.8)$. The figure is taken from Ref. [65] which contains the details and references for the data sets.

5.5. nPDFs for $x_N \ge 1$

The rescaled PDFs allow us to compare distributions from different nuclei in a meaningful manner as illustrated in Fig. 5.1. Operationally, all current global nuclear PDF analyses work on the interval $x_N \in [0, 1]$ even though the technically allowed range is $x_N \in [0, A]$. Although the details of the $x_N > 1$ region are beyond the scope of this study, we do want to briefly mention experimental measurements that provide insights in this extreme kinematic region.

The Fermi Region

The effects of nuclear binding can modify the partonic momentum, and thus dramatically impact the resulting structure functions. Fig. 5.3 shows the characteristic form of the nuclear correction factor F_2^{Fe}/F_2^D with shadowing at small x_N ($x_N < 0.1$), anti-shadowing ($x_N \sim 0.1$), the EMC region for intermediate x, and the Fermi region for large x. The nuclear binding effects are especially evident at larger momentum fractions (large x) in the Fermi region where the PDFs are steeply falling. There are a variety of theoretical approaches to describe this region, as well as experimental measurements of these nuclear binding effects. We provide only a brief overview below and refer to the reference for additional details.

Challenges of the Fermi Region

The limitations of constraining the PDFs in the $x_N \sim 1$ region are evident when examining the nuclear ratio F_2^A/F_2^p in the large x_N region. As x increases, the $\mathcal{R} = F_2^A/F_2^p$ ratio transitions from $\mathcal{R} \leq 1$ in the EMC region, to $\mathcal{R} \gtrsim 1$ as $x_N \to 1$ in the Fermi region.

We find that $F_2^p(x)$ with A = 1 for the proton must vanish at $x_N = 1$, whereas $F_2^A(x)$ can be finite since $x_N \in [0, A]$. This means that the denominator of \mathcal{R} is vanishing while the numerator is finite. Hence, \mathcal{R} will rapidly increase at $x \sim 1$, and this is consistent with experimental measurements. More generally, we expect $F_2^{A_1} > F_2^{A_2}$ for $A_1 > A_2$, and this is the case for $F_2^{12}C/F_2^D$ as shown in Fig. 5.4.



Figure 5.4: We display the ratio F_2^A/F_2^d for selected nuclei data sets in the large x region. The different curves show the impact of the ε parameter for the rescaling $x'_N \to [x_N - \varepsilon x_N^\kappa \log_{10} A]$ with $\kappa = 10$. This figure is taken from Ref. [19].

Theoretical Implementations

On the theoretical side, there are a number of approaches that have been proposed to model the nPDFs into the $x_N > 1$ region, and we point to a selection of references [11, 37, 102–110]. Additionally, there is a LO solution of DGLAP equation for quarks taking into account the full support (including $x_N > 1$ region) which is detailed in Ref. [111].

One technique to describe the $x_N > 1$ region is to use a convolution of the nucleon PDFs f(x,Q) defined on the interval $x \in [0,1]$ with a smearing kernel S_A which will shift the PDFs out to larger x values:

$$f_A(x,Q) = \int_x^A \frac{dy}{y} S_A(y,Q) f(x/y,Q) \quad .$$
 (5.25)

Here, the smearing kernel S_A is typically a Gaussian with an A-dependent width; larger nuclei have a larger smearing width, and hence a larger proportion of the f_A distribution will populate the x > 1region. The behavior of the PDF convolution can also be approximated using an A-dependent rescaling: $x'_N \rightarrow [x_N - \varepsilon x_N^{\kappa} \log_{10} A]$, c.f., Ref. [19]. This mimics the PDF convolution method described above, but the rescaling avoids the convolution integral, so it is computationally fast to evaluate.

In Fig. 5.4 we display F_2^A/F_2^d for selected nuclei and data sets, and indicate various levels of rescaling (controlled by the ε parameter). The data points are described in Ref. [19], and the solid curve (blue) with ε =0 is the default theory with no x-rescaling. The solid points are within the kinematic cuts of the global fit (Q > 1.3 GeV, $W_N > 1.7$ GeV), and the hollow points are outside these cuts. As we increase the ε parameter, we see this increases the ratio in the large x region, and can improve the comparison of the data and theory.

Judging from this sample of data, it appears that a detailed modeling of the Fermi region can improve the description of the data in this extreme kinematic region (low W_N , low Q, large x_N). These corrections could be a combination of TMCs, $x_N > 1$ effects, higher twist, or other non-perturbative corrections. Thus, a complete analysis must include and balance all these contributions.

Experimental Investigations

There are a number of studies that have explored the structure functions in the Fermi region at very large x_N , and we mention a few examples below.

For example: Refs. [112, 113] examined SLAC DIS data from a variety of nuclear targets and found that in the larger x_N region ($x_N \ge 0.5$) the Fermi motion effects are similar to those in the deuteron, and increase for heavier nuclei. Measurements in this region are particularly challenging, and are subject to a variety of uncertainties as outlined in the Appendix of Ref. [112].

For the CC neutrino DIS process, CCFR [114] measured ~2000 events in the region $x_N > 0.75$, and used this to study F_2 at very large x_N . They fit the high x_N behavior to a decaying exponential $\propto \exp(-sx_N)$ normalized at $x_N = 0.65$, and find a slope of $s = 8.3 \pm 0.7$. They note this compares favorably with E133 at SLAC [115], which observed $s \sim [7,8]$.

Hall C at Jefferson Lab measured the NC structure function F_2 on a variety of nuclei ranging from ²H and ³He up to ¹⁹⁷Au in the large x_N region [116, 117] out to $x_N \sim 1.4$. This work extends an earlier study of Ref. [118], and related measurements performed at SLAC [119]. Taking account of the kinematic pre-factors and the $\{h_2, g_2\}$ contributions, they extract $F_2^{(0)}(\xi_N, Q^2)$, where ξ_N is the Nachtmann variable, and fit this to a decaying exponential of the form $\exp(-s\xi_N)$ in the region $\xi_N \geq 0.75$. The large ξ_N fits finds the slope is in the general range $s \sim 15 \pm 2$. The slope has a mild increase with increasing Q^2 , and a mild decrease with increasing nuclear A. These results also compare favorably with the BCDMS muon scattering data [120] which finds $s = 16.5 \pm 0.6$ for $x_N \sim [0.75, 1.05]$.

Parton Momentum in the Fermi Region

Based on the above large x_N measurements, we can ask the question: How much parton momentum is in the region $x_N > 1$?

The experimental measurements discussed above parameterize F_2 at large x_N as $\exp[-sx_N]$, where s can range from about ~8.3 to ~14. With s=8.3, the fraction of F_2 in the range $x_N \in [1, \infty]$ is less than ~6% that in the range $x_N \in [0.65, 1]$. Given that the average momentum fraction in the interval $x_N \in [0.65, 1]$ is $\leq 2\%$ of the total nucleon momentum, this gives a rough estimate of the potential momentum fraction beyond $x_N > 1$ of $6\% \times 2\% \leq 0.1\%$ of the total momentum fraction. For $s \sim 14$, the decrease at large x_N is even steeper; here, the F_2 ratio of $x_N \in [0.65, 1]$ compared to $x_N \in [1, \infty]$ drops to $\leq 0.1\%$, and the potential momentum fraction beyond $x_N > 1$ estimate becomes $1\% \times 2\% \leq 0.02\%$ of the total.

In both of the above cases, the integrated parton momentum is very small in the $x_N > 1$ region, but the impact on the $F_2^{A_1}/F_2^{A_2}$ ratio still yields the characteristic increase in the $x_N \sim 1$ Fermi region.

5.6. Threshold problem and higher twist contributions

On the basis of the correspondence between the limits $x_N \to 1$ and $W_N^2 \to M_A^2$, the very high x_N behavior of DIS structure functions at fixed Q^2 is sensitive to the presence of power-suppressed corrections beyond leading twist (twist-4 for unpolarized processes). It has been conjectured that the problematic threshold behavior in target-mass corrected structure functions in the free nucleon case are at least partly due to the absence of higher-twist contributions. As an approximate test of the inclusion of twist-4 contributions, Ref. [121] proposed a $1/Q^2$ expansion of the OPE mass-corrected structure functions, which indeed suppresses the problematic threshold behavior. These calculations were done for F_2 , and later, for other tensor structure function in Ref. [122]. While this was done for the free nucleon, a similar exercise might be carried out for nuclear structure functions as another TMC prescription. In particular, Kulagin and Petti [121] showed that by expanding the target mass corrected structure functions to leading order in $1/Q^2$, the resulting functions have the correct $x_N \rightarrow 1$ limits. While avoiding the threshold problem, this prescription, however, raises the question of whether the $1/Q^2$ approximation is sufficiently accurate for structure functions near $x_N \approx 1$ at moderate Q^2 .

To test the convergence of the $1/Q^2$ expansion at large x, Ref. [122] further expands the OPE result(s) to include $\mathcal{O}(1/Q^4)$ corrections. In fact, one can demonstrate that for a structure function that behaves at large x as $(1-x)^n$, the target mass corrected result will vanish in the $x_N \to 1$ limit up to order $1/Q^{2n-2}$ in the expansion. For $n \approx 3$, as is typical for nucleon structure functions, the threshold problem will therefore appear only at order $1/Q^6$.

6. The parton model with quark and hadron masses

We now turn to a discussion of the parton model, including the effects of quark and hadron mass. Such mass effects can be rigorously included in the QCD-improved parton model, as shown in a general factorization proof with quark masses by Collins [46]. The underlying scheme is also known in the literature as the Aivazis-Collins-Olness-Tung (ACOT) scheme [57, 58, 123], which is an example of what is called a General Mass Variable Flavor Number Scheme (GM-VFNS) in the modern literature [58, 91–94, 124]. In the following subsections we summarize DIS structure functions in the ACOT scheme, including quark and hadron (nuclear) mass effects, and compare and contrast these expressions to the structure functions in the OPE. As discussed in the introduction (Sec. 1), we consider the theoretical status of both approaches, the OPE and the collinear-factorization-based QCD-improved parton model, on equal footing.

In the naïve, massless parton model, we have a hadron of momentum p_A that emits a collinear parton of momentum $k = x_A p_A$. If we try and extend this simple picture to the case of massive partons, the presence of the parton mass m_i violates the collinear kinematics and we encounter ambiguities proportional to the parton mass. We work in the ACOT formalism [57, 58, 123] to compute the structure functions in the helicity basis. This has the advantage that the polarization vectors are boost invariant between the hadron and parton frame, and provides advantages for incorporating both the hadron and parton masses, as we shall illustrate below. We observed in Sec. 4 that the hadron mass modifies the scaling variable via the relation $\xi_A = R_M x_A$, where the combination $R_M x_A$ is the Nachtmann variable. Within the context of the OPE and the parton model, the effect of the quark masses factorizes from the hadron modification. The rescaling variable is modified by an additional factor R_{ij} , which is a function of the quark masses $\{m_i, m_j\}$ and is defined in Table 4.2. Thus, we have $\xi_A = R_{ij} R_M x_A$.

In the following, we also review the structure functions in the light-front formalism, and compare with the TMC expressions obtained from the OPE. There has been extensive discussion in the literature about computations with massive partons [58, 91–94]. The present discussion is focused on the detailed organization of the massive parton contributions in the formalism outlined by Collins [46]. This work extended the factorization proof to the case of massive partons and demonstrated that all contributions proportional to the parton mass can be fully factorized. Thus, we include all terms of order $\mathcal{O}(m_i/Q)^2$, and the non-factorizable terms are suppressed by powers of $(\Lambda_{\rm NP}/Q)^2$.

6.1. Helicity Formulation of the Parton Model

When we generalize the massless QCD parton model to the massive case, there are potential ambiguities proportional to the parton mass m_i , which may enter if we are not careful about organizing our perturbative expansion. This organization is more easily dealt with in the helicity basis as we outline. The original ACOT formalism for massive DIS was computed in the helicity basis, and this provides some advantages in working with both hadron and parton masses. For example: in the helicity formalism, we can define polarization vectors (Sec. 6.2) in a boost-invariant manner so that they are the same for both the hadron and parton reference frames (Sec. 6.3). This has the advantage that there is no mixing of terms between the partonic and hadronic structure functions in the helicity basis. Additionally, by working with light-front momenta, $\{p^{\pm}, k^{\pm}\}$, we can take advantage of the fact that $p^{+} \gg p^{-}$ and $k^{+} \gg k^{-}$, and define our scaling variable ξ as $k^{+} = \xi p^{+}$, which is valid for both massless and massive partons.

For the present discussion, we will focus on the target mass corrections which enter the hadronic structure functions. We will find the helicity formalism matches the OPE master formula for the leading terms. We compare these result both analytically in Sec. 6.4 and numerically in Sec. 6.5 (and Sec. 7).

6.2. Helicity Decomposition

In analogy to what was done in Sec. 2, we decompose the hadronic tensor $W_{\mu\nu}$ by projecting out Lorentz tensors to obtain the six structure functions F_i . We can also project out the helicity components using polarization vectors $\epsilon_n^{\nu}(p,q)$:

$$F_{\lambda} \equiv F_{mn} = \epsilon_m^{\mu *}(p,q) W_{\mu\nu}(p,q) \epsilon_n^{\nu}(p,q) .$$
(6.1)

Here, helicity indices m, n span the helicity components plus (+), minus (-), longitudinal (0), and scalar (s). Angular momentum conservation reduces these to six combination {++, --, 00, ss, 0s, s0}, just as we have six hadronic structure functions in the tensor basis F_i with {i = 1...6}. Additionally, the three combinations { F_{ss}, F_{s0}, F_{0s} } are suppressed by the lepton mass in a manner analogous to { F_4, F_5, F_6 }, so typically these are ignored. The remaining combinations are { F_{++}, F_{00}, F_{--} }, which we also indicate with the short-hand notation F_{λ} with $\lambda = \{+, 0, -\}$.

We obtain the hadron-level cross section by contracting the hadronic and leptonic tensors:

$$d\sigma \sim L^{\mu\nu}(q,k) \ W_{\mu\nu}(p,q) \equiv L^{m'n'}(q,k) \ d_{m'}^{m}(\psi)^{-1} \ d_{n'}^{n}(\psi) \ F_{mn}(p,q) \ ,$$
(6.2)

where we show the corresponding relations in both the tensor and helicity basis. Here the Wigner rotation matrix $d_{n'}^n(\psi)$ is the SO(2,1) analog of the familiar 3×3 "rotation" matrix (since the exchanged particle is spin-1 with 3 polarization states), and ψ is the boost angle that transforms between the hadronic Breit ("brick wall") frame, where p || q, and the leptonic frame where k || k', with

$$\cosh \psi = \frac{2p \cdot (k+k')}{\Delta [-Q^2, p^2, p_x^2]} = \frac{\xi^2 M^2 - Q^2 + 2\xi (s-M^2)}{\xi^2 M^2 + Q^2} \xrightarrow[M \to 0]{} \frac{2-y}{y} \quad .$$
(6.3)

Here, $\Delta^2[a, b, c] = a^2 + b^2 + c^2 - 2(ab + bc + ca)$. An elegant feature of Eq. (6.2) is that the leptonic (L^{mn}) and hadronic (F_{mn}) helicity structure functions are simple and all the kinematic complications due to the target mass are contained in the $d(\psi)$ rotation matrices. This simplicity is a consequence of the underlying group-theoretic approach to the factorized structure.⁹

 $^{^{9}}$ While the helicity formalism elegantly exhibits the underlying symmetries of the interaction, as we go to higher-order the computation of the Wigner rotations for multiple intermediate particles can become more complex [58].

	TMC	ACOT-TMC	ACOT-TMC TMC
F_1	$\frac{x}{\xi r}$	1	$\frac{\xi r}{x} \sim 1 + \varepsilon^2$
F_2	$\frac{x^2}{\xi^2 r^3}$	$\frac{x}{\xi r^2}$	$\frac{\xi r}{x} \sim 1 + \varepsilon^2$
F_3	$\frac{x}{\xi r^2}$	$\frac{1}{r}$	$\frac{\xi r}{x} \sim 1 + \varepsilon^2$

Table 6.1: Tabulation of prefactors from the master formulas for the TMC (obtained using the OPE) of Eq. (3.24) and ACOT-TMC light-front derivations of Eq. (6.10), where $\varepsilon = (xM/Q)$, and $r = r_A = r_N = \sqrt{1 + 4\varepsilon^2}$. While the F_i 's are generally evaluated at x_N , using Table 4.1 we have $x_N/\xi_N = x_A/\xi_A$ and these ratios are equivalent. The TMC and ACOT-TMC prefactors match to $\mathcal{O}(\varepsilon^2)$, and this is plotted in Fig. 6.1. The source of this factor is discussed in the text.

6.3. Boost Invariant Polarizations

If we work in light-front coordinates $\{x^+, \vec{x}, x^-\}$ with $x^{\pm} = (x^0 \pm x^3)/\sqrt{2}$ and $\vec{x} = \{x^1, x^2\}$, then

$$p^{\mu} = \left\{ p^{+}, \vec{0}, \frac{M^{2}}{2p^{+}} \right\} \quad \text{and} \quad q^{\mu} = \left\{ -\xi p^{+}, \vec{0}, \frac{Q^{2}}{2\xi p^{+}} \right\} , \quad (6.4)$$

such that $p^2 = M^2$ and $q^2 = -Q^2$. Additionally, $2p \cdot q = Q^2/x = Q^2/\xi - \xi M^2$. We now choose to define the parton momentum for a massive state as

$$k^{\mu} = \left\{ \xi p^{+}, \vec{0}, \frac{m^{2}}{2\xi p^{+}} \right\} , \qquad (6.5)$$

such that $k^2 = m^2$ and $k^+ = \xi p^+$. This scaling relation holds for both massless and massive partons.

We require two reference vectors to define the polarizations. For the hadron polarizations $\epsilon_n^{\nu}(p,q)$ we choose the vectors $\{p,q\}$, while for the parton polarizations $\epsilon_n^{\nu}(k,q)$ we choose the vectors $\{k,q\}$. For both the hadron and parton polarizations, the scalar polarization is given by $\epsilon_0^{\mu} = q^{\mu}/Q$, and the transverse polarizations are defined as usual to be $\epsilon^{\pm} = (0, \pm 1, -i, 0)/\sqrt{2}$. The longitudinal polarizations are constructed from the reference vectors using a completeness relation [57]. The polarization vectors only depend on the reference vectors p^{μ} and k^{μ} to the extent that they define the t - z plane in conjunction with q^{μ} ; see Ref. [57] for details.

Thus, the polarization vectors are invariant between the hadron and parton reference frame. Additionally, there is a direct relation between the hadronic helicity structure functions F_{λ} and the partonic helicity structure functions ω_{λ} :

$$F_{\lambda} = \delta^{\lambda\lambda'} f \otimes \omega_{\lambda'} \quad . \tag{6.6}$$

Here, f is the PDF, \otimes represents a convolution, $\delta^{\lambda\lambda'}$ is the usual Kronecker δ -function, and λ sums over the six helicity configurations. Specifically, we note there is no mixing as indicated by the $\delta^{\lambda\lambda'}$ function. This is in contrast to the structure functions in the tensor basis:

$$F_i = c^{ij} f \otimes \omega_j \tag{6.7}$$

where the mixing coefficients c^{ij} contain off-diagonal elements, with indicies i, j summing over the six tensor structure functions [57].

6.4. Relation to Master Formulas

Finally, coming back to the hadronic level, we can relate the hadronic helicity structure functions F_{λ} to the tensor structure functions F_i using Eq. (6.1) and obtain:¹⁰

$$F_1 = \frac{1}{2} \{F_+ + F_-\} , \qquad F_2 = \frac{x}{r^2} \{F_+ + F_- + 2F_0\} , \qquad F_3 = \frac{1}{r} \{-F_+ + F_-\} , \qquad (6.8a)$$

and the inverse relations

$$F_{\pm} = F_1 \mp \frac{r}{2} F_3 , \qquad F_0 = -F_1 + \frac{r^2}{2x} F_2 , \qquad (6.8b)$$

where $r = r_A = r_N = \sqrt{1 + 4\varepsilon^2}$ and $\varepsilon = (xM/Q)$. The F_0 helicity structure function is related to the longitudinal structure F_L of Eq. (3.26) via

$$F_L = 2x F_0 = r^2 F_2 - 2x F_1 . ag{6.9}$$

Translating the above relations into the master formula for the TMCs for the light-front approach yields:

$$F_1^{\text{ACOT-TMC}}(x) = F_1^{(0)}(\xi)$$
 (6.10a)

$$F_2^{\text{ACOT-TMC}}(x) = \frac{x}{\xi r^2} F_2^{(0)}(\xi)$$
(6.10b)

$$F_3^{\text{ACOT-TMC}}(x) = -\frac{1}{r} F_3^{(0)}(\xi) \quad , \qquad (6.10c)$$

and $F_L^{\text{ACOT-TMC}}(x)$ can be constructed using Eq. (3.26). In the above formula, we can incorporate the parton masses by using the full rescaling variable $\xi_A = R_{ij} R_M x_A$, and computing $F_i^{(0)}(\xi)$ including the quark mass effects. The proof of factorization for heavy quarks is detailed in Ref. [46].

The correspondence between the ACOT-TMC and TMC expressions are summarized in Table 6.1. In the right-most column, we display the relative conversion factor $(\xi r/x)$, which is the same for all three $F_i^{\text{ACOT-TMC}}(x)$ results. While we do not compute $F_{4,5,6}$, we expect these structure functions also follow this pattern. For F_i , we see that the TMC and ACOT-TMC approach align to $\mathcal{O}(\varepsilon^2)$, where $\varepsilon = (xM/Q)$ vanishes outside the dominant TMC region of large x and small Q. Note, these comparisons are for the leading term $F_i^{\text{Leading-TMC}}$ only (see below and Sec. 7), as the higher order terms in $(M/Q)^2$ with the h_i and g_i functions are not included in the ACOT expressions.

6.5. Relation of Parton Model to OPE

The parton model ACOT-TMC results differ from the TMC results obtained with OPE in two respects: i) there is an overall factor of $(\xi r/x)$, and ii) the OPE result contains additional $(M/Q)^2$ and $(M/Q)^4$ terms. We will discuss the origin of these differences below.

Correspondence of the leading factor $(\xi r/x)$: The TMC (OPE) results of Eq. (4.15) contain an overall factor of $(\xi r/x)$ as compared with the ACOT-TMC results of Eq. (6.10). The source of this factor comes from fully including the hadron mass in the calculation as outlined in Appendix A.5

¹⁰Note that while the F_i structure functions of Eq. (4.15) are evaluated at x_N , the relations of this section hold in general; hence, we omit the subscripts on x_N here.


Figure 6.1: We display the relative factor $(\xi r/x)$ between the ACOT and leading OPE terms as tabulated in Table 6.1 for $Q = \{1.3, 1.5, 2, 3, 4, 6\}$ GeV as a function of x with $M = M_{proton}$. The lower two Q values $\{1.3, 1.5\}$ are dashed to indicate these values are below typical kinematic cuts $(Q_{cut} \sim 2 \text{ GeV})$ for global PDF fits.

which computes both the massless and massive results. The relative factor is most obvious in the case of F_1 , where the ACOT-TMC result is simply given by $F_1^{(0)}(\xi)$.

In contrast, the massive OPE results of Eq. (A.146) include the additional $(\xi r/x)$ factor which arises from the delta function of Eq. (A.124). This delta function enforces the relationship between the Bjorken and Nachtmann variables, and is a manifestation of momentum conservation. The same factor is present in all the OPE results for F_i as computed in Appendix A.5, and is summarized in Table 6.1 for $F_{1,2,3}$. Additionally, this factor is also present in Ref. [83]; c.f., Eq. (1.24) where, matching notation, $(1 + \mu\xi^2) = (r\xi/x)$ with $\mu = (M/Q)^2$.

In Fig. 6.1 we show the size of this relative factor as a function of x for selected Q values. For small x and large Q this factor is close to unity, but grows for large x and small Q values where the TMCs are typically substantial.

Correspondence of the M/Q **Powers:** The OPE master equation, Eq. (3.24), includes higher powers of (M^2/Q^2) which are formally twist-2. The source of these terms comes from the intrinsic transverse momentum of the parton [55, 83]. The partonic cross sections are evaluated with parton momenta on-shell, but not collinear with the parent hadron. The details are provided in Refs. [55, 83], which include the transverse momentum in the description of the partons, [*c.f.*, $f(x, k_T^2)$ in Eq. (1.22) of Ref. [83]]. As the transverse momentum is bounded to be of order M^2 , these contributions give rise to the twist-2 (M^2/Q^2) and (M^4/Q^4) terms of Eq. (3.24). Thus, if the transverse momentum is accounted for in the parton model, one matches the OPE result.

Recap: We have derived the structure function results for the parton model using the ACOT formalism in a helicity basis as summarized in Eq. (6.10). We can obtain a complete correspondence between the parton model and OPE structure function results if i) we fully account for the hadron mass in the delta function (c.f., Eq. (A.124)), which ensures the relationship between the Bjorken and Nachtmann variables, and ii) we account for the transverse momentum of the parton (c.f., Ref. [83]).

7. Numerical results

Having discussed various theoretical and phenomenological aspects of TMCs for nuclei from an analytic perspective, we now turn our focus to the quantitative aspect of TMCs. In this section, we explore

the numerical impact of TMCs on first the proton structure functions (Sec. 7.1), and then the nuclear structure functions (Sec. 7.2). We will examine both the leading-TMC and sub-leading-TMC contributions, and explore the variation across the nuclear A range. These observations will allow us to design a simple parameterization of the TMCs which are computationally efficient (Sec. 7.3). Finally, we will compute the impact of the TMCs for DIS reduced cross sections with sample kinematics from HERA, JLab, EIC, and LBNF (Sec. 7.4).

In order to build intuition for the numerical results throughout this section, it is useful to recall several relations between structure functions and quark and antiquark PDFs, but which hold only at LO in the absence of CKM mixing and masses. In particular, for neutrino and antineutrino DIS on an arbitrary nuclear target (A), where A may also be just a proton, the charged current structure functions ($F_i^{W^{\pm}}$) and PDFs are related by:

$$F_1^{\nu A} = (d + s + \bar{u} + \bar{c}), \qquad F_1^{\bar{\nu}A} = (u + c + \bar{d} + \bar{s}), \tag{7.1a}$$

$$F_2^{\nu A} = 2x \left(d + s + \bar{u} + \bar{c} \right), \qquad F_2^{\bar{\nu} A} = 2x \left(u + c + \bar{d} + \bar{s} \right), \tag{7.1b}$$

$$F_3^{\nu A} = +2 \left(d + s - \bar{u} - \bar{c} \right), \qquad F_3^{\bar{\nu}A} = -2 \left(u + c - \bar{d} - \bar{s} \right), \tag{7.1c}$$

in the limit of four quarks. Here $\{u, d, ...\}$ are the PDFs of a full nucleus A. Likewise, when the exchange of Z bosons can be neglected in charged lepton DIS, the neutral current structure functions (F_i^{γ}) are related to PDFs at LO by

$$F_2^{l^{\pm}A} = x \frac{1}{9} \left[4(u+\bar{u}) + (d+\bar{d}) + 4(c+\bar{c}) + (s+\bar{s}) \right].$$
(7.2)

While the LO relations shown above are intuitively useful, our calculations are performed at full NLO in QCD including the quark mass contributions; specifically, we use the S-ACOT(χ) scheme.

Since we will compare separate components of the TMC contributions, to reduce ambiguity we use the following nomenclature throughout our presentation.

- **TMC:** We identify the full set of contributions to the structure function, given by the OPE master equation of Eq. (4.15), as the "TMC" result; this label is without qualifiers.
- **No-TMC:** As we take the $(M^2/Q^2) \to 0$ limit, the pre-factors of $F_i^{A,(0)}$ become unity, and the higherorder $(M/Q)^2$ and $(M/Q)^4$ terms containing the $\{h_i, g_i\}$ functions vanish.¹¹ Additionally, we have $\xi \to x$ so that the expressions in Eq. (4.15) reduce to $F_i^{A,(0)}(x)$ alone. We refer to this as the "No-TMC" result.
- **Leading-TMC:** The "Leading-TMC" structure functions are obtained from Eq. (4.15) by only keeping the first term on the RHS which is proportional to $(M/Q)^0$. Specifically, we are neglecting the terms proportional to $(M/Q)^2$ and $(M/Q)^4$ which contain the $\{h_i, g_i\}$ functions.
- **ACOT-TMC:** We obtain the "ACOT-TMC" structure functions from the ACOT master equation of Eq. (6.10). This result is similar to the "Leading-TMC," but the pre-factors differ by $(\xi r/x)$ as detailed in Table 6.1. Importantly, the "ACOT-TMC" result does not include the higher-order $(M/Q)^2$ and $(M/Q)^4$ terms containing the $\{h_i, g_i\}$ functions.

¹¹Note, we always retain the full quark mass dependence in all calculations as this is factorized from the hadron-level kinematics. C.f., Ref. [60] for details.

7.1. Proton Structure Functions with TMC

In comparison to structure functions for massless protons, structure functions with TMCs for nuclei contain two additional layers of complexity. The first, of course, is the larger nucleon content; the second are the TMCs themselves. Therefore, in order to establish a baseline intuition of TMCs for nuclear structure functions, we consider briefly TMCs for proton structure functions. For more comprehensive studies of TMCs for protons, see Ref. [60] and references therein.

We begin with Fig. 7.1, where in the upper panel we plot the absolute structure functions F_i for charged current (W^-) and neutral current (γ, Z) exchange showing "No-TMC" (solid), full "TMC" (short dash), "Leading-TMC" (dash-dot), and "ACOT-TMC" (long dash-dot), at Q = 1.5 GeV and 10 GeV as a function of the Bjorken scaling variable x. In the lower panel we show the ratio of the full "TMC" to the "No-TMC" results. The inset panel displays the ratio of the "Leading-TMC" to the "ACOT-TMC" which is simply a function of the relative ($\xi r/x$) factor discussed in Sec. 6.

 F_i Structure Functions: (Fig. 7.1 Upper Panels) We focus first on the upper panels of Fig. 7.1 which shows the proton structure functions vs. x. All the structure functions are steeply decreasing for large x. This reflects the underlying PDF structure of Eqs. (7.1) and (7.2): PDFs, and hence structure functions, must decrease at large x due to momentum conservation.

For both Q = 1.5 GeV and Q = 10 GeV, we observe that the full "TMC," "Leading-TMC" and "ACOT-TMC" results roughly coincide. The good agreement between the full "TMC" and "Leading-TMC" allows us to infer that the sub-leading contributions containing the $\{h_i, g_i\}$ terms are small; this is to be expected as they are suppressed by powers of $(M/Q)^2$ and $(M/Q)^4$. We will further examine the impact of these sub-leading terms in Sec. 7.2.

The approximate agreement of the "Leading-TMC" and the "ACOT-TMC" reflects the impact of the relative factor of $(\xi r/x)$ discussed in Sec. 6.4. To better display this difference, in the inset panels we plot the ratio of these two structure functions for the lower Q = 1.5 GeV value; at Q = 10 GeV, the difference is minimal as is evident in Fig. 6.1.

In contrast, at Q = 1.5 GeV, the "No-TMC" results are dramatically different from the other three curves. The "No-TMC" expression is most similar to the "Leading-TMC" as illustrated below for the case of F_1 :

$$F_1^{\text{TMC-leading}}(x_N) = \frac{x_N}{\xi_N r_N} F_1^{(0)}(\xi_N) , \qquad (7.3a)$$

$$F_1^{\text{No-TMC}}(x_N) = F_1^{(0)}(x_N) ,$$
 (7.3b)

where $r_N = \sqrt{1 + 4\varepsilon^2}$. There are two differences between these expressions: i) for "No-TMC" the prefactor is set to unity, and ii) the argument of $F_i^{(0)}$ is set to x instead of ξ . Given that $\xi = 2x/(1+r) \simeq x(1-x^2M^2/Q^2)$, the expression for $F_1^{\text{No-TMC}}$ is neglecting the (M/Q) mass effects.

We can separately explore the impact of these two components and determine which is dominant. For F_1 the relative factor $x/(\xi r)$ is precisely the inverse of what is displayed in Fig. 6.1. Thus, we observe that for Q = 1.5 GeV and x = 0.9, this factor yields a shift of ~20%. Comparing with Fig. 7.2, we see this is clearly not sufficient to explain the observed large difference between the "No-TMC" and "Leading-TMC" results. While F_2 and F_3 have slightly different prefactors, we obtain the same conclusion that these prefactors do not generate the sizeable differences observed between the "No-TMC" and the other results.

Thus, the dominant variation of $F_i^{\text{No-TMC}}$ and $F_i^{\text{TMC-Leading}}$ must be due to the different arguments: x vs. ξ . This correspondence of x and ξ is displayed in Fig. 2.1 where we observe that $x \sim \xi$ for

Symbol	A	Z	Symbol	A	Z	Symbol	A		Symbol	A	
Н	1	1	Be	9	4	Ca	40	20	Xe	131	54
D	2	1	С	12	6	Fe	56	26	W	184	74
³ He	3	2	Ν	14	7	Cu _{iso}	64	32	Au	197	79
He	4	2	Ne	20	10	Kr _{iso}	84	42	Au _{iso}	197	98.5
Li	6	3	Al	27	13	Agiso	108	54	Pb_{iso}	207	103.5
Li	7	3	Ar	40	18	$\mathrm{Sn}_{\mathrm{iso}}$	119	59.5	Pb	208	82

Table 7.1: List of nuclear PDFs considered in this work. We use the nCTEQ15 nPDF set [17]. The nuclei indicated with "iso" subscript are isoscalar corrected; thus, the Z value can be half-integer. We will also show results with a neutron PDF computed using isospin symmetry.

small x and large Q, but for large x and small Q we find $\xi < x$. Because the PDFs are steeply falling at large x (where x and ξ have the largest deviation), it is the difference of ξ vs. x in the arguments on the RHS of Eq. (7.3) that is driving the variation observed in Fig. 7.2.

While we have detailed the comparison of F_1 in Fig. 7.2(a), the same rationale applies to the other structure functions displayed.

In contrast to the sizeable differences between the "No-TMC" result and the other three curves for the lower Q value (1.5 GeV), the "No-TMC" result is roughly comparable to the others for Q = 10 GeV. In total, these comparisons affirm that TMCs are negligible for small-x or for intermediate to large Q.

 F_i Structure Function Ratios: (Fig. 7.1 Lower Panels) To highlight the importance of including the TMCs, we now examine the lower panels of Fig. 7.1 which display the ratio of F_i^{TMC} to $F_i^{\text{No-TMC}}$. The ratio plots accentuate subtle features that are not evident on the log-scale plots of the upper panels.

If we first focus on the Q = 1.5 GeV results, we observe the deviation between F_i^{TMC} and $F_i^{\text{No-TMC}}$ differs markedly for the separate $F_{1,2,3}$ structure functions. These differences arise from: i) the different prefactors, see e.g. Eq. (4.15), ii) the different argument of $F_i^{(0)}$ (as noted previously), and also iii) the sub-leading $\{h_i, g_i\}$ contributions.

It is the combination of these components that cause the variation among the Fig. 7.1 sub-figures. For example, we see the F_1 curves deviate at smaller x values as compared with the F_2 and F_3 results. Additionally, the F_2 and F_3 ratios dip below unity for intermediate x values ($x \sim 0.3 - 0.5$) while the F_1 ratio is above unity.

7.2. Nuclear Structure Functions with TMCs

Having examined the proton structure functions in Fig. 7.1, we now consider the corresponding nuclear structure functions of Fig. 7.2. The list of nuclear isotopes that we consider are summarized in Tab. 7.1.

Full TMCs for Nuclei

In Fig. 7.2 we plot the ratio of the full "TMC" structure functions F_i^{TMC} to the "No-TMC" structure functions $F_i^{\text{No-TMC}}$ for charged current (W^-) and neutral current (γ, Z) exchange for the nuclei of Table 7.1. This ratio highlights the impact of the TMCs.

We observe the general behaviour of the nuclear ratios displayed in Fig. 7.2 are very similar to the corresponding proton results in the lower panels of Fig. 7.1. Because the nuclear and proton results



Figure 7.1: Upper panels: Proton structure functions for charged current W^- (a) F_1 , (b) F_2 , and (c) F_3 and (d) neutral current $\gamma/Z F_2$ as a function of momentum fraction x_N . We display results for the full "TMC" (solid blue), "Leading-TMC" (dashed green), "No-TMC" (dot-dash orange, and "ACOT-TMC" (dotted red) at Q = 1.5 GeV and 10 GeV. Lower panels: Ratio of F_i with full "TMC" to "No-TMC" at Q = 1.5 GeV and 10 GeV. Insets: Ratio of F_i with "Leading-TMC" to "ACOT-TMC" at Q = 1.5 GeV.

are so similar, we only plot a single Q value (1.5 GeV) in Fig. 7.2 as other Q values (e.g., Q = 10 GeV) will be similar to the proton results. We will study the detailed Q dependence in the following.

The results of Fig. 7.2 clearly demonstrate that the TMCs for the nuclei are effectively independent of A. We attribute this curious finding to the re-scaling property $(x_A M_A = x_N M_N)$ as shown in Sec. 4. This re-scaling allows us to identically rewrite the TMCs for structure functions in Eq. (3.23), which are functions of nuclear-level quantities x_A and M_A , in terms of nucleon-level quantities x_N and M_N . As a result, the variable nucleon content of nuclei, which is the defining characteristic of nuclei, is



Figure 7.2: Ratio of nuclear structure functions F_i for the full "TMC" over "No-TMC" vs. x_N at Q = 1.5 GeV. We display ratios for charged current W^- (a) F_1 , (b) F_2 , and (c) F_3 and (d) neutral current $\gamma/Z F_2$ for the nuclei listed in Table 7.1. The proton is indicated with the black dashed line, and the neutron with the blue dotted line. The narrow band of colored lines shows the various nuclear results.

averaged out, resulting in near universal behavior.

To illustrate the emergence of this universality, let us Taylor-expand the nuclear structure function ratio $F_1^{\text{TMC-Leading}}/F_1^{\text{No-TMC}}$ to show this is independent of the nuclear A value up to corrections $\varepsilon^2 = (xM/Q)^2$. We use the expressions of Eq. (7.3) and computing the derivative as $F_i^{(0)}(\xi_N) \approx F_i(x_N) + \delta x_N [dF_i(y)/dy]_{y=x_N}$, where $\delta x_N = (\xi_N - x_N) \approx (x^3 M_p^2/Q^2) \ll 1$, If we apply this approximation to the F_1 ratio, for example, we then obtain:

$$\frac{F_1^{\text{TMC-Leading}}(x_N)}{F_1^{\text{No-TMC}}(x_N)} \sim \left(1 - \varepsilon^2\right) \frac{F_1^{(0)}(\xi_N)}{F_1^{(0)}(x_N)} \sim \left(1 - \varepsilon^2\right) \frac{F_1^{(0)}(x_N) + \mathcal{O}(\varepsilon^2)}{F_1^{(0)}(x_N)} \sim \left(1 - \varepsilon^2\right) + \mathcal{O}\left(\varepsilon^4\right) .$$
(7.4)



Figure 7.3: Upper panels: We display the nuclear structure function ratios of $F_i^{\text{TMC}}/F_i^{\text{TMC}-\text{Leading}}$ for selected CC and NC processes for $Q = \{1.3, 1.5, 2, 3, 4, 6\}$ GeV (from top to bottom) vs. x_N . Lower panels: We display the variation of this ratio compared to the average variation as computed in Eq. (7.6) for Q = 2 GeV. In the top row we show F_2 results for (a) NC γ exchange, (b) CC W^+ exchange, and (c) CC W^- exchange, and the bottom row shows F_3 results for the same processes. In all panels, the proton is indicated with the black dashed line, and the neutron with the blue dotted line. The narrow band of colored lines shows the various nuclear results.

where we have defined $\varepsilon = (xM/Q)$. Remarkably, we find that the ratio $F_1^{\text{TMC-Leading}}(x_N)/F_1^{\text{No-TMC}}(x_N)$ is independent of A up to ε^2 . Recall the full TMC ratio $F_1^{\text{TMC}}(x_N)/F_1^{\text{No-TMC}}(x_N)$ differs only by the additional $\{h_i, g_i\}$ terms which are suppressed by powers of $(M/Q)^2$. This implies that nuclear TMCs themselves, when defined in terms of averaged quantities, are essentially universal.

We now explore this approximate universal property of the TMCs, and determine how we can exploit this property to simplify certain nuclear TMC calculations.

Comparison of Full vs. Leading-TMCs for Nuclei

In Fig. 7.1 we observed that the "Leading-TMC" yielded a reasonable approximation to the full "TMC" result, even at low Q and large x values. Computationally, the "Leading-TMC" result is much simpler to compute as the full "TMC" result include contributions from the $\{h_i, g_i\}$ terms, each of which includes an integral. Additionally, if we can take advantage of the approximate A independence observed in the previous section, this may allow us to greatly simplify the calculation of TMCs for the many different nuclei present in a typical nPDF fit.

Magnitude of Ratios: (Fig. 7.3 Upper panels) We start by comparing the ratio of the "Leading-TMC" to the full "TMC" result as shown in the upper panels of Fig. 7.3. The results are shown for selected CC and NC processes with representative values $Q = \{1.3, 1.5, 2, 3, 4, 6\}$ GeV.

We display results for all the nuclei of Table 7.1 In these plots, the individual nuclear A values are not labeled, but the ratios coalesce into clearly defined bands for each Q value. This coalescence holds for all permutations of structure functions and boson exchanges. The different nuclei are shown

as colored solid lines in the figure. We also include results for the proton (dashed black) and neutron (dotted blue).

Focusing first on the bands of the top panels, we see that for F_2^A the differences between the full TMCs and Leading-TMCs are as large as $\mathcal{O}(10\% - 15\%)$ at $x \sim 0.5$ for Q = 1.3 GeV, and $\mathcal{O}(4\% - 5\%)$ for F_3^A . For larger Q values we expect these differences to be reduced as they are proportional to powers of $(M/Q)^2$. At Q = 2 GeV, which is a typical Q_{cut} value for many global PDF analyses, the ratio for F_2^A is $\leq 5\%$, and the ratio of F_3^A is $\leq 2\%$. For $Q \geq 3 - 4$ GeV, we find that these differences reduces to the sub-percent level, and effectively vanish at Q = 6 GeV.

Nuclear A Dependence of Ratios: Having discussed the magnitude of the $F_i^{\text{TMC}}/F_i^{\text{TMC-Leading}}$ ratios, we now examine the A dependence of these ratios. A distinctive feature of Fig. 7.3 is the coalescence of the results into individual bands, and this suggests that the nuclear A dependence of this ratio is minimal. [The proton and neutron ratios (shown as dashed and dotted black lines, respectively), do not necessarily lie within the bands, and we will discuss these separately.] This apparent universality of the full TMC/Leading-TMC ratios can be traced back to: i) the fact that this ratio has only mild dependence on the underlying PDF, and ii) the fact that for nuclei, in the large x region it is the average u + d (isoscalar) PDF that dominates this result.

One approach to understand these features is to consider the analytically computed upper bound for these ratios. If we assume the structure functions are monotonically decreasing (an entirely reasonable assumption in the large x region), it is possible to obtain the following constraints on these ratios [60]:

$$\frac{F_2^{\text{TMC}}}{F_2^{\text{TMC-leading}}}(x,Q^2) \le 1 + \left(\frac{M}{Q}\right)^2 \frac{6x\xi}{r}(1-\xi) + \left(\frac{M}{Q}\right)^4 \frac{12x^2\xi^2}{r^2}(-\ln\xi - 1 + \xi)$$
(7.5a)

$$\frac{F_3^{\text{TMC}}}{F_3^{\text{TMC-leading}}}(x,Q^2) \le 1 - \left(\frac{M}{Q}\right)^2 \frac{2x\xi}{r} \ln \xi \quad .$$
(7.5b)

Note, these bounds have absolutely no dependence on the PDF. Here, we also explicitly see the powers of $(M/Q)^2$ which drive the ratios to unity for large Q.

In Fig. 7.4, we plot these bounds as a function of x for selected Q values. Comparing the magnitudes of the bounds with the values of Fig. 7.3, we see the bounds are quite conservative. Note that these bounds are valid for any nPDFs in the large x region (where F_i^A is decreasing monotonically), and that is the relevant region of interest for the TMC effects. Consequently, the independence of the bound on the PDF helps us understand the comparable nuclear A independence of the $F_i^{\text{TMC}}/F_i^{\text{TMC-Leading}}$ ratios observed in Fig. 7.3.

Nuclear A Variation of Ratios: The observation that these ratios are relatively insensitive to the nuclear A value suggests a "short cut" which can be used to efficiently implement TMCs into numerical calculations.

More specifically, computing the full TMCs for each nucleus A requires a calculation of the $h_i^A(\xi_N, Q^2)$ and $g_i^A(\xi_N, Q^2)$ terms in Eq. (4.15). Each term requires a separate integration over a nuclear structure function. Clearly, such steps would require significant CPU time were they included inside the fitting loop of a global PDF analysis. If it is true that the size of TMCs to nuclear structure functions are relatively insensitive to the nuclear A value, as our works suggests, then TMCs can be estimated through the application of a "universal" correction factor, $R_i^V(x, Q^2)$. Such a factor can be



Figure 7.4: We show the upper bounds of $F_i^{\text{TMC}}/F_i^{\text{TMC-Leading}}$ for a) F_2 and b) F_3 for selected $Q = \{1.3, 1.5, 2, 3, 4, 6\}$ GeV, (from top to bottom) vs. x_N as computed with Eqs. (7.5a) and (7.5b).

constructed by taking an average TMC summed over the various n_A nuclei:

$$R_i^V(x_N, Q^2) = \frac{1}{n_A} \sum_{k}^{n_A} \frac{F_i^{A_k, \text{TMC}}(x_N, Q^2)}{F_i^{A_k, \text{TMC-Leading}}(x_N, Q^2)} = \frac{1}{n_A} \sum_{k}^{n_A} R_i^{k, V}$$
(7.6)

where "V" labels the type of the exchanged gauge boson (γ, Z, W^{\pm}) , and the sum over k excludes the proton and neutron.

The utility of the "universal" correction factor $R_i^V(x, Q^2)$ is that we can compute this once at the beginning of a fit, and then apply this correction factor inside the fitting loop without the need for recomputing additional integrals over h_i and g_i functions.

To determine the potential precision of such an approximation, we show in the lower panels of Fig. 7.3 a ratio of ratios constructed by taking $(F_i^{\text{TMC}}/F_i^{\text{TMC}-\text{Leading}})$ over R_i^V for Q = 2 GeV, which is a typical Q_{cut} value for global fits. It is remarkable how narrow the band of nuclear PDFs lie in comparison to the universal R_i^V function. For F_2 we see the nuclear bands lie within 0.2% of the universal curve, while for F_3 the bands lie within 0.1%. For larger Q values, the precision is further increased. This is clearly sufficient accuracy for any TMC calculation given the typical nPDF uncertainty in the large x region.

Variation of Proton & Neutron Ratios: In Fig. 7.3 we observed that all the nuclear PDFs were uniformly within a narrow band. In contrast, the proton and neutron results displayed a wider variation lying outside these nuclear bands. We will explain the source of this variation, and understand why the universal correction factor $R_i^V(x, Q^2)$ previously introduced, still works well in the case of nuclei.

Specifically, in Fig. 7.3 we see that the proton ratios lie above the nuclear band for the NC and CC W^- plots, and below for the CC W^+ plots; furthermore, the behavior of the neutron ratios is exactly the opposite of the proton. We can understand these features by identifying the dominant contribution for each process. As $x \to 1$, we neglect the gluon and sea quarks to obtain: $F_2(\gamma/Z) \sim \frac{x}{9}[4u(x)+d(x)]$ and $F_2(W^-) \sim 2xu(x)$, while $F_2(W^+) \sim 2xd(x)$. Recall that for a proton PDF, the d/u ratio in the large x region can typically be as small as ~ 0.1; that is, the up quark is an order of magnitude larger than the down quark.

We now understand the pattern of Fig. 7.3. The structure function for the NC and CC W^- processes are driven by the up quark, while the CC W^+ is driven by the down quark. Thus, the proton



Figure 7.5: The nuclear structure function ratio $F_2^{\text{TMC}}/F_2^{\text{TMC-Leading}}$ as function of x_N for $Q = \{1.3, 1.5, 2, 3, 4, 6\}$ GeV for (a) the proton, (b) deuteron, and (c) ³He. For each nucleus, we overlay the result for NC exchange (solid green), CC W^+ exchange (dot orange), and CC W^- exchange (dot-dashed blue). Note that for the deuteron in Fig. (b), the curves for the three CC/NC processes coincide.

results are above the others for NC and CC W^- but below for CC W^+ in plots of Fig. 7.3. Since we obtain the neutron PDFs by isospin symmetry $(u \leftrightarrow d)$, so we also understand why the neutron results are the opposite of the proton.

This exercise also demonstrates why the results for the nuclear PDFs lie together in a uniform band. Most of the nuclei are closer to the isoscalar state $(A \sim 2Z)$ than either the proton (Z = A) or neutron state (Z = 0). For an isoscalar nuclei we have u = d, and it is effectively the average of the up and down PDFs (u + d)/2 which provides the dominant contributions to Fig. 7.3. Consequently, we find that the nuclear band is effectively an average of the proton and neutron result, and thus lies between the two in Figs. 7.3. The one slight variation of this pattern is the parity-violating F_3 NC structure function; this effect must come from the Z boson contribution because the parity-conserving photon contribution vanishes.

Approximate Nuclear A **Independence:** We illustrate this further in Fig. 7.5 which shows the ratios for the NC and CC processes for the a) proton, b) deuteron, and c) ³He. We choose these examples because the deuteron is an isoscalar, and ³He is one of the most "non-isoscalar" nuclei (with the exception of hydrogen/proton).

The pattern in Fig. 7.5 is reminiscent of Fig. 7.3. In Fig. 7.5(a) for the proton, we see the (u-quark dominated) NC and CC W^- processes lie above the (d-quark dominated) CC W^+ process. In Fig. 7.5(b) for the isoscalar deuteron, we see all the curves coincide since u = d for this nuclei. Finally, in Fig. 7.5(c) for the ³He, we see the NC and CC W^- processes lie slightly above the CC W^+ process, but the difference is not as dramatic as for the proton case. Note in these cases, the u : d ratio is 2 : 1 in the proton, 1 : 1 in the deuteron, 5 : 4 in ³He. For any heavier nuclei, the u : d ratio will be comparatively closer to unity, and the results of the individual NC and CC processes will tend to coincide.

The outcome of these findings is that given the typical uncertainty of nuclear PDFs in the large x region, applying an average, A-independent set of TMCs yields precision which is sufficient for a wide variety of real-world calculations. We shall exploit this observation in Sec. 7.3 and derive an approximate parameterization for these TMCs which can be applied to the general nuclear case.

Setting aside the proton and neutron cases, at Q = 2 GeV we find the deviation of individual nuclei from the average to be less than ~ 0.2% for F_2 , and less than ~ 0.1% for F_3 . And of course, these



Figure 7.6: To highlight the effect of the sub-leading terms $\{h_i, g_i\}$, we display the ratios $F_i^{\text{TMC}+h_2}/F_i^{\text{TMC}-\text{Leading}}$ (dotted red) and $F_i^{\text{TMC}}/F_i^{\text{TMC}-\text{Leading}}$ (solid blue), $i = \{1, 2\}$, for selected $Q = \{1.3, 1.5, 2.3, 4, 6\}$ GeV (from top to bottom) as a function of x_N . We show $F_{1,2}$ for neutral current (NC) and charged current (W^+) processes using the nCTEQ15 PDF for ¹²C. Note F_2^{TMC} contains both $\{h_i, g_i\}$ contributions, $F_2^{\text{TMC}+h_2}$ contains only $\{h_i\}$, and $F_2^{\text{TMC}-\text{Leading}}$ contains no sub-leading contributions.

differences decrease for larger Q.

Sub-leading TMCs for Nuclei

As a final study, we want to demonstrate the individual size of the sub-leading h_2^A and g_2^A components of the full TMCs in Eq. (4.15). In Fig. 7.6 we present the results for NC (γ/Z) and CC (W^{\pm}) processes, and we display a pair of ratios $F_2^{\text{TMC+h2}}/F_2^{\text{TMC-Leading}}$ and $F_2^{\text{TMC-Leading}}$, for each Q value.

Recall that $F_2^{\text{TMC-Leading}}$ neglects the sub-leading h_2^A and g_2^A contributions, where h_2^A is suppressed by $(M/Q)^2$, and g_2^A is suppressed by $(M/Q)^4$. The $F_2^{\text{TMC+h2}}$ expression includes the h_2^A contribution, but neglects g_2^A . Finally, the full F_2^{TMC} expression includes both h_2^A and g_2^A ; this could equivalently be called $F_2^{\text{TMC+h2+g2}}$.

Since the g_2^A term is suppressed by $(M/Q)^4$, we expect this will only contribute for very large x at low Q^2 values. Therefore, we expect $F_2^{\text{TMC+h2}}$ and F_2^{TMC} to coincide throughout most of the parameter space. This expectation is validated in Fig. 7.6 where we do see that $F_2^{\text{TMC+h2}}$ and F_2^{TMC}



Figure 7.7: We present the nuclear structure function ratio $F_i^{\text{TMC}}/F_i^{\text{TMC}-\text{Leading}}$ averaged over the nuclei of Table 7.1 as defined in Eq. (7.6). The ratios are shown as a function of x_N for $Q = \{1.3, 1.5, 2, 3, 4, 6\}$ GeV (from top to bottom) using the nCTEQ15 nPDFs. We compare the exact results (green solid) of Eq. (4.15) to the approximate parameterizations (red dashed) based on Eqs. (7.8) with parameters given in Table 7.2. For Q > 2 GeV, we observe the parameterization can match the exact results to better than 0.2%.

differ only for the lowest few Q values in the large x region. For example, at Q = 2 GeV the differences are barely discernible in the figure, and the effect is even smaller as Q increases.

Thus, h_2^A provides the dominant contribution throughout the kinematic range, but g_2^A does contribute for low Q values. The pattern for the other structure functions $F_{1,3}^A$ is similar.

7.3. Parameterizing $F_i^{\text{TMC}}/F_i^{\text{Leading-TMC}}$

In Sec. 7.2 we demonstrated that the full TMC/Leading-TMC ratios were effectively insensitive to the nuclear A value. In this section, we provide a parameterization for these ratios which allows us to efficiently compute the full TMCs given the Leading-TMC structure functions.

Starting from Eqs. (4.15), we divide $F_i^{\text{TMC}}(x)$ by the leading term $F_i^{\text{Leading-TMC}}(x)$:

$$\frac{F_1^{\text{TMC}}(x)}{F_1^{\text{Leading-TMC}}(x)} = 1 + \frac{M^2}{Q^2} \frac{x\xi}{r} \frac{h_2(\xi)}{F_1^{(0)}(\xi)} + \frac{M^4}{Q^4} \frac{2x^2\xi}{r^2} \frac{g_2(\xi)}{F_1^{(0)}(\xi)} , \qquad (7.7a)$$

$$\frac{F_2^{\text{TMC}}(x)}{F_2^{\text{Leading-TMC}}(x)} = 1 + \frac{M^2}{Q^2} \frac{6x\xi^2}{r} \frac{h_2(\xi)}{F_2^{(0)}(\xi)} + \frac{M^4}{Q^4} \frac{12x^2\xi^2}{r^2} \frac{g_2(\xi)}{F_2^{(0)}(\xi)} , \qquad (7.7b)$$

$$\frac{F_3^{\text{TMC}}(x)}{F_3^{\text{Leading-TMC}}(x)} = 1 + \frac{M^2}{Q^2} \frac{2\xi x}{r} \frac{h_3(\xi)}{F_3^{(0)}(\xi)} \quad .$$
(7.7c)

The goal of this section is to parameterize these ratios.

As explained in detail in Appendix B, if we assume the structure functions vanish at x = 1and expand them in a Taylor series, we can obtain simple expressions for the ratios in terms of the parameter function $\gamma_i(Q)$, which characterizes the first derivative of the structure function. This allows us to obtain the approximate expressions for the ratios $h_i/F_i^{(0)}$ and $g_2/F_i^{(0)}$:

$$\frac{h_2(\xi)}{F_2^{(0)}(\xi)} = \frac{1-\xi}{\xi} + \gamma_2(Q) \frac{1-\xi}{\xi^2} \sum_{j=1}^{j_{\text{max}}} \frac{(-1)^j}{j!(j+1)} \,_2F_1\left(2,j+1,j+2,1-\frac{1}{\xi}\right) \,, \tag{7.8a}$$

$$\frac{h_3(\xi)}{F_3^{(0)}(\xi)} = -\ln(\xi) + \gamma_3(Q) \frac{1-\xi}{\xi} \sum_{j=1}^{j_{\text{max}}} \frac{(-1)^j}{j!(j+1)} \,_2F_1\left(1,j+1,j+2,1-\frac{1}{\xi}\right) \,, \tag{7.8b}$$

$$\frac{h_2(\xi)}{F_1^{(0)}(\xi)} = 2\xi r^2 \left[\frac{1-\xi}{\xi} + \gamma_1(Q) \frac{1-\xi}{\xi^2} \sum_{j=1}^{j_{\text{max}}} \frac{(-1)^j}{j!(j+1)} \, _2F_1\left(2,j+1,j+2,1-\frac{1}{\xi}\right) \right] \,, \tag{7.8c}$$

$$\frac{g_2(\xi)}{F_2^{(0)}(\xi)} = -\ln(\xi) - (1-\xi) + \gamma_2(Q) \frac{(1-\xi)^2}{\xi^2} \sum_{j=1}^{j_{\max}} \frac{(-1)^j}{j!(j+2)} \,_2F_1\left(2, j+2, j+3, 1-\frac{1}{\xi}\right) \,, \tag{7.8d}$$

$$\frac{g_2(\xi)}{F_1^{(0)}(\xi)} = 2\xi r^2 \left[-\ln(\xi) - (1-\xi) + \gamma_1(Q) \frac{(1-\xi)^2}{\xi^2} \sum_{j=1}^{j_{\text{max}}} \frac{(-1)^j}{j!(j+2)} \, _2F_1\left(2, j+2, j+3, 1-\frac{1}{\xi}\right) \right] \,, \quad (7.8e)$$

where ${}_{2}F_{1}(a, b, c, z)$ is a hypergeometric function. Although the summation over the index j can, in principle, go to infinity, we truncate the series at $j = j_{\text{max}}$. The 1/(j!) prefactor of the hypergeometric function helps ensure the series converges quickly, and we find $j_{\text{max}} = 4$ yields results that are $\leq 1\%$ accuracy.

The function $\gamma_a(Q)$ is given by

$$\gamma_a(Q) = \lambda_a \ln(Q)^{\delta_a} \quad . \tag{7.9}$$

The values of $\{\lambda_a, \delta_a\}$ are obtained by fitting the parameterization to the exact results for each F_i^{TMC} structure function. Note that the values of $\{\lambda_a, \delta_a\}$ are independent of the type of the exchanged bosons; they have only a mild dependence on the specific structure function F_i^A as displayed in Table 7.2.

Additionally, while we have computed these results assuming an NLO DGLAP evolution, these results should be similar for an NNLO evolution. This is because: i) we are working at relatively low Q values where the effects of the DGLAP evolution from Q_0 to Q will be minimal, and ii) since this quantity is a ratio of structure functions, the modification of the ratio is expected to be less than the

	F_1^{TMC}	$F_1^{TMC}/F_1^{leading}$		$/F_2^{leading}$	$F_3^{TMC}/F_3^{leading}$		
nPDFs	λ_1	δ_1	λ_2	δ_2	λ_3	δ_3	
nCTEQ15	2.275	-0.014	2.144	0.100	2.183	0.029	
EPPS16	2.226	-0.026	2.086	0.095	2.142	0.022	
nNNPDF2.0	2.226	-0.025	2.086	0.103	2.197	0.023	
TUJU19	2.290	-0.0170	2.167	0.099	2.200	0.031	

Table 7.2: We approximate the $\{h_i(\xi), g_i(\xi)\}$ contributions to F_i^{TMC} using $F_i^{\text{TMC-Leading}}$ together with a 2-parameter function $\gamma_a(Q)$. Here, $\gamma_a(Q)$ implicitly depends on the parameters $\{\lambda_a, \delta_a\}$ which are fitted for each structure function $\{F_{1,2,3}\}$ and for selected PDFs displayed in the table. The TMC corrections are accurate to within ~ 0.3% for Q > 1.3 GeV. The $\{\lambda_i, \delta_i\}$ parameters are independent of the exchanged boson (γ, Z, W^{\pm}) , and are relatively insensitive to the specific underlying nPDF. Once the $\{\lambda_i, \delta_i\}$ parameters are determined for a particular nPDF, the full TMC corrections can be efficiently computed (without convolution integrations) to high precision as illustrated in Fig. 7.7.

impact on the individual structure functions.

We have also computed the results for a variety of different PDF sets to ensure this parameterization is largely insensitive to the specific PDF. This also is displayed in Table 7.2, where we observe a mild variation of the parameters between PDF sets.

In Figure 7.7, we show the comparison between our parameterization with the exact results obtained using nCTEQ15 nPDFs. We can see that our parameterization works very well to reproduce the exact results. The parameterization has the advantage that it can be computed efficiently whereas the full F_i^{TMC} results require additional convolution integrations for the $\{h_i(\xi), g_i(\xi)\}$ functions. For values of $Q \ge 2$ GeV, which is a typical cut for the global nPDF fits, we see the parameterization describes the full TMC results to better than 0.2%.

Summary: In the previous sections we established several results which will facilitate efficient calculation of TMCs for nuclear structure functions. We briefly summarize the key observations below.

- TMCs can yield large corrections in the double limit of large x and small Q.
- The "Leading-TMC" result provides an excellent approximation to the full "TMC" result as the $h_{2,3}$ contributions are suppressed by additional $(M/Q)^2$, and the g_2 contributions by additional $(M/Q)^4$ powers.
- The nuclear dependence of the "TMC" result is approximately independent of A. This observation suggests we can determine an A-independent "universal" correction factor which can be used to quickly calculate the nuclear structure function TMCs for any nuclei. As illustrated in the bottom panels of Fig. 7.3, this "universal" approximation holds to $\leq 0.1\%$ for $Q \geq 2$ GeV.
- As demonstrated in Fig. 7.7, we can parameterize this "universal" correction factor with Eq. (7.8) into a 2-parameter form that can be easily incorporated in a efficient nuclear PDF fitting program. This parameterization matches the exact results to better than 0.2% for $Q \ge 2$ GeV.

Given the "Leading-TMC" result, the above allows us to efficiently approximate the full "TMC" result for any nuclear structure function by applying "universal" correction factor. This avoids the computation of the numerical integrals contained in the $\{h_2, h_3, g_2\}$ functions.

7.4. Reduced Cross Sections with TMCs for Nuclei

When extracting structure functions from DIS data or comparing theoretical expectations for DIS to data, one often works with the "reduced cross section" $\sigma_{\text{Red.}}$. These quantities are derived from double differential cross sections scaled by kinematic factors, and provide a direct connection between DIS kinematics and structure functions.

The general cross section for neutral and charged current DIS can be expressed as: [75]

$$\frac{d^2\sigma^i}{dx\,dy} = x(s-M^2)\,\frac{d^2\sigma^i}{dxdQ^2} = \frac{4\pi\alpha^2}{xyQ^2}\,\eta^i\left\{\left(1-y-\frac{x^2y^2M^2}{Q^2}\right)F_2^i + \left(y^2x\right)F_1^i \mp \left(y-\frac{y^2}{2}\right)xF_3^i\right\},\tag{7.10}$$

where i=NC, CC corresponds to neutral-current or charged-current processes. Additionally, $\eta^{NC} = 1$ and $\eta^{CC} = 4\eta_w$, with $\eta_w = \frac{1}{2} \left[\frac{G_F M_W^2}{4\pi \alpha} \frac{Q^2}{Q^2 + M_W^2} \right]^2$. In the last term of Eq. (7.10), the "–" sign is taken for an incoming anti-lepton $\{e^+, \bar{\nu}\}$ and the "+" sign for an incoming lepton $\{e^-, \nu\}$.

Neutral Current (NC) Cross Section: We find it convenient to define the reduced cross section by dividing out the leading factors.

$$\frac{d^2 \sigma^{\rm NC}}{dx \, dy} = x(s - M^2) \frac{d^2 \sigma^{\rm NC}}{dx dQ^2} = \frac{4\pi \alpha^2}{xyQ^2} \left[\frac{Y_+}{2} \sigma^{\rm NC}_{\rm Red.} \right] , \qquad (7.11)$$

$$\sigma_{\text{Red.}}^{NC} = \left(1 + \frac{2y^2 \varepsilon^2}{Y_+}\right) F_2^{\text{NC}} \mp \frac{Y_-}{Y_+} x F_3^{\text{NC}} - \frac{y^2}{Y_+} F_L^{\text{NC}} , \qquad (7.12)$$

with $F_L = r^2 F_2 - 2xF_1$, $r = \sqrt{1 + 4\varepsilon^2}$, $\varepsilon = (xM/Q)$ and $Y_{\pm} = 1 \pm (1 - y)^2$; the r^2 term in F_L is essential to obtain the correct sign for the ε^2 term of Eq. (7.12). Note that at LO ($F_L \sim 0$) for parity-conserving photon exchange ($F_3 = 0$), so that we have σ_{Red}^{NC} simply reduces to F_2^{NC} .

Charged Current (CC) Cross Section: For the charged current process, we have:

$$\frac{d^2 \sigma^{\rm CC}}{dx \, dy} = x(s - M^2) \frac{d^2 \sigma^{\rm CC}}{dx dQ^2} = \frac{G_F^2}{2\pi x} \frac{Q^2}{y} \left(\frac{M_W^2}{Q^2 + M_W^2}\right)^2 \left[\sigma_{\rm Red.}^{CC}\right], \tag{7.13}$$

$$\sigma_{\text{Red.}}^{CC} = \frac{Y_{+}}{2} \left(1 + \frac{2y^{2}\varepsilon^{2}}{Y_{+}} \right) F_{2}^{\text{CC}} \mp \frac{Y_{-}}{2} x F_{3}^{\text{CC}} - \frac{y^{2}}{2} F_{L}^{\text{CC}}$$
(7.14)

Here we have followed the HERA convention [125] where the CC differs from NC by a factor of $Y_+/2$.

For both the NC and CC cases, the factor of $2y^2\varepsilon^2/Y_+$ with $\varepsilon = (xM/Q)$ multiplying F_2 arises directly from the DIS kinematics; however, this is often neglected as it is small for typical HERA kinematics.

TMCs for Nuclear Reduced Cross Sections

As described in the preceding subsections, TMCs can have a large numerical impact on nuclear structure functions in the low Q^2 and large x double limit. To illustrate the TMC impact, in Fig. 7.8 we present results for experimental scenarios including HERA, JLab, EIC, and LBNF. The upper panels of Fig. 7.8 display the reduced cross sections, and the lower panels show the ratio of the full



Figure 7.8: We present reduced cross sections for the following scenarios: (a) HERA inspired e^+p neutral current scattering with $E_{CM} = 318$ GeV for Q = 1.3 GeV; (b) JLab inspired e^-A charged current scattering with $E_{beam} = 24$ GeV for Q = 1.3 GeV; (b) EIC inspired e^-A neutral current scattering with $E_{CM} = 100$ GeV for Q = 1.3 GeV; (d) LBNF inspired νA charged current scattering with $E_{CM} = 10$ GeV for Q = 1.3 GeV; (d) LBNF inspired νA charged current scattering with $E_{CM} = 10$ GeV for Q = 1.3 GeV; (d) LBNF inspired νA charged current scattering with $E_{CM} = 10$ GeV for Q = 1.3 GeV. Upper panels: The upper panels display the full "TMC" results. Lower panels: The lower panels display the ratio of the full "TMC" to "No-TMC". The proton is indicated with the black dashed line, and the neutron with the blue dotted line. The narrow band of colored lines shows the various nuclear results.

"TMC" to the "No-TMC" result. As before, all the nuclear isotopes of Table 7.1 are within the colored band, and the proton and neutron are displayed separately.

In Fig. 7.8-(a) we show the HERA-inspired e^+p neutral current σ_{Red}^{NC} at $\sqrt{s} = 318$ GeV at Q = 1.3 GeV for the full "TMC" results. Although HERA ran proton beams, we also show the nuclear results which may be useful for a future LHeC/FCCeh facility. The TMCs yield a slight suppression at intermediate x values (~ 0.4), and an enhancement at larger x values (≥ 0.6) which can be significant. The behavior of σ_{Red}^{NC} strongly resembles that of F_2 in Fig. 7.1; this is expected as given our choices of Q and E_{CM} we find $y \ll 1$ and $Y_- \sim 0$, so we see from Eq. (7.12) that the F_2 contribution dominates. For the higher Q values (not shown) TMCs are reduced as for F_2 in Fig. 7.1, and are generally negligible except in the very large x region.

In Fig. 7.8-(b) we show a JLab-inspired e^-A neutral-current σ_{Red}^{NC} with $E_{\text{Beam}} = 24$ GeV at Q = 1.3 GeV. The character of the curves for $E_{\text{Beam}} = 12$ GeV (not shown) is quite similar to the displayed results. The results here are qualitatively similar to the F_2 result of Fig. 7.1 and the HERA case above with a reduction for intermediate x (~ 0.4), and an enhancement at larger x.

In Fig. 7.8-(c) we show an EIC-inspired e^-A neutral-current σ_{Red}^{NC} with $\sqrt{s} = 100$ GeV at Q = 1.3 GeV. As with the above HERA and JLab cases, these results largely follow the F_2 structure function.

In Fig. 7.8-(d) we show an LBNF-inspired $\nu - A$ charged-current scattering with $E_{\text{Beam}} = 10$ GeV at Q = 1.3 GeV. In the low x region, we see the various nuclear curves separate due to the differing nuclear mass values.

From this exercise, one can see that for large \sqrt{s} , we generally have small y and σ_{Red} closely follows F_2 . For smaller \sqrt{s} values, we have larger y and the F_3 and F_L structure functions can now contribute.

8. Conclusions

To exploit the current and future nuclear measurements and to obtain the desired accuracy of theory calculations, it is essential to include TMCs to structure functions in analyses. The present study provides both pedagogical material and a practical reference for computing and investigating TMCs for nuclear processes such as at the upcoming Electron Ion Collider (EIC) and Deep Underground Neutrino Experiment (DUNE). We briefly outline the key components and results in the following:

- We extend the OPE formalism to the case of interacting nuclei and explicitly provide the correspondence between TMC expressions for a nucleon and a nucleus, given in Eq. (3.24). To derive these relations (see Sec. 3.2 and Appendix A), we use only fundamental symmetry principles of the nuclei, and make no assumptions about the parton or other models.
- We specifically investigate analytically and numerically the magnitude of TMCs and show it is the quantity $(x_A M_A/Q) = (x_N M_N/Q)$ that enters the master TMC formula, Eq. (4.15). Given that the dominant support of nPDFs is in the region $x_N \in [0, 1]$, TMCs are governed by powers of (M_N/Q) and not (M_A/Q) .
- In Sec. 4.2, we derive the master TMC expression in terms of the rescaled structure functions. We also investigate the power-suppressed terms proportional to h_i and g_i , and determine their rescaling properties.
- In terms of the rescaled structure functions, we define the rescaled nPDFs, which are the ones commonly studied and presented in comparisons between different nuclear data. In Sec. 5.1, we also discuss the corresponding set of DGLAP equations and associated sum rules, including isospin symmetry and the complication of separately extraction u(x) and d(x) nPDFs.
- In particular, we establish the relations between the kinematic variables including the nuclear scaling variables $\{x_A, \xi_A\}$ and the nucleon scaling variables $\{x_N, \xi_N\}$. Additionally, we indicate in Sec. 5.4 some subtle complications we encounter such as defining an appropriate W^2 cut for the nucleon case.
- We obtain a complete correspondence between the parton model and OPE structure function results identifying the source of the differences. Additionally, in Sec. 6 we review the ACOT helicity approach with light-cone kinematics, and compare with the direct OPE results.
- In Sec. 7, we provide variety of numerical comparisons to demonstrate the size of the TMCs in various kinematic regions. These comparisons are shown for both structure functions, and the reduced cross section for some realistic machine kinematics.

• We also present numerically the dependence of the TMCs on the nuclear A value and the exchanged vector boson $\{\gamma, Z, W^{\pm}\}$. We find the relative magnitude of the TMCs is weakly dependent on the exchanged boson, and very weakly dependent on the nuclear A. This latter observation allow us to derive an approximate (and computationally expedient) parameterization for the nuclear TMCs suitable for including inside a global fit function.

Just as Ref. [60] provided the details for the proton case, we hope this report provides foundational and reference material which can facilitate incisive investigations of nuclear phenomena.

Acknowledgments

The authors would like to thank Eric Christy, John, C. Collins, Stefan Krezer, and Mary Hall Reno for many useful comments and discussions.

A.K. and R.R. acknowledge the support of Narodowe Centrum Nauki under Grant No. 2019/34/E /ST2/00186. R.R also acknowledges the support of the Polska Akademia Nauk (grant agreement PAN.BFD.S.BDN. 613. 022. 2021 - PASIFIC 1, POPSICLE). This work has received funding from the European Union's Horizon 2020 research and innovation program under the Skłodowska-Curie grant agreement No. 847639 and as part of the Skłodowska-Curie Innovative Training Network MCnetITN3 (grant agreement no. 722104), and from the Polish Ministry of Education and Science. The work at WWU Münster was funded by the DFG through the Research Training Group 2149 "Strong and Weak Interactions - from Hadrons to Dark Matter" and the SFB 1225 "Isoquant," project-id 273811115. M.K. thanks the School of Physics at the University of New South Wales in Sydney, Australia for its hospitality and financial support through the Gordon Godfrey visitors program. The work of A.A. was supported by the U.S. Department of Energy under Grant No. DE-SC0008791. The work of T.J.H. was supported by the U.S. Department of Energy under contract DE-AC02-06CH11357. The research of P. D. was funded as a part of the Center of Excellence in Quark Matter of the Academy of Finland (project 346326). The work of C.K. was supported by the U.S. Department of Energy contract DE-AC05-06OR23177, under which Jefferson Science Associates LLC manages and operates Jefferson Lab. J.G.M. has been supported by Fermi Research Alliance, LLC under Contract No. DE-AC02-07CH11359 with the U.S. Department of Energy, Office of Science, Office of High Energy Physics. The work of F.O. was supported by the U.S. Department of Energy under Grant No. DE-SC0010129. The work of I.S. was supported by the French CNRS via the IN2P3 project GLUE@NLO.

Appendix A. Nuclear structure functions with TMCs in the OPE

In this appendix, we build structure functions for massive, unpolarized nuclear targets following a modified prescription of Ref. [49, 50]. The purpose of this is two fold. First is to establish that the TMC formulae used for free nucleons are applicable to nuclei. Second is to provide a explicit derivation of the TMC master formulae with details beyond what is generally available in the literature. It is well-documented that constructing TMCs to structure functions is a tedious endeavor, with many intermediate steps omitted from the literature. Some notable exceptions exist [49, 50, 54, 60, 81, 126]. For example: Refs. [49, 50] give a largely complete treatment of F_2 but implicitly employ derivative, integral, and summation identities. Some of these intermediate steps are more completely documented in Refs. [81, 126], but other steps may not be fully justified under their assumptions [85]. Similarly, while the structure functions W_4 and W_5 (or equivalently F_4 and F_5) are documented in Refs. [54, 60], neither works out the many permutations of contractions with the momentum factor $\Pi^{\mu_1...\mu_{2k}}$ from Eq. (3.19). Keeping track of these permutations requires care, as demonstrated for F_1 and F_2 in Ref. [126]. Notably, Ref. [60] documents some differences and similarities between the above treatments. The treatment in this Appendix builds on the steps of Refs. [49, 50, 54, 63, 64, 81, 85, 126].

We first build a formula for the inclusive DIS cross section in Appendix A.1 based on the notation defined in section 2.1. In Appendix A.2, we derive the matrix element, squared matrix element, and inclusive hadronic tensor in DIS. In Appendix A.3, the OPE is applied to the inclusive hadronic tensor, with the leading-power result organized according to Wilson coefficients. In Appendix A.4, OPE terms are reorganized by Lorentz structures, resulting in structure-function mixing. Structure functions for massless and massive nuclear targets are then constructed in Appendix A.5.

Appendix A.1. Preliminaries and the inclusive DIS cross section formula

To build nuclear structure functions with TMCs in the OPE formalism, we consider the DIS process as defined in Eq. (2.1) of section 2.1 and depicted in Fig. 1.1. Given this configuration, the total cross section for a generic $2 \rightarrow (n+1)$ scattering process is given by the formula

$$\sigma(\ell_1 + A \to \ell_2 + X_n) = \int dP S_{n+1} \frac{d\sigma}{dP S_{n+1}}, \qquad \frac{d\sigma}{dP S_{n+1}} = \frac{1}{\mathcal{F}} \frac{1}{\mathcal{S}_{\ell_1} \mathcal{S}_A} \sum_{\{\lambda\}} \sum_{\text{dof}} |\mathcal{M}|^2, \tag{A.1a}$$

$$\mathcal{F} = 4\sqrt{(p_A \cdot k_1)^2 - p_A^2 k_1^2} \stackrel{k_1^2 = m_1^2 \to 0}{\approx} 2(s - M_A^2).$$
(A.1b)

Here, \mathcal{F} is the Moller flux factor and reduces to $\mathcal{F} \approx 2(s - M_A^2)$ when the mass $k_1^2 = m_1^2$ vanishes. $d\sigma/dPS_{n+1}$ is the totally differential, nucleus-level cross section. S_{ℓ_1} and S_A are the spin multiplicities for ℓ_1 and A in unpolarized scattering. The double summations in the differential cross section run over external helicities $\{\lambda\}$ and other discrete quantum numbers/degrees of freedom (dof). We introduce the subscript n on X_n to make explicitly that it is an n-body state. In addition, \mathcal{M} is the scattering amplitude for the $2 \rightarrow (n+1)$ process. The associated (n+1)-body phase space measure is given by

$$dPS_{n+1}(k_1 + p_A; k_2, p_X) = (2\pi)^4 \delta^4 \left(k_1 + p_A - k_2 - \sum_{i=1}^n p_{X_i} \right) \frac{d^3 k_2}{(2\pi)^3 2k_2^0} \times \prod_{i=1}^n \frac{d^3 p_{X_i}}{(2\pi)^3 2p_{X_i}^0}$$
(A.2)

$$= dPS_n(q + p_A; p_X) \times \frac{d^3k_2}{(2\pi)^3 2k_2^0}$$
(A.3)

$$= \int d^4 z \ e^{i(q+p_A-p_X)\cdot z} \ \prod_{i=1}^n \frac{d^3 p_{Xi}}{(2\pi)^3 2p_{Xi}^0} \ \times \frac{d^3 k_2}{(2\pi)^3 2k_2^0} \ . \tag{A.4}$$

In the above lines, we employed momentum conservation and δ -function identities to rewrite the measure in ways that will be used below. Specifically in the third line, we used

$$(2\pi)^4 \,\delta^4(q-p) = \int d^4z \,e^{i(q-p)\cdot z} , \qquad (A.5)$$

where q, p and z are 4-vectors. The point of these identities is to identify the phase space integral over the hadronic matrix element also as the Fourier transformation over composite operators. The modeling of X_n 's phase space is not precisely correct in many presentations of structure functions in DIS. Often, p_X is treated as a one-body configuration despite being an *n*-body configuration. While this mis-modelling is ultimately a technicality, it does impact the formal definition of $W_{\mu\nu}$; see, for instance, Ref. [127]. We reiterate that X_n here (or X in the main text) is an *n*-body state.

Using the above phase space decomposition, we can write an expression for the DIS cross section that (1) is differential with respect to the kinematics of $\{\ell_2, k_2\}$, and (2) is explicitly inclusive over all hadronic activity. This expression is given by

$$k_2^0 \times \frac{d\sigma}{d^3 k_2} = \frac{1}{(16\pi^2)} \frac{1}{\pi} \frac{1}{\mathcal{F}} \frac{1}{\mathcal{S}_{\ell_1} \mathcal{S}_A} \sum_{n=1}^{\infty} \int d^4 z \ e^{i(q+p_A-p_X)\cdot z} \prod_{i=1}^n \frac{d^3 p_{Xi}}{(2\pi)^3 2 p_{Xi}^0} \sum_{\{\lambda\}} \sum_{\text{dof}} |\mathcal{M}|^2 \ . \tag{A.6}$$

Here, $S_{\ell_1} = 2$ for an unpolarized ℓ_1 , and $S_A = 2$ for an unpolarized, spin-1/2 nuclei A. Additionally, $S_{\ell_1} = 1$ when ℓ_1 is an incoming neutrino or antineutrino because neutrino beams are effectively 100% polarized. More broadly, for unpolarized nuclear targets with spin $s_A > 1/2$, the spin-averaging factor S_A generalizes to $S_A = (2s_A + 1) > 2$.

In Eq. (A.6), we inserted a sum over the multiplicity n of X_n . Conceptually, this is equivalent to the relationship

$$\ell_1 + A \rightarrow \ell_2 + \text{any hadronic activity} = \sum_{n=1}^{\infty} \left[\ell_1 + A \rightarrow \ell_2 + X_n \right].$$
 (A.7)

For example, when n = 2 this sums over the configurations $pp, pn, p\pi, pK, \pi\pi, \ldots$, which are all included in the inclusive sum.

Appendix A.2. The DIS matrix element and inclusive hadronic tensor

The hadronic tensor $W^A_{\mu\nu}$ of Eq. (2.7) is built from the (squared) matrix element \mathcal{M} . For $\ell_1 A \to \ell_2 X$ scattering via an arbitrary electroweak boson V(q), we parameterize this by

$$-i\mathcal{M} = \langle X\ell_2 | J^{\mu}_{\ell_2\ell_1}(0) \cdot \Delta^V_{\mu\sigma}(q) \cdot J^{\sigma}_{XA}(0) | A\ell_1 \rangle$$
(A.8)

$$= \langle \ell_2 | J^{\mu}_{\ell_2 \ell_1}(0) | \ell_1 \rangle \cdot \Delta^V_{\mu \sigma}(q) \cdot \langle X | J^{\sigma}_{XA}(0) | A \rangle, \tag{A.9}$$

where the leptonic part of the matrix element L^{μ} and current $J^{\mu}_{\ell_2\ell_1}$ in terms of 4-component helicity spinors, a generic coupling normalization \tilde{g} , and chiral couplings g_V, g_A , are

$$L^{\mu} \equiv \langle \ell_2 | J^{\mu}_{\ell_2 \ell_1}(0) | \ell_1 \rangle = -i \tilde{g} \, \overline{u}(k_2, \lambda_2) \left[g^{\ell}_V \gamma^{\mu} + g^{\ell}_A \gamma^{\mu} \gamma^5 \right] u(k_1, \lambda_1).$$
(A.10)

We assume arbitrary vector and axial-vector couplings between the $(\ell_2 \ell_1)$ system and V in order to accommodate the possibility of V being a photon or weak boson. The conversion between the different possibilities is given in Tab. A.1. Similarly, we write the hadronic part of the matrix element in terms

Vertex	Coupling strength	g_R^ℓ	g_L^ℓ	g_V^ℓ	g^ℓ_A
$\frac{V - \ell_1 - \ell_2}{\gamma - \ell - \ell}$	${egin{array}{c} {ar g} \\ eQ^\ell \end{array}}$	$(g_V^\ell + g_A^\ell)$ 1	$(g_V^\ell - g_A^\ell) \ 1$	$\frac{(g_R+g_L)}{2}$ 1	$\frac{(g_R - g_L)}{2}$
$\overset{'}{Z} - \ell - \ell$ $W - \ell_1 - \ell_2$	$\frac{g_W}{\cos\theta_W}$ $\frac{g_W}{\sqrt{2}}$	$-Q^{\ell}\sin^2\theta_W$	$(T_3^\ell)_L - Q^\ell \sin^2 \theta_W$ 1	$\frac{\frac{1}{2}(T_3^\ell)_L - Q^\ell \sin^2 \theta_W}{\frac{1}{2}}$	$-\frac{1}{2}(T_3^\ell)_L -\frac{1}{2}$

Table A.1: Electroweak chiral couplings and coupling strength normalizations used for fermions ℓ_1, ℓ_2 with weak isospin charge $(T_3^4)_L = \pm 1/2$ and electric charge Q^ℓ , with normalization $Q^e = -1$.

of an arbitrary hadronic current $J_{XA}^{\sigma}(0)$:

$$H^{A\sigma} \equiv \langle X|J_{XA}^{\sigma}(0)|A\rangle \equiv \overline{u}_X(p_X,\lambda_X)[\dots]^{\sigma}u(p_A,\lambda_A).$$
(A.11)

We assume here and below that J_{XA}^{σ} is always a renormalized object in QCD.

In Eq. (A.9), $\Delta_{\mu\sigma}^{V}(q)$ is the propagator of V. When we assume the ℓ_i are massless, the Dirac equation and $\mathrm{SU}(2)_L$ invariance ensure that the longitudinal modes of $\Delta_{\mu\sigma}^{V}$ do not contribute to the scattering process.¹² More specifically, the longitudinal component of Δ scales as $\Delta_{\mu\sigma}(q)|_{\mathrm{long}} \propto q_{\mu}q_{\sigma}$. Since $k_i u(k_i) = m_i u(k_i) = 0$ and $q_{\mu} = (k_1 - k_2)_{\mu}$, one finds

$$L^{\mu} \cdot \Delta^{V}_{\mu\sigma}|_{\text{long}} \propto q_{\mu} \cdot \overline{u}(k_{2}, \lambda_{2}) \left[g^{\ell}_{V} \gamma^{\mu} + g^{\ell}_{A} \gamma^{\mu} \gamma^{5} \right] u(k_{1}, \lambda_{1})$$
(A.12)

$$= \overline{u}(k_2, \lambda_2) \left[g_V^{\ell}(k_1 - k_2) + g_A^{\ell}(k_1 - k_2)\gamma^5 \right] u(k_1, \lambda_1) = 0.$$
(A.13)

The above implies that only the transverse components of $\Delta^V_{\mu\sigma}$ contributes to DIS for massless leptons. For $V = \gamma, W, Z$ with mass M_V (where $M_{\gamma} = 0$), one can then write

$$-i\mathcal{M} = L^{\mu}(k_2, k_1)\Delta^{V}_{\mu\sigma}(q)H^{A\sigma}(p_X, p_A) = L^{\mu}\frac{-ig_{\mu\sigma}}{q^2 - M_V^2}H^{A\sigma} = \frac{-i}{q^2 - M_V^2}L^{\mu}H^A_{\mu}.$$
 (A.14)

After squaring and summing over external helicity states, one obtains

$$\sum_{\{\lambda\}} |\mathcal{M}|^2 = \frac{1}{(q^2 - M_V^2)^2} \sum_{\{\lambda\}} L^{\mu\nu} \cdot \sum_{\{\lambda\}} H^A_{\mu\nu} .$$
(A.15)

The leptonic tensor $L^{\mu\nu}$ denotes the square of L^{μ} and can be solved exactly using trace technology,

 $^{^{12}}$ We typically neglect the mass of the leptons, except for the tau-lepton.

helicity amplitudes, or other standard technology. It is given by

$$\sum_{\{\lambda\}} L^{\mu\nu} = \sum_{\{\lambda\}} \left(L^{\dagger} \right)^{\mu} L^{\nu} = \sum_{\{\lambda\}} \langle \ell_1 | \left(J^{\dagger}_{\ell_2 \ell_1}(0) \right)^{\mu} | \ell_2 \rangle \langle \ell_2 | J^{\nu}_{\ell_2 \ell_1}(0) | \ell_1 \rangle$$
(A.16)

$$=4\tilde{g}^{2}\left\{(g_{V}^{2}+g_{A}^{2})(k_{1}^{\mu}k_{2}^{\nu}+k_{1}^{\nu}k_{2}^{\mu})-\left[(g_{V}^{2}+g_{A}^{2})(k_{1}\cdot k_{2})-(g_{V}^{2}-g_{A}^{2})m_{1}m_{2}\right]g^{\mu\nu}-2i(g_{V}g_{A})k_{1\alpha}k_{2\beta}\epsilon^{\mu\nu\alpha\beta}\right\}$$
(A.17)

$${}^{m_1,m_2 \to 0} 4\tilde{g}^2 \Big\{ (g_V^2 + g_A^2) (k_1^{\mu} k_2^{\nu} + k_1^{\nu} k_2^{\mu}) - (g_V^2 + g_A^2) (k_1 \cdot k_2) g^{\mu\nu} - 2i (g_V g_A) k_{1\alpha} k_{2\beta} \epsilon^{\mu\nu\alpha\beta} \Big\}$$
(A.18)

For completeness, we provide the expression for $\sum_{\{\lambda\}} L^{\mu\nu}$ with massive leptons. If using this formula, note that Eq. (A.14) must be modified since $k_i u(k_i) = m_i u(k_i) \neq 0$. For the remainder of this work, we keep $m_1, m_2 = 0$. Note that the expression here for the leptonic tensor only sums over initialand final-state polarizations; a symmetry factor of $1/S_{\ell_1} = 1/2$ for unpolarized initial-states has not yet been applied. Following the conventions of Table A.1, for the case of QED, where $g_V = 1$, $g_A = 0$, and $\tilde{g} = e$, we recover the usual expression:

$$\sum_{\{\lambda\}} L^{\mu\nu} \Big|_{\text{QED}} = 4e^2 \Big\{ k_1^{\mu} k_2^{\nu} + k_1^{\nu} k_2^{\mu} - (k_1 \cdot k_2) g^{\mu\nu} \Big\}.$$
(A.19)

For the case of W boson exchange, where $g_V = 1/2$, $g_A = -1/2$, and $\tilde{g} = g_W/\sqrt{2}$, we have

$$\sum_{\{\lambda\}} L^{\mu\nu} \Big|_{W} = g_{W}^{2} \Big\{ k_{1}^{\mu} k_{2}^{\nu} + k_{1}^{\nu} k_{2}^{\mu} - (k_{1} \cdot k_{2}) g^{\mu\nu} + i k_{1\alpha} k_{2\beta} \epsilon^{\mu\nu\alpha\beta} \Big\}.$$
(A.20)

The exclusive hadronic tensor $H^A_{\mu\nu}$ for an *n*-body final state X denotes the square of H_{μ} . For a fixed *n*, it is given by

$$\sum_{\{\lambda\}} H^A_{\mu\nu} = \sum_{\{\lambda\}} \left(H^{A\dagger} \right)_{\mu} H^A_{\nu} = \sum_{\{\lambda\}} \langle A | \left(J^{\dagger}_{XA}(0) \right)_{\mu} | X \rangle \langle X | J_{XA\nu}(0) | A \rangle .$$
(A.21)

From $H^A_{\mu\nu}$ we can build the *inclusive* hadronic tensor $W^A_{\mu\nu}$ that sums over all final states.

Building the inclusive hadronic tensor

In inclusive DIS, one measures the kinematics of ℓ_2 and remains totally inclusive regarding the hadronic system X. (Though in practice, the invariant mass of X is often measured.) The inclusiveness criterion means summing over all possible final-state multiplicities n (where n is same n in Eq. (A.2)), discrete dof, and continuous dof, i.e., momenta. Performing this summation allows us to define the *inclusive*

hadronic tensor as

$$W_{\mu\nu}^{A} \equiv \frac{1}{4\pi} \sum_{n=1}^{\infty} \int dP S_{n}(q+p_{A};p_{X}) \sum_{\text{dof } \{\lambda\}} H_{\mu\nu}^{A}$$
(A.22)

$$= \frac{1}{4\pi} \int d^{4}z \ e^{i(q+p_{A}-p_{X})\cdot z} \sum_{n=1}^{\infty} \int \prod_{i=1}^{n} \frac{d^{3}p_{Xi}}{(2\pi)^{3}2p_{Xi}^{0}} \\ \times \sum_{\text{dof}} \sum_{\{\lambda\}} \langle A|J_{XA\mu}^{\dagger}(0)|X\rangle \langle X|J_{XA\nu}(0)|A\rangle$$
(A.23)

$$= \frac{1}{4\pi} \int d^4 z \ e^{i(q+p_A-p_X)\cdot z} \ \oint \langle A|J^{\dagger}_{XA\mu}(0)|X\rangle \langle X|J_{XA\nu}(0)|A\rangle.$$
(A.24)

In the second line, we used the phase space identities of Eq. (A.2). Following Ref. [127] (and related references), we denote the 3^n -dimensional integration over momenta and triple summation by f. Importantly, the integration and summation constitute a summation over all possible configurations spanning the space of $\{|X\rangle\}$. Completeness dictates that

$$\oint |X\rangle\langle X| = \sum_{n=1}^{\infty} \int \prod_{i=1}^{n} \frac{d^3 p_{Xi}}{(2\pi)^3 2 p_{Xi}^0} \sum_{\text{dof } \{\lambda\}} |X\rangle\langle X| = 1.$$
(A.25)

Before employing Eq. (A.25), we use translation invariance to build a spacetime separation between $J_{A\mu}^{\dagger}$ and $J_{A\nu}$ that is constrained by momentum conservation. Such separations in the DIS limit are restricted to be near the light cone, i.e., with $z^2 \sim 0$ (*c.f.*, Ref. [64]). Applying translation invariance to the current $J_{XA\mu}^{\dagger}(0)$ gives

$$\langle A|J_{XA\mu}^{\dagger}(0)|X\rangle = \langle A|\left[e^{-i\hat{P}\cdot z}J_{XA\mu}(z)e^{i\hat{P}\cdot z}\right]^{\dagger}|X\rangle$$
(A.26)

$$= \langle A|e^{-i\hat{P}\cdot z}J^{\dagger}_{XA\mu}(z)e^{i\hat{P}\cdot z}|X\rangle = e^{-i(p_A - p_X)\cdot z} \langle A|J^{\dagger}_{XA\mu}(z)|X\rangle, \tag{A.27}$$

where \hat{P}^{α} is the four-momentum operator that generates the eigenvalue equation $\hat{P}^{\alpha}|Y(p_Y)\rangle = p_Y^{\alpha}|Y(p_Y)\rangle$. Combining the preceding expressions leads to familiar expression for $W_{\mu\nu}^A$:

$$W^{A}_{\mu\nu} = \frac{1}{4\pi} \int d^{4}z \; e^{iq \cdot z} \langle A | J^{\dagger}_{XA\mu}(z) \; J_{XA\nu}(0) | A \rangle \; . \tag{A.28}$$

In electromagnetism, currents are Hermitian. This leads to the omission of the \dagger in many texts. In other texts, Eq. (A.28) is written using a commutator of currents [63, 64]. However, Refs. [47, 85] note that commutators are not necessary to define $W^A_{\mu\nu}$. And indeed, Refs. [63, 64] show that the commutator-based result reduce to Eq. (A.28) in certain limits.

We are now in position to write the differential cross section for the DIS process $\ell_1 + A \rightarrow \ell_2 +$ anything in terms of the inclusive hadronic tensor $W^A_{\mu\nu}$. To do this, we take the differential cross section formula of Eq. (A.6) and insert the squared matrix element in Eq. (A.15). We then substitute the summations and integration over the exclusive hadronic tensor $H_{\mu\nu}$ with the inclusive hadronic tensor $W_{\mu\nu}$ to obtain the formula,

$$k_2^0 \times \frac{d\sigma}{d^3 k_2} = \frac{1}{(16\pi^2)s} \frac{2}{\mathcal{S}_{\ell_1} \mathcal{S}_A} \frac{1}{(q^2 - M_V^2)^2} \left(\sum_{\{\lambda\}} L^{\mu\nu}\right) \cdot W^A_{\mu\nu} .$$
(A.29)

This formula holds for arbitrary A, including unbound nucleons. Expressions for the leptonic tensor (summed over all external spins) are found in Eqs. (A.18)-(A.20). The factor of 2 in the numerator originates from the adopting the $(1/4\pi)$ normalization of $W_{\mu\nu}$; when A is a spin-1/2 object, it cancels the spin-averaging factor S_A . The inclusive hadronic tensor is given by Eq. (A.28), but now we relate it to the time-order matrix element for virtual Compton scattering, which has a known expansion in the OPE at leading twist.

Appendix A.3. Nuclear structure functions from the OPE I: organization

As summarized in Sec. 3.2, building structure functions in the OPE for massless and massive hadronic targets is a multi-step process. The advantage, however, is clear: the OPE, if it holds for QCD, facilitates an all-orders, operational definitions for parton densities and structure functions. Moreover, the power counting within the OPE formalism allows one to organize contributions that are not clearly captured in fixed-order perturbation theory.

To do this in the manner of Refs. [49, 50], one starts with the inclusive hadronic tensor $W^A_{\mu\nu}$ in the DIS limit (large Q, fixed x_A) as defined above and relates it to the time-ordered matrix element $T^A_{\mu\nu}$ for virtual Compton scattering¹³ in the short-distance limit (large Q, fixed x_A/Q). The OPE of $T^A_{\mu\nu}$ is then organized by Lorentz structures. By comparing the Lorentz structures of the expansions of $W^A_{\mu\nu}$ and $T^A_{\mu\nu}$, one can identify the structure functions of $W^A_{\mu\nu}$ in terms of the OPE. When the mass of a nucleus A is neglected, the OPE of $T^A_{\mu\nu}$ simplifies to familiar expressions. The differences between the fully massive and simplified massless expressions are the TMCs to the structure functions of $W^A_{\mu\nu}$.

Following this outline, we start by taking the expression for $W^A_{\mu\nu}(p_A, q)$ in Eq. (A.28) and decomposing it into a sum of tensor-valued coefficients multiplied by dimensionless, scalar-valued functions \tilde{W}^A_i . The \tilde{W}^A_i are the structure functions. Lorentz symmetry and hermiticity dictate that only certain combinations of p_A and q are allowed as coefficients. The most general combination allowed by symmetries for unpolarized A is

$$W_{\mu\nu}^{A}(p_{A},q) = \frac{1}{4\pi} \int d^{4}z \ e^{iq \cdot z} \langle A | J_{\mu}^{\dagger}(z) \ J_{\nu}(0) | A \rangle$$

$$= -g_{\mu\nu} \tilde{W}_{1}^{A} + \frac{p_{A\mu}p_{A\nu}}{M_{A}^{2}} \tilde{W}_{2}^{A} - i\epsilon_{\mu\nu\rho\sigma} \frac{p_{A}^{\rho}q^{\sigma}}{M_{A}^{2}} \tilde{W}_{3}^{A} + \frac{q_{\mu}q_{\nu}}{M_{A}^{2}} \tilde{W}_{4}^{A}$$

$$+ \frac{p_{A\mu}q_{\nu} + p_{A\nu}q_{\mu}}{M_{A}^{2}} \tilde{W}_{5}^{A} + \frac{p_{A\mu}q_{\nu} - p_{A\nu}q_{\mu}}{M_{A}^{2}} \tilde{W}_{6}^{A}$$
(A.30)
(A.30)
(A.31)

$$= -g_{\mu\nu}\tilde{F}_{1}^{A} + \frac{p_{A\mu}p_{A\nu}}{Q^{2}}2x_{A}\tilde{F}_{2}^{A} - i\epsilon_{\mu\nu\alpha\beta}\frac{p_{A}^{\alpha}q^{\beta}}{Q^{2}}x_{A}\tilde{F}_{3}^{A} + \frac{q_{\mu}q_{\nu}}{Q^{2}}2\tilde{F}_{4}^{A} + \frac{(p_{A\mu}q_{\nu} + p_{A\nu}q_{\mu})}{Q^{2}}2x_{A}\tilde{F}_{5}^{A} + \frac{(p_{A\mu}q_{\nu} - p_{A\nu}q_{\mu})}{Q^{2}}2x_{A}\tilde{F}_{6}^{A} .$$
(A.32)

Formally, the normalizations and organization of the \tilde{W}_i^A and \tilde{F}_i^A are conventional. Following Ref. [60], the \tilde{W}_i^A are normalized such that each is dimensionless and have at most a prefactor of M_A^{-2} . Subsequently, the \tilde{F}_i^A are normalized such that

 $^{{}^{13}\}gamma^*(q)A(p_A) \rightarrow \gamma^*A$ for electromagnetic currents, $W^*(q)A(p_A) \rightarrow W^*A$ for charged currents, etc.

$$\tilde{F}_{1}^{A} = \tilde{W}_{1}^{A}, \qquad (A.33a) \qquad \tilde{F}_{3}^{A} = \left(\frac{Q^{2}}{x_{A}M_{A}^{2}}\right) \tilde{W}_{3}^{A}, \qquad (A.33c) \\
\tilde{F}_{i}^{A} = \left(\frac{Q^{2}}{2x_{A}M_{A}^{2}}\right) \tilde{W}_{i}^{A}, \quad \text{for} \quad i = 2, 5, 6 \quad (A.33b) \qquad \tilde{F}_{4}^{A} = \left(\frac{Q^{2}}{2M_{A}^{2}}\right) \tilde{W}_{4}^{A}, \qquad (A.33d)$$

Using \tilde{F}_i^A instead of \tilde{W}_i^A factors out known Q^2 dependence from the structure functions. This makes scaling with respect to x_A more manifest. Under this normalization, \tilde{W}_i^A and \tilde{F}_i^A are real for i = 1-5, whereas \tilde{W}_6^A and \tilde{F}_6^A are imaginary [85]. Moreover, \tilde{W}_6^A and \tilde{F}_6^A are only nonzero if charge-parity symmetry is violated in QCD. However, the coefficient of \tilde{W}_6^A and \tilde{F}_6^A will vanish when contracted with the symmetric $L^{\mu\nu}$ tensor. Choosing \tilde{W}_i^A , \tilde{F}_i^A , or other normalizations does not change the underlying physics. However, seemingly innocuous differences in their definitions can impact the final form of TMCs due to structure function mixing, i.e., the off-diagonal terms in A_i^i and B_i^i in Eq. (3.23).

The next step is to recognize that $\tilde{W}^A_{\mu\nu}$ in the DIS limit can be related to the time-ordered matrix element $T^A_{\mu\nu}$ for $AV^* \to AV^*$ scattering in short-distance limit. This is given by [40]

$$T^{A}_{\mu\nu}(p_{A},q) \equiv \int d^{4}z \ e^{iq\cdot z} \ \langle A|\mathcal{T}J^{\dagger}_{\mu}(z) \ J_{\nu}(0)|A\rangle$$

$$= \int d^{4}z \ e^{iq\cdot z} \ \langle A|J^{\dagger}_{\mu}(z) \ J_{\nu}(0)|A\rangle \ \theta(z^{0} > 0)$$
(A.34)

+
$$\int d^4 z \; e^{iq \cdot z} \langle A | J_{\nu}(0) \; J^{\dagger}_{\mu}(z) | A \rangle \; \theta(z^0 < 0)$$
 (A.35)

$$\equiv T_{z^0>0}(p_A, q) + T_{z^0<0}(p_A, q), \tag{A.36}$$

where θ is the usual Heaviside step function normalized to unity. The matrix elements under the integral can be identified as the (inverse) Fourier transforms (FT) of $W^A_{\mu\nu}$ over q:

$$\operatorname{FT}[W^{A}_{\mu\nu}(p_{A},q)](x) = \int \frac{d^{4}q}{(2\pi)^{4}} e^{-iq \cdot x} W^{A}_{\mu\nu}(p_{A},q) = \frac{1}{4\pi} \langle A|J^{\dagger}_{\mu}(x) J_{\nu}(0)|A\rangle , \qquad (A.37a)$$

$$\operatorname{FT}[\overline{W}^{A}_{\nu\mu}(p_{A},q)](x) = \int \frac{d^{4}q}{(2\pi)^{4}} e^{-iq \cdot x} \overline{W}^{A}_{\nu\mu}(p_{A},q) = \frac{1}{4\pi} \langle A|J_{\nu}(0) J^{\dagger}_{\mu}(x)|A\rangle .$$
(A.37b)

Equation (A.37b) defines the "conjugate" hadronic tensor $\overline{W}^A_{\nu\mu}$. It is related to $W^A_{\mu\nu}$ by

$$\overline{W}^{A}_{\nu\mu}(p_{A},q) = \frac{1}{4\pi} \int d^{4}z \; e^{-i(-q)\cdot z} \langle A|J_{\nu}(0) \; J^{\dagger}_{\mu}(z)|A\rangle = \left[W^{A}_{\mu\nu}(p_{A},-q)\right]^{\dagger} \; . \tag{A.38}$$

Note that the Hermitian conjugation operator † should be applied to both the structure functions and the tensor-valued coefficients of $W^A_{\mu\nu}$. For the case of electromagnetic currents, these are Hermitian and Eq. (A.38) reduces to $\overline{W}^{A}_{\nu\mu}(p_{A},q)|_{\rm EM} = W^{A}_{\mu\nu}(p_{A},-q).$ After exchanging the order of integration, the first term in Eq. (A.36) is

$$T_{z^0>0}(p_A,q) = 4\pi \int \frac{d^4k}{(2\pi)^4} W^A_{\mu\nu}(p_A,k) \int d^4z \ e^{i(q-k)\cdot z} \ \theta(z^0>0)$$
(A.39)

$$= 4\pi \int \frac{d^4k}{(2\pi)^4} W^A_{\mu\nu}(p_A,k) (2\pi)^3 \delta(\vec{q}-\vec{k}) \left[\pi \delta(q^0-k^0) + \frac{i}{q^0-k^0}\right].$$
(A.40)

The three-dimensional $\delta(\vec{q} - \vec{k})$ comes from integrating over all of \vec{z} -space and the bracketed factor

comes from integrating only over $z^0 > 0$. Following the argument of Ref. [85], momentum conservation dictates that $(2p_A \cdot q) > Q^2$, which implies that $W^A_{\mu\nu}$ is zero for $(2p_A \cdot q) < Q^2$, or equivalently that $W^A_{\mu\nu}$ is zero for $x_A > 1$ and $x_A < 0$. Fixing Q^2 and taking $W^A_{\mu\nu}$ to be analytic on the plane $\omega_A \equiv x_A^{-1}$ further implies branch cuts along $|\omega_A| \ge 1$ as depicted in Fig. 3.1. Therefore, since the pole in Eq. (A.40) can be decomposed into its principal \mathcal{P} and singular parts when deformed, i.e.,

$$\frac{1}{q^0 - k^0 \mp i\varepsilon} = \mathcal{P} \pm i\pi\delta(q^0 - k^0), \qquad (A.41)$$

the tensor $W^A_{\mu\nu}$ can be identified as the discontinuity of $T_{z^0>0}$. Specifically, one finds:

disc
$$T_{z^{0}>0} = \lim_{\varepsilon \to 0} \left[T_{z^{0}>0}(q^{0} + i\varepsilon) - T_{z^{0}>0}(q^{0} - i\varepsilon) \right]$$
 (A.42)

$$= 4\pi \int \frac{d^{4}k}{(2\pi)^{4}} W^{A}_{\mu\nu}(p_{A}, k) (2\pi)^{3} \delta(\vec{q} - \vec{k})$$

$$\times \left\{ \left[i\mathcal{P} - i^{2}\pi\delta(q^{0} - k^{0}) \right] - \left[i\mathcal{P} + i^{2}\pi\delta(q^{0} - k^{0}) \right] \right\}$$
(A.43)

$$= 4\pi W^{A}(p_{A}, q)$$
(A.44)

$$= 4\pi \ W_{\mu\nu}(p_A, q) \ . \tag{A.44}$$

A similar result holds for $T_{z^0<0}$. Subsequently, one can write in terms of $\omega_A = x_A^{-1}$ the relations [40, 85]

$$T^{A}_{\mu\nu}(p_{A},q)\Big|_{\omega_{A}-i\varepsilon}^{\omega_{A}+i\varepsilon} = 4\pi \ W^{A}_{\mu\nu}(p_{A},q), \quad \text{for} \quad \omega_{A} > 0 \ , \tag{A.45a}$$

$$T^{A}_{\mu\nu}(p_{A},q)\Big|_{\omega_{A}+i\varepsilon}^{\omega_{A}-i\varepsilon} = 4\pi \left[W^{A}_{\mu\nu}(p_{A},-q)\right]^{\dagger}, \quad \text{for} \quad \omega_{A} < 0 \ . \tag{A.45b}$$

and with $W^A_{\mu\nu}$ and $W^{A\dagger}_{\nu\mu}$ vanishing for $\omega_A < 1$ and $\omega_A > -1$, respectively. Intuitively, Eq. (A.45) states that $T^A_{\mu\nu}$ and $W^{A\dagger}_{\mu\nu}$, which are defined in different kinematic limits (the short-distance and DIS limits, respectively), are nevertheless related through analytic continuation. Therefore, one can take $T^A_{\mu\nu}(p_A,q)$, decompose it into a contour integral over x^{-1}_A using Cauchy's integral formula, and deform the contour around the discontinuities at $x^{-1}_A < 1$ and $x^{-1}_A > 1$ as shown in Fig. 3.1.

Coefficient functions of $T^A_{\mu\nu}$ and moments of structure functions

The relationship in Eq. (A.45) can be refined by decomposing $T^A_{\mu\nu}$ in three different ways. The first is according to Lorentz structures as was done for $W^A_{\mu\nu}$ in Eq. (A.31):

$$T_{\mu\nu}^{A} = -g_{\mu\nu}\Delta\tilde{T}_{1}^{A} + \frac{p_{A\mu}p_{A\nu}}{M_{A}^{2}}\Delta\tilde{T}_{2}^{A} - i\epsilon_{\mu\nu\alpha\beta}\frac{p_{A}^{\alpha}q^{\beta}}{M_{A}^{2}}\Delta\tilde{T}_{3}^{A} + \frac{q_{\mu}q_{\nu}}{M_{A}^{2}}\Delta\tilde{T}_{4}^{A} + \frac{(p_{A\mu}q_{\nu} + p_{A\nu}q_{\mu})}{M_{A}^{2}}\Delta\tilde{T}_{5}^{A} + \frac{(p_{A\mu}q_{\nu} - p_{A\nu}q_{\mu})}{M_{A}^{2}}\Delta\tilde{T}_{6}^{A}.$$
 (A.46)

Component-by-component, the dispersion relationship of Eq. (A.45) becomes

$$\Delta \tilde{T}_{i}^{A}(p_{A},q)\Big|_{\omega_{A}-i\varepsilon}^{\omega_{A}+i\varepsilon} = 4\pi \ \tilde{W}_{i}^{A}(p_{A},q), \quad \text{for} \quad \omega_{A} > 0 \ , \tag{A.47a}$$

$$\Delta \tilde{T}_i^A(p_A, q) \Big|_{\omega_A + i\varepsilon}^{\omega_A - i\varepsilon} = 4\pi \ (-1)^{b_i} \ \tilde{W}_i^A(p_A, -q), \quad \text{for} \quad \omega_A < 0 \ , \tag{A.47b}$$

and
$$b_i = 0$$
 (1), for $i = 1, 2, 3, 4, 6$ (5). (A.47c)

The factor of $(-1)^{b_i}$ comes from applying the \dagger operator in Eq. (A.45b) to \tilde{W}_i^A and its prefactor in combination with propagating the argument -q into the prefactor of \tilde{W}_i^A . We also take into account whether \tilde{W}_i is real or imaginary, as discussed below Eq. (A.33); recall \tilde{W}_6^A is imaginary.

The second way of decomposing $T^A_{\mu\nu}$ is by the OPE (done below) and gives expressions for $\Delta \tilde{T}^A_i$. The third decomposition is by Taylor expansion in the short-distance limit. The distinction between this limit and the DIS limit is important: In the DIS limit, $(Q^2/M_A^2) \rightarrow \infty$ while $x_A = (Q^2/2p_A \cdot q)$ is fixed. In the short-distance limit, $(Q^2/M_A^2) \rightarrow \infty$ while x_A/Q is fixed, meaning that x_A grows with (Q/M_A) . Formally, the short-distance limit corresponds to the limit $\omega_A = 2(p_A \cdot q)/Q^2 \rightarrow 0$. We therefore assume that we can write

$$\zeta_i^A \times \omega_A^{a_i} \times \Delta \tilde{T}_i^A(p_A, q) = \sum_{k=0}^{\infty} t_{i(k)}^A(Q^2) \,\omega_A^k \,, \quad \text{where}$$
(A.48a)

$$a_i = 0 \ (1) \text{ for } i = 1,4 \ (2,3,5,6) \ , \text{ and } \zeta_1^A = 1 \ , \zeta_{2,4,5,6}^A = \left(\frac{Q^2}{2M_A^2}\right) \ , \zeta_3^A = \left(\frac{Q^2}{M_A^2}\right) \ .$$
 (A.48b)

Here, $t_{i(k)}^{A}$ is the k^{th} coefficient of the expansion and is only a function of Q^2 . The factors ζ_i^A and $\omega_A^{a_i}$, which are fixed in the short distance limit, are the same factors relating the structure functions \tilde{W}_i and \tilde{F}_i in Eqs. (2.9) and (A.33). Following Ref. [85] and using Cauchy's integral formula

$$F(y) = \frac{1}{2\pi i} \int_{\mathcal{C}} \frac{dz}{z - y} F(z), \qquad (A.49)$$

where \mathcal{C} is an appropriately chosen contour, it is possible to analytically continue $\Delta \tilde{T}_i^A$:

$$\zeta_i^A \times \omega_A^{a_i} \times \Delta \tilde{T}_i^A(p_A, q) = \frac{\zeta_i^A}{2\pi i} \int_{\mathcal{C}} \frac{d\omega_A'}{(\omega_A' - \omega_A)} (\omega_A')^{a_i} \Delta \tilde{T}_i^A(p_A, q')$$
(A.50)

$$= \frac{\zeta_i^A}{2\pi i} \int_{\mathcal{C}} \frac{d\omega'_A}{\omega'_A (1 - \frac{\omega_A}{\omega'_A})} (\omega'_A)^{a_i} \Delta \tilde{T}_i^A(p_A, q')$$
(A.51)

$$= \sum_{k=0}^{\infty} \frac{\zeta_i^A}{2\pi i} \int_{\mathcal{C}} d\omega'_A \left(\frac{\omega_A}{\omega'_A}\right)^k (\omega'_A)^{a_i-1} \Delta \tilde{T}_i^A(p_A, q').$$
(A.52)

The geometric series is obtained by fixing the integration variable ω'_A , which can be large, but taking the external variable $\omega_A \to 0$.

Comparing Eq. (A.52) to Eq. (A.48) and using Eqs. (A.47), one obtains

$$t_{i(k)}^{A}(Q^{2}) = \frac{\zeta_{i}^{A}}{2\pi i} \int_{\mathcal{C}} d\omega_{A}' (\omega_{A}')^{a_{i}-k-1} \Delta \tilde{T}_{i}^{A}(p_{A},q')$$

$$= \frac{4\pi \zeta_{i}^{A}}{2\pi i} \int_{1}^{\infty} d\omega_{A}' (\omega_{A}')^{a_{i}-k-1} \tilde{W}_{i}^{A}(p_{A},q')$$

$$+ \frac{4\pi \zeta_{i}^{A}}{2\pi i} \int_{-\infty}^{-1} d\omega_{A}' (\omega_{A}')^{a_{i}-k-1} (-1)^{b_{i}} \tilde{W}_{i}^{A}(p_{A},-q')$$

$$= (-2i)\zeta_{i}^{A} \int_{0}^{1} dx_{A}' (x_{A}')^{k-1-a_{i}} \tilde{W}_{i}^{A}(x_{A}',Q^{2})$$

$$+ (-2i)\zeta_{i}^{A} \int_{-1}^{0} dx_{A}' (x_{A}')^{k-1-a_{i}} (-1)^{b_{i}} \tilde{W}_{i}^{A}(-x_{A}',Q^{2}) .$$
(A.53)

The last line is obtained by changing to the variable $x'_A = (\omega'_A)^{-1}$. Changing the integration variable

of the second term to $z_A = -x'_A$, collecting factors of (-1), using $(-1)^{a_i} = (-1)^{-a_i}$, and relabeling gives

$$t_{i(k)}^{A}(Q^{2}) = (-2i) \int_{0}^{1} dx'_{A} (x'_{A})^{k-1} \\ \times \left[\zeta_{i} (x'_{A})^{-a_{i}} \tilde{W}_{i}^{A}(x'_{A}, Q^{2}) + (-1)^{k-1+a_{i}+b_{i}} \zeta_{i} (x'_{A})^{-a_{i}} \tilde{W}_{i}^{A}(x'_{A}, Q^{2}) \right]$$
(A.56)

$$= (-2i) \left[1 + (-1)^{k-1+a_i+b_i} \right] \int_0^1 dx'_A (x'_A)^{k-1} \tilde{F}_i^A(x'_A, Q^2).$$
(A.57)

We now denote the N^{th} Mellin moment of the function M(z) by M^N and fix normalizations such that a Mellin transformation and its inverse (over a path c) are:

$$M^{N} = \int_{0}^{1} dz \ z^{N-1} \ M(z) \quad \text{with} \quad M(z) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} dN \ z^{-N} \ M^{N}.$$
(A.58)

Under this normalization, the moments of $\tilde{W}_i^A(x_A)$ are related to those of $\tilde{F}_i^A(x_A)$ by

$$\tilde{F}_{1}^{AN} = \tilde{W}_{1}^{AN} , \qquad (A.59a) \qquad \tilde{F}_{3}^{AN} = \left(\frac{Q^{2}}{M_{A}^{2}}\right) \tilde{W}_{3}^{A(N-1)} , \qquad (A.59c)$$

$$\tilde{F}_{i}^{AN} = \left(\frac{Q^{2}}{2M_{A}^{2}}\right) \tilde{W}_{i}^{A(N-1)} \quad \text{for} \quad i = 2, 5, 6 , \quad (A.59b) \qquad \tilde{F}_{4}^{AN} = \left(\frac{Q^{2}}{2M_{A}^{2}}\right) \tilde{W}_{4}^{AN} . \tag{A.59d}$$

Both are related to the N^{th} coefficient function $t^A_{i(k=N)}(Q^2)$ by

$$t_{i(N)}^{A}(Q^{2}) = -2i \left[1 + (-1)^{N-1+a_{i}+b_{i}}\right] \tilde{F}_{i}^{AN}(Q^{2})$$
(A.60a)

$$= \begin{cases} 0, & N = \text{even} \\ -4i \ \tilde{F}_i^{AN}(Q^2), & N = \text{odd} \end{cases}, \quad \text{for} \quad i = 1, 4, 5 , \qquad (A.60b)$$

$$= \begin{cases} -4i \ \tilde{F}_i^{AN}(Q^2), \ N = \text{even} \\ 0, \qquad N = \text{odd} \end{cases}, \quad \text{for} \quad i = 2, 3, 6 \ . \tag{A.60c}$$

This allows us to rewrite the expansion in Eq. (A.48) as

$$\zeta_i^A \times \omega_A^{a_i} \times \Delta \tilde{T}_i^A(Q^2, \omega_A) = -4i \sum_N^\infty \tilde{F}_i^{AN}(Q^2) \,\omega_A^N \,, \tag{A.61}$$

where it is implied that the index N runs only over odd or even integers.

We now make a few brief comments. First, in order for Eq. (A.61) to hold, $|\omega_A| < 1$ must be satisfied, i.e., one must be in the DIS limit. Second, as N increases, the dominant contribution to \tilde{F}_i^{AN} is when the argument of $\tilde{F}_i^A(z)$ approaches unity since (according to the definition of Eq. (A.58)) \tilde{F}_i^{AN} would otherwise be suppressed by a small number. Finally, since $\omega_A = 1$ corresponds to the elastic limit (see Eq. (2.2c)), Q^2 must be made increasingly large for large-N moments to be well-defined and for the invariant mass of the hadronic system to remain in the perturbative regime.

Expanding $T^A_{\mu\nu}$ with the OPE

Using the OPE, the leading behavior of $T^A_{\mu\nu}$ in powers of $(1/Q^p)$ can be decomposed into a sum of operators $\mathcal{O}^{\mu_1,\ldots,\mu_k}$ and Wilson coefficients $c^{\tau,\epsilon}_{\mu\nu\mu_1\ldots,\mu_k}$. This expansion is given by [40, 49, 50]

$$\lim_{z \to 0} T^A_{\mu\nu}(p_A, q) \stackrel{\text{OPE}}{=} -2i \sum_{k,\iota} c^{\tau=2,\iota}_{\mu\nu\mu_1\dots\mu_k}(q) \left\langle A(p_A) | \mathcal{O}^{\mu_1\dots\mu_k}_{\iota,\tau=2} | A(p_A) \right\rangle + \mathcal{O}(\tau > 2) .$$
(A.62)

At leading power, the composite operators $\mathcal{O}_{\iota,\tau}^{\mu_1...\mu_k}$ are symmetric, quark billinears (or pairs of gluon field strengths) that sandwich uncontracted covariant derivatives. Tallying up the number of uncontracted Lorentz indicies implies that $\mathcal{O}_{\iota,\tau}^{\mu_1...\mu_k}$ carries a spin of k. The operators are organized according to their twist $\tau \equiv d - n$, where d is the dimensionality of $\mathcal{O}_{\iota,\tau}^{\mu_1...\mu_k}$ in the standard sense of dimension power counting in an effective field theory. For a fixed spin and twist, the index ι catalogs all the Lorentz structures that can possibly contract with $\mathcal{O}_{\iota,\tau}^{\mu_1...\mu_k}$. Coupling and renormalization factors are sequestered into the effective Wilson coefficient $c_{\mu\nu\mu_1...\mu_k}^{\tau,\iota}(q)$. For fixed spin k, a larger twist τ corresponds to a larger (1/Q) suppression in the Wilson coefficient. Schematically, Eq. (A.62) stipulates that in the short-distance limit, the time-ordered matrix element $T_{\mu\nu}^A$, which is a function of p_A and q, can be expressed in terms of Wilson coefficients, which are only functions of q, and hadronic matrix elements, which are only functions of p_A .

The operators $\mathcal{O}^{\mu_1,\ldots,\mu_k}$ can be decomposed into symmetric (traceless) and non-symmetric parts, with $\mathcal{O}^{\mu_1,\ldots,\mu_k} \sim \hat{P}^{\mu_1}_A \ldots \hat{P}^{\mu_k}_A$ + Tr. The trace term captures all contributions proportional to the spacetime metric and derivatives, which after contractions or applications of equations of motion give rise to powers of quark masses [49, 50]. Neglecting quark masses¹⁴, the matrix elements that follow from acting on $\mathcal{O}^{\mu_1,\ldots,\mu_{2k}}$ are given by

$$\langle A|\mathcal{O}_{\iota,\tau=2}^{\mu_1\dots\mu_{2k}}|A\rangle = A_{\tau=2}^{2k} \times \tilde{\Pi}^{\mu_1\dots\mu_{2k}}, \quad \text{where}$$
(A.63a)

$$\tilde{\Pi}^{\mu_1\dots\mu_{2k}} = \sum_{j=0}^k (-1)^j \; \frac{(2k-j)!}{2^j (2k)!} \; \eta(j, 2k-2j) \underbrace{\{g...g\}}_{j \; g^{\mu_n \mu_m's}} \; \underbrace{\{p_A...p_A\}}_{(2k-2j) \; p_A^{\mu_n's}} \; (p_A^2)^j \; . \tag{A.63b}$$

Note the index change from k to 2k. The factor $A_{\tau=2}^{2k}$ is the scalar-valued "reduced" hadronic matrix element and describes long-distance hadronic dynamics. (The 2k in $A_{\tau=2}^{2k}$ is an index, not an exponent.) The index j sums over all permutations of $p_A^{\mu_n}$ and $g^{\mu_n\mu_m}$, with $\mu_n, \mu_m \in {\mu_1, \ldots, \mu_{2k}}$, that contract with a particular Wilson coefficient. For a given k and j, the two ${\ldots}$ brackets denote j spacetime metrics $g^{\mu_n\mu_m}$ and (2k-2j) factors of momentum $p_A^{\mu_m}$. The permutation multiplicity η is

$$\eta(j \text{ metrics}, 2k-2j \text{ factors of } p_A) = \frac{1}{2^j} \frac{(2k)!}{j!(2k-2j)!}$$
 (A.64)

The numerator of η is calculated from (2k)! = [(2k-2j)+(2j)]! The denominator factor of 2^j accounts for the two-fold symmetry of j symmetric metric tensors, i.e., $g^{\mu_n\mu_m} = g^{\mu_m\mu_n}$. If a metric or momentum factor is pulled from either $\{\dots\}$ bracket, η is updated accordingly.

In the massless target limit, i.e., when $(M_A^2/Q^2) \to 0$, one neglects j > 0 terms since they generate powers of $(p_A^2)^j = M_A^{2j}$. For this reason, the j > 0 terms are sometimes called "kinematical power corrections" [55, 83]. In the j = 0 limit, Eq. (A.63) reduces to 2k factors of $p_A^{\mu_m}$:

$$\left. \left\langle A | \mathcal{O}_{\iota,\tau=2}^{\mu_1,\dots,\mu_{2k}} | A \right\rangle \right|_{(M_A/Q)^2 \to 0} = A_{\tau=2}^{2k}(p_A^2) \times \tilde{\Pi}_{\mu_1\dots,\mu_{2k}} \Big|_{j=0} = A_{\tau=2}^{2k}(p_A^2) \times (p_A^{\mu_1}\dots p_A^{\mu_{2k}}) . \tag{A.65}$$

¹⁴In practice, finite quark masses can be incorporated in TMCs through a rescaling of the Nachtmann scaling variable. See Appendix A.2 of Ref. [60] and references therein.

At leading power of τ and for a fixed 2k > 2, the c functions in Eq. (A.62) can be decomposed as

$$c_{\mu\nu\mu_{1},\dots,\mu_{2k}}^{\tau=2,\iota}(q) = \left[-2g_{\mu\nu}q_{\mu_{1}}q_{\mu_{2}}C_{1}^{2k} + g_{\mu\mu_{1}}g_{\nu\mu_{2}}Q^{2}C_{2}^{2k} - i\epsilon_{\mu\nu\alpha\beta}g_{\mu_{1}}^{\alpha}q^{\beta}q_{\mu_{2}}C_{3}^{2k} + 4\frac{q_{\mu}q_{\nu}}{Q^{2}}q_{\mu_{1}}q_{\mu_{2}}C_{4}^{2k} + 2(g_{\mu\mu_{1}}q_{\nu}q_{\mu_{2}} \pm g_{\nu\mu_{1}}q_{\mu}q_{\mu_{2}})C_{5,6}^{2k}\right] \times \frac{2^{2k}}{(Q^{2})^{2k}} \times \left(\prod_{m=3}^{2k}q_{\mu_{m}}\right).$$
(A.66)

Here, the $C_{\iota=1,\ldots,6}^{2k}$ are scalar-valued coefficients that parameterize the normalization of each $c_{\mu\nu\mu_1,\ldots,\mu_{2k}}^{\tau=2,\iota}$. The C_{ι} are defined to all orders in QCD but can be identified and matched to quantities in fixed-order perturbation theory. The tensor part, i.e., the part carrying Lorentz indices, can then be organized according to Lorentz structures as in Eq. (A.46). Conventional factors of 2 in Eq. (A.66) are pulled from the C_{ι}^{2k} in order to simplify later expressions.

Assembling these ingredients and contracting over all $2k > k_{\min}$ indices, the OPE gives

$$\lim_{z \to 0} T^{A}_{\mu\nu}(q, p_{A}) \stackrel{\text{OPE}}{=} -2i \sum_{k=k_{\min}}^{\infty} \left[-2g_{\mu\nu}q_{\mu_{1}}q_{\mu_{2}}C_{1}^{2k} + g_{\mu\mu_{1}}g_{\nu\mu_{2}}Q^{2}C_{2}^{2k} - i\epsilon_{\mu\nu\alpha\beta}g^{\alpha}_{\mu_{1}}q^{\beta}q_{\mu_{2}}C_{3}^{2k} + 4\frac{q_{\mu}q_{\nu}}{Q^{2}}q_{\mu_{1}}q_{\mu_{2}}C_{4}^{2k} + 2(g_{\mu\mu_{1}}q_{\nu}q_{\mu_{2}} \pm g_{\nu\mu_{1}}q_{\mu}q_{\mu_{2}})C_{5,6}^{2k} \right] \\ \times \frac{2^{2k}}{(Q^{2})^{2k}} \times \left(\prod_{m=3}^{2k}q_{\mu_{m}}\right) \times A^{2k}_{\tau=2}(p^{2}_{A}) \times \tilde{\Pi}^{\mu_{1}\dots\mu_{2k}} + \mathcal{O}(\tau > 2)$$
(A.67)

$$\equiv \Delta \tilde{T}^{A}_{1\mu\nu} + \Delta \tilde{T}^{A}_{2\mu\nu} + \Delta \tilde{T}^{A}_{3\mu\nu} + \Delta \tilde{T}^{A}_{4\mu\nu} + \Delta \tilde{T}^{A}_{5\mu\nu} + \Delta \tilde{T}^{A}_{6\mu\nu} + \mathcal{O}(\tau > 2)$$
(A.68)

$$= -g_{\mu\nu}\Delta\tilde{T}_{1}^{A} + \frac{p_{A\mu}p_{A\nu}}{M_{A}^{2}}\Delta\tilde{T}_{2}^{A} - i\epsilon_{\mu\nu\alpha\beta}\frac{p_{A}q^{\prime}}{M_{A}^{2}}\Delta\tilde{T}_{3}^{A} + \frac{q_{\mu}q_{\nu}}{M_{A}^{2}}\Delta\tilde{T}_{4}^{A}$$
$$+ \frac{(p_{A\mu}q_{\nu} + p_{A\nu}q_{\mu})}{M_{A}^{2}}\Delta\tilde{T}_{5}^{A} + \frac{(p_{A\mu}q_{\nu} - p_{A\nu}q_{\mu})}{M_{A}^{2}}\Delta\tilde{T}_{6}^{A} + \mathcal{O}(\tau > 2).$$
(A.69)

The starting point for the summation over k depends on the particular Wilson coefficient; specifically, $k_{\min} = 2$ for C_2 while $k_{\min} = 1$ for the other C_i coefficients. In Eq. (A.69), the scalar-valued coefficients $\Delta \tilde{T}_{i=1,\dots,6}^A$, i.e., the quantities without external Lorentz indices, are related to the hadronic structure functions W_i^A through the dispersion relationships of Eqs. (A.45) and (A.47), up to (1/Q) corrections. The $\Delta \tilde{T}_i$ are given explicitly in terms of summations over k and j below in Eqs. (A.96)-(A.106).

In the intermediate step Eq. (A.68), each $\Delta \tilde{T}_{i=1,\dots,6}^{A} \mu \nu$ denotes the collection of contractions that are respectively proportional to the coefficient C_{ι} . We introduce this step because $\Delta \tilde{T}_{i}^{A} \mu \nu$ and $\Delta \tilde{T}_{i}^{A}$ do not have a one-to-one correspondence when $M_{A} \neq 0$. As a consequence of the normalizations adopted for \tilde{W}_{i}^{A} in Eq. (A.31), and subsequently those for $\Delta \tilde{T}_{i}^{A}$ in Eq. (A.46), some $\Delta \tilde{T}_{i}^{A}$ are sourced by multiple OPE operators when $M_{A} \neq 0$. This phenomenon of \tilde{W}_{i}^{A} (or $\Delta \tilde{T}_{i}^{A}$) being sources by two or more C_{ι} is sometimes called "structure function mixing," and is discussed in Appendix A.4.

We now derive all six $\Delta \tilde{T}^A_{\iota \ \mu\nu}$ from Eq. (A.67) by contracting all μ_m indices. After, we reorganize $\Delta \tilde{T}^A_{\iota \ \mu\nu}$ and group terms according to their Lorentz structure, e.g., collect all terms proportional to $g_{\mu\nu}$ or all terms proportional to $\epsilon_{\mu\nu\alpha\beta}p^{\alpha}_Aq^{\beta}$. We then identify the $\Delta \tilde{T}^A_i$, which are in terms of hadronic matrix elements $A^{2k}_{\tau=2}$, Wilson coefficients C^{2k}_{ι} , and nested summations. In Appendix A.5, the summations are first evaluated for the case of a massless target. The results for the $(M_A/Q^2) \neq 0$ case are then expressed in terms of structure functions for massless targets.

Organization of Lorentz contractions in the OPE

Explicit evaluation of all contractions of Lorentz indices in Eq. (A.67) is an exacting task. Therefore, as a first step, we organize everything according to $\Delta \tilde{T}^A_{\iota \ \mu\nu}$, and hence C_{ι} . As a consequence, patterns in $\Delta \tilde{T}^A_{\iota \ \mu\nu}$ appear and simplify the work. The outcome is summarized in Appendix A.4.

(i) $\Delta \tilde{T}^{A}_{1\mu\nu}$: We start with $\Delta \tilde{T}^{A}_{1\mu\nu}$ for $M_A \neq 0$ because, like $\Delta \tilde{T}^{A}_{4\mu\nu}$, the covariant tensor structure for $\mu_{m=1} \dots \mu_{m=2k}$ consists entirely of momentum factors q. That is to say, all lowered internal indices μ_m are carried by q_{μ_m} . This results in the simplest possible permutations of metrics and momenta in $\tilde{\Pi}^{\mu_1\dots\mu_{2k}}$. Combining terms, one gets

$$\Delta \tilde{T}^{A}_{1\mu\nu} = -4i \sum_{k=1}^{\infty} \left[-g_{\mu\nu} C_{1}^{2k} A_{\tau=2}^{2k} \right] \times \frac{2^{2k}}{(Q^{2})^{2k}} \times \left(\prod_{m=1}^{2k} q_{\mu_{m}} \right) \times \tilde{\Pi}^{\mu_{1}\dots\mu_{2k}} + \mathcal{O}(\tau > 2), \tag{A.70}$$

where the production factor $(\prod_{m=1}^{2k} q_{\mu_m})$ has (re)absorbed q_{μ_1} and q_{μ_2} .

The contraction of $\prod_{m=1}^{2k} q_{\mu_m}$ and the momentum factor $\tilde{\Pi}$ combines all possible permutations of 2k instances of q_{μ_m} with j metrics $g^{\mu_m\mu_n}$ in $\{g \dots g\}$ and (2k-2j) momenta $p_A^{\mu_m}$ in $\{p_A \dots p_A\}$. This results in (2k-2j) products of $(q \cdot p_A)$. The remaining (2k-0) - (2k-2j) = 2j instances of q_{μ_m} then contract with the 2j indices from j metrics. As no instance of $g^{\mu_m\mu_n}$ or $p_A^{\mu_m}$ have been pulled from $\tilde{\Pi}$, its permutation factor η is the same as in Eq. (A.64). The outcome is

$$\begin{pmatrix}
\sum_{m=1}^{2k} q_{\mu_m} \\
\prod^{\mu_1 \dots \mu_{2k}} = \sum_{j=0}^{k} (-1)^j \frac{(2k-j)!}{2^j (2k)!} \frac{1}{2^j} \frac{(2k)!}{j! (2k-2j)!} \\
\times \qquad (q^2)^j (q \cdot p_A)^{(2k-2j)} (p_A^2)^j$$
(A.71)

$$= \frac{(Q^2)^{2k}}{2^{2k}} \sum_{j=0}^{k} \frac{(2k-j)!}{j!(2k-2j)!} \left(\frac{M_A^2}{Q^2}\right)^j x_A^{-(2k-2j)} , \qquad (A.72)$$

where $x_A = Q^2/(2p_A \cdot q)$ is the Bjorken scaling variable and $p_A^2 = M_A^2$. Altogether,

=

$$\Delta \tilde{T}^{A}_{1\mu\nu} = -g_{\mu\nu} \times (-4i) \sum_{k=1}^{\infty} \left[C^{2k}_{1} A^{2k}_{\tau=2} \right] \sum_{j=0}^{k} \frac{(2k-j)!}{j!(2k-2j)!} \left(\frac{M^{2}_{A}}{Q^{2}} \right)^{j} x^{-(2k-2j)}_{A} + \mathcal{O}(\tau > 2).$$
(A.73)

We stress that the quantity to the right of $g_{\mu\nu}$ here is not $\Delta \tilde{T}_1^A$ in Eq. (A.69). In general, the coefficients $\Delta \tilde{T}_i^A$ receive contributions from more than one $\Delta \tilde{T}_{\iota \ \mu\nu}^A$.

(ii) $\Delta \tilde{T}^{A}_{4\mu\nu}$: The next term we construct is $\Delta \tilde{T}^{A}_{4\mu\nu}$ for $M_A \neq 0$, which has an identical tensor structure of contracted covariant indices as $\Delta \tilde{T}^{A}_{1\mu\nu}$. In other words, like $\Delta \tilde{T}^{A}_{1\mu\nu}$, all the lowered internal indices μ_m are carried exclusively by factors of q_{μ_m} . The result is then,

$$\Delta \tilde{T}^{A}_{4\mu\nu} = \frac{q_{\mu}q_{\nu}}{Q^{2}} \times (-8i) \sum_{k=1}^{\infty} \left[C_{4}^{2k} A_{\tau=2}^{2k} \right] \sum_{j=0}^{k} \frac{(2k-j)!}{j!(2k-2j)!} \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} x_{A}^{-(2k-2j)} + \mathcal{O}(\tau > 2).$$
(A.74)

(iii) $\Delta T_{3\mu\nu}^A$: Moving onto $\Delta T_{3\mu\nu}^A$ for $M_A \neq 0$, exactly one covariant index from the collection $\mu_{m=1} \dots \mu_{m=2k}$ is carried by the metric $g_{\mu_1}^{\alpha}$. All other covariant instances of μ_m are carried by (2k-1) instances

of q_{μ_m} . This difference among the lowered indices introduces a complication that was absent in the previous cases but nevertheless allows us to write

$$\Delta \tilde{T}^{A}_{3\mu\nu} = -2i \sum_{k=1}^{\infty} \left[-i\epsilon_{\mu\nu\alpha\beta} q^{\beta} C_{3}^{2k} A_{\tau=2}^{2k} \right] \times \frac{2^{2k}}{(Q^{2})^{2k}} \times g^{\alpha}_{\mu_{1}} \times \left(\prod_{m=2}^{2k} q_{\mu_{m}}\right) \times \tilde{\Pi}^{\mu_{1}\dots\mu_{2k}} + \mathcal{O}(\tau > 2) .$$
(A.75)

In principle, $g_{\mu_1}^{\alpha}$ can contract with a metric from the collection $\{g \dots g\}$ when it contains the index $\mu_{m=1}$. This extracts a factor of q^{α} out of the production factor $(\prod_{m=2}^{2k} q_{\mu_m})$, and hence potential permutations, through the chain $q^{\alpha} = g_{\mu_1}^{\alpha} q_{\mu_n} g^{\mu_n \mu_1}$. However, this generates a term proportional to $\epsilon_{\mu\nu\alpha\beta}q^{\beta}q^{\alpha}$, which is zero due to the antisymmetric tensor.

It is also possible for the metric $g_{\mu_1}^{\alpha}$ to contract with a momentum factor from the collection $\{p_A \dots p_A\}$. This extracts a factor of $p_A^{\alpha} = g_{\mu_1}^{\alpha} p_A^{\mu_1}$, and reduces the possible number of permutations in the momentum factor $\tilde{\Pi}$. The contractions within $\Delta \tilde{T}_{3\mu\nu}^{A}$ then involve (2k-1) instances of q_{μ_m} contracting with (2k-2j-1) instances of $p_A^{\mu_m}$, to make (2k-2j-1) powers of $(q \cdot p_A)$. The remaining (2k-1) - (2k-2j-1) = 2j instances of q_{μ_m} then contract with the remaining j metrics. Finally, since the number of available $p_A^{\mu_m}$ has shifted (by one unit), the multiplicity factor in Eq. (A.64) must be updated accordingly. With the updated η factor, the outcome is

$$g_{\mu_{1}}^{\alpha} \left(\prod_{m=2}^{2k} q_{\mu_{m}}\right) \tilde{\Pi}^{\mu_{1}...\mu_{2k}} = p_{A}^{\alpha} \sum_{j=0}^{k-1} (-1)^{j} \frac{(2k-j)!}{2^{j}(2k)!} \frac{1}{2^{j}} \frac{(2k-1)!}{j!(2k-2j-1)!} \times (q^{2})^{j} (q \cdot p_{A})^{(2k-2j-1)} (p_{A}^{2})^{j}$$
(A.76)

$$= \frac{2p_A^{\alpha}}{Q^2} \frac{(Q^2)^{2k}}{2^{2k}} \sum_{j=0}^{k-1} \frac{(2k-j)!(2k-1)!}{(2k)!j!(2k-2j-1)!} \left(\frac{M_A^2}{Q^2}\right)^j x_A^{-(2k-2j-1)} .$$
(A.77)

Note that the summation over j extends only to j = k - 1 since there are fewer allowed permutations; this can be surmised from the denominator factor 1/(2k - 2j - 1)!, which would reduce to 1/(-1)! were j = k. After assembling the different terms, we obtain

$$\Delta \tilde{T}^{A}_{3\mu\nu} = -i\epsilon_{\mu\nu\alpha\beta} \frac{p^{\alpha}_{A}q^{\beta}}{Q^{2}} \times (-4i) \sum_{k=1}^{\infty} \left[C^{2k}_{3} A^{2k}_{\tau=2} \right] \\ \times \sum_{j=0}^{k-1} \frac{(2k-j)!(2k-1)!}{(2k)!j!(2k-2j-1)!} \left(\frac{M^{2}_{A}}{Q^{2}} \right)^{j} x^{-(2k-2j-1)}_{A} + \mathcal{O}(\tau > 2) .$$
(A.78)

Unlike the other $\Delta \tilde{T}_i^A$ in Eq. (A.69), we can identify the quantity to the right of the antisymmetric tensor $\epsilon_{\mu\nu\alpha\beta}$ as $\Delta \tilde{T}_3^A$ since there is a one-to-one correspondence.

(iv) $\Delta \tilde{T}_{5\mu\nu}^{A}$: The steps to determine $\Delta \tilde{T}_{5\mu\nu}^{A}$ (and $\Delta \tilde{T}_{6\mu\nu}^{A}$) for $M_{A} \neq 0$ closely follow those of $\Delta \tilde{T}_{3\mu\nu}^{A}$ due to the presence of only one metric carrying a lowered index from the collection $\mu_{m=1} \dots \mu_{m=2k}$. The difference, however, is the absence of an antisymmetric tensor. The contractions of $g_{\mu\mu_{1}}$ and $g_{\nu\mu_{1}}$ with the collections $\{g \dots g\}$ and $\{p_{A} \dots p_{A}\}$ will thus generate two sets of terms each.

To obtain both sets of terms, we first write

$$\Delta \tilde{T}_{5\mu\nu}^{A} = -4i \sum_{k=1}^{\infty} \left[(g_{\mu\mu_{1}}q_{\nu} + g_{\nu\mu_{1}}q_{\mu})C_{5}^{2k}A_{\tau=2}^{2k} \right] \times \frac{2^{2k}}{(Q^{2})^{2k}} \times \left(\prod_{m=2}^{2k} q_{\mu_{m}}\right) \times \tilde{\Pi}^{\mu_{1}\dots\mu_{2k}} + \mathcal{O}(\tau > 2) .$$
(A.79)

For the case of the metric $g_{\alpha\mu_1}$ contracting with $\{p_A \dots p_A\}$, we have from Eq. (A.77)

$$g_{\alpha\mu_{1}}\left(\prod_{m=2}^{2k}q_{\mu_{m}}\right)\tilde{\Pi}^{\mu_{1}\dots\mu_{2k}}\bigg|_{\mu_{1}\in\{p_{A}\dots,p_{A}\}} = \frac{2p_{A\alpha}}{Q^{2}} \left.\frac{(Q^{2})^{2k}}{2^{2k}} \sum_{j=0}^{k-1} \frac{(2k-j)!(2k-1)!}{(2k)!j!(2k-2j-1)!} \left(\frac{M_{A}^{2}}{Q^{2}}\right)^{j} x_{A}^{-(2k-2j-1)} \right.$$
(A.80)

For the second case, $g_{\alpha\mu_1}$ contracts with a metric in $\{g \dots g\}$ and extracts a factor of $q_{\alpha} = g_{\alpha\mu_1}q_{\mu_n}g^{\mu_n\mu_1}$ out of the production factor $(\prod_{m=2}^{2k}q_{\mu_m})$. The precise factor of q_{μ_n} that is extracted can be any one of the (2k-1) elements in the product factor, implying an additional multiplicity factor of this size [54]. The remaining (2k-2) instances of q_{μ_m} then contract with (2k-2j) instances of $p_A^{\mu_m}$ to make (2k-2j) powers of $(q \cdot p_A)$. The remaining (2k-2) - (2k-2j) = 2j-2 factors of q_{μ_m} then contract with the 2(j-1) indices of the remaining (j-1) metrics. After updating η and noting that the sum now starts at j = 1, the outcome is

$$g_{\alpha\mu_{1}}\left(\prod_{m=2}^{2k}q_{\mu_{m}}\right)\tilde{\Pi}^{\mu_{1}\dots\mu_{2k}}\Big|_{\mu_{1}\in\{g\dots g\}} = q_{\alpha}\sum_{j=1}^{k}(-1)^{j}\frac{(2k-j)!}{2^{j}(2k)!}\frac{1}{2^{j-1}}\frac{(2k-2)!}{(j-1)!(2k-2j)!} \times (2k-1)$$

$$\times (q^{2})^{(j-1)}(q\cdot p_{A})^{(2k-2j)}(p_{A}^{2})^{j} \tag{A.81}$$

$$= -\frac{2q_{\alpha}}{Q^2} \frac{(Q^2)^{2k}}{2^{2k}} \sum_{j=1}^k \frac{(2k-j)!(2k-1)!}{(2k)!(j-1)!(2k-2j)!} \left(\frac{M_A^2}{Q^2}\right)^j x_A^{-(2k-2j)} .$$
(A.82)

Assembling all terms for both $g_{\mu\mu_1}$ and $g_{\nu\mu_1}$, we obtain the expression

$$\begin{split} \Delta \tilde{T}_{5\mu\nu}^{A} &= \frac{\left(p_{A\mu}q_{\nu} + p_{A\nu}q_{\mu}\right)}{Q^{2}} \\ &\times \left(-8i\right) \sum_{k=1}^{\infty} \left[C_{5}^{2k}A_{\tau=2}^{2k}\right] \sum_{j=0}^{k-1} \frac{\left(2k-j\right)!(2k-1)!}{(2k)!j!(2k-2j-1)!} \left(\frac{M_{A}^{2}}{Q^{2}}\right)^{j} x_{A}^{-(2k-2j-1)} \\ &- \frac{\left(q_{\mu}q_{\nu} + q_{\nu}q_{\mu}\right)}{Q^{2}} \\ &\times \left(-8i\right) \sum_{k=1}^{\infty} \left[C_{5}^{2k}A_{\tau=2}^{2k}\right] \sum_{j=1}^{k} \frac{\left(2k-j\right)!(2k-1)!}{\left(2k\right)!(j-1)!(2k-2j)!} \left(\frac{M_{A}^{2}}{Q^{2}}\right)^{j} x_{A}^{-(2k-2j)} \\ &+ \mathcal{O}(\tau > 2) \;. \end{split}$$
(A.83)

Importantly, there are two tensor structures: The first is $(p_{A\mu}q_{\nu} + p_{A\nu}q_{\mu})$, whose coefficient contributes to $\Delta \tilde{T}_5^A$ in Eq. (A.69). The second is $q_{\mu}q_{\nu}$, whose coefficient contributes to $\Delta \tilde{T}_4^A$ in Eq. (A.69). Note that the second term can equally be written as $4(q_{\mu}q_{\nu})$; we do not (yet) condense the result in order to emphasize that $g_{\mu\mu_1}$ and $g_{\nu\mu_1}$ each generates two terms.

(v) $\Delta \tilde{T}^{A}_{6\mu\nu}$: The next term, $\Delta \tilde{T}^{A}_{6\mu\nu}$ for $M_A \neq 0$, can be obtained directly from $\Delta \tilde{T}^{A}_{5\mu\nu}$ by making the replacement $q_{\mu} \rightarrow -q_{\mu}$ and $C_5 \rightarrow C_6$. When making this replacement, the second part of Eq. (A.83) vanishes since $(q_{\mu}q_{\nu} + q_{\nu}q_{\mu}) \rightarrow (q_{\mu}q_{\nu} - q_{\nu}q_{\mu}) = 0$. The final result for $\Delta \tilde{T}^{A}_{6\mu\nu}$ is

$$\Delta \tilde{T}_{6\mu\nu}^{A} = \frac{\left(p_{A\mu}q_{\nu} - p_{A\nu}q_{\mu}\right)}{Q^{2}} \times (-8i) \times \sum_{k=1}^{\infty} \left[C_{6}^{2k}A_{\tau=2}^{2k}\right] \\ \times \sum_{j=0}^{k-1} \frac{(2k-j)!(2k-1)!}{(2k)!j!(2k-2j-1)!} \left(\frac{M_{A}^{2}}{Q^{2}}\right)^{j} x_{A}^{-(2k-2j-1)} + \mathcal{O}(\tau > 2) .$$
(A.84)

(vi) $\Delta \tilde{T}^A_{2\mu\nu}$: The final term $\Delta \tilde{T}^A_{2\mu\nu}$ for $M_A \neq 0$, which is given by

$$\Delta \tilde{T}^{A}_{2\mu\nu} = (-2i) \sum_{k=2}^{\infty} \left[Q^2 C_2^{2k} A_{\tau=2}^{2k} \right] \times \frac{2^{2k}}{(Q^2)^{2k}} \times g_{\mu\mu_1} g_{\nu\mu_2} \times \left(\prod_{m=3}^{2k} q_{\mu_m} \right) \times \tilde{\Pi}^{\mu_1 \dots \mu_{2k}} + \mathcal{O}(\tau > 2) .$$
(A.85)

is the most nuanced because the covariant tensor structure involves two metrics. That is to say, two lowered indices from the collection $\mu_{m=1} \dots \mu_{m=2k}$ are carried by $g_{\mu\mu_1}$ and $g_{\nu\mu_2}$. The contraction of these metrics with the collections $\{g \dots g\}$ and $\{p_A \dots p_A\}$, in turn, generates five terms with four distinct Lorentz structures. The five possible ways are:

- (a) both $g_{\mu\mu_1}$ and $g_{\nu\mu_2}$ contract with momenta from $\{p_A \dots p_A\}$ and extract $p_{A\mu}p_{A\nu}$;
- (b) both $g_{\mu\mu_1}$ and $g_{\nu\mu_2}$ contract with *different* metrics from $\{g \dots g\}$ and extract the quantity $q_{\mu}q_{\nu}$ from the production factor $(\prod_{m=3}^{2k} q_{\mu_m})$;
- (c) both $g_{\mu\mu_1}$ and $g_{\nu\mu_2}$ contract with the same metric from $\{g \dots g\}$ and extract $g_{\mu\nu}$;
- (d) $g_{\mu\mu_1}$ contracts with $\{p_A \dots p_A\}$ and $g_{\nu\mu_2}$ with $\{g \dots g\}$ to extract $(p_{A\mu}q_{\nu})$;
- (e) same as (d) but with the index exchange $\mu \leftrightarrow \nu$ to extract the quantity $(p_{A\nu}q_{\mu})$.

In case (a), contracting the metrics with the collection $\{p_A \dots p_A\}$ leaves (2k - 2j - 2) instances of $p_A^{\mu_m}$ to contract with (2k - 2) instances of q_{μ_m} . The remaining (2k - 2) - (2k - 2j - 2) = 2jinstances of q_{μ_m} then contracts with the 2j indices of the j metrics in the collection $\{g \dots g\}$. Updating the permutation factor, we get

$$g_{\mu\mu_1}g_{\nu\mu_2}\left(\prod_{m=3}^{2k}q_{\mu_m}\right)\tilde{\Pi}^{\mu_1\dots\mu_{2k}}\Big|_{\mu_1,\mu_2\in\{p_A\dots p_A\}} = p_{A\mu}p_{A\nu}\sum_{j=0}^{k-1}\frac{(2k-j)!}{2^j(2k)!}\frac{1}{2^j}\frac{(2k-2)!}{j!(2k-2j-2)!}$$

$$\times (-1)^j(q^2)^j \ (q\cdot p_A)^{(2k-2j-2)} \ (p_A^2)^j \tag{A.86}$$

$$=\frac{4p_{A\mu}p_{A\nu}}{Q^4} \frac{(Q^2)^{2k}}{2^{2k}} \sum_{j=0}^{k-1} \frac{(2k-j)!(2k-2)!}{(2k)!j!(2k-2j-2)!} \left(\frac{M_A^2}{Q^2}\right)^j x_A^{-(2k-2j-2)} .$$
(A.87)

Note that the summation over j extends only to j = k - 1 since there are fewer allowed permutations; this can be inferred from the denominator factor 1/(2k - 2j - 2)!.

In case (b), contracting $g_{\mu\mu_1}$ and $g_{\nu\mu_2}$ with different metrics from the collection $\{g \ldots g\}$ leaves (2k-4) instances of q_{μ_m} to contract with (2k-2j) instances of $p_A^{\mu_m}$. The precise q_{μ_m} that are extracted from the product factor $(\prod_{m=3}^{2k} q_{\mu_m})$ can be any two of its (2k-2) elements, implying the additional multiplicity factor $(2k-2) \times (2k-3)$. The remaining (2k-4) - (2k-2j) = (2j-4)

factors of q_{μ_m} then contract with the (j-2) metrics remaining in $\{g \dots g\}$. Updating the η factor and noting the summation over j starts at j = 2, we get

$$g_{\mu\mu_{1}}g_{\nu\mu_{2}}\left(\prod_{m=3}^{2k}q_{\mu_{m}}\right)\tilde{\Pi}^{\mu_{1}...\mu_{2k}}\Big|_{\mu_{1},\mu_{2}\in\{g...g\}}$$

$$=q_{\mu}q_{\nu}\sum_{j=2}^{k}\frac{(2k-j)!}{2^{j}(2k)!}\frac{1}{2^{j-2}}\frac{(2k-4)!}{(j-2)!(2k-2j)!} \times (2k-3)(2k-2)$$

$$\times (-1)^{j}(q^{2})^{(j-2)}(q\cdot p_{A})^{(2k-2j)}(p_{A}^{2})^{j} \qquad (A.88)$$

$$=\frac{4q_{\mu}q_{\nu}}{Q^4} \frac{(Q^2)^{2k}}{2^{2k}} \sum_{j=2}^k \frac{(2k-j)!(2k-2)!}{(2k)!(j-2)!(2k-2j)!} \left(\frac{M_A^2}{Q^2}\right)^j x_A^{-(2k-2j)}.$$
 (A.89)

In case (c), contracting $g_{\mu\mu_1}$ and $g_{\nu\mu_2}$ with the same metric from the collection $\{g \dots g\}$ leaves (2k-2) instances of q_{μ_m} to contract with all (2k-2j) instances of $p_A^{\mu_m}$. The (2k-2) - (2k-2j) = (2j-2) uncontracted instances of q_{μ_m} are then matched with the (j-1) metrics that are left in $\{g \dots g\}$. Updating η and the summation over j, which now starts from j = 1, we get

$$g_{\mu\mu_{1}}g_{\nu\mu_{2}} \left(\prod_{m=3}^{2k} q_{\mu_{m}}\right) \tilde{\Pi}^{\mu_{1}\dots\mu_{2k}} \bigg|_{g^{\mu_{1}\mu_{2}} \in \{g...g\}} = g_{\mu\nu} \sum_{j=1}^{k} \frac{(2k-j)!}{2^{j}(2k)!} \times \frac{1}{2^{j-1}} \frac{(2k-2)!}{(j-1)!(2k-2j)!} (-1)^{j} (q^{2})^{(j-1)} (q \cdot p_{A})^{(2k-2j)} (p_{A}^{2})^{j}$$
(A.90)

$$= -\frac{2g_{\mu\nu}}{Q^2} \frac{(Q^2)^{2k}}{2^{2k}} \sum_{j=1}^k \frac{(2k-j)!(2k-2)!}{(2k)!(j-1)!(2k-2j)!} \left(\frac{M_A^2}{Q^2}\right)^j x_A^{-(2k-2j)}.$$
 (A.91)

In case (d), $g_{\mu\mu_1}$ contracts with $\{p_A \dots p_A\}$, leaving (2k - 2j - 1) instances of $p_A^{\mu_m}$, while $g_{\nu\mu_2}$ contracts with $\{g \dots g\}$, and eventually $(\prod_{\mu=3}^{2k} q_{\mu_n})$, leaving a total of (2k - 3) instances of q_{μ_m} . The result is (2k - 2j - 1) powers of $(q \cdot p_A)$ and (2k - 3) - (2k - 2j - 1) = (2j - 2) factors of q_{μ_m} to contract with (j - 1) metrics. The q_{μ_m} that is extracted from the product factor can be any one of its initial (2k - 2) elements, implying an additional multiplicity factor of this size. Updating η and noting the summation over j runs from j = 1 to j = k - 1, we obtain

$$g_{\mu\mu_{1}}g_{\nu\mu_{2}}\left(\prod_{m=3}^{2k}q_{\mu_{m}}\right)\tilde{\Pi}^{\mu_{1}...\mu_{2k}}\Big|_{\mu_{2}\in\{g...g\}}^{\mu_{1}\in\{p_{A}...p_{A}\}} = p_{A\mu}q_{\nu}\sum_{j=1}^{k-1}\frac{(2k-j)!}{2^{j}(2k)!}$$

$$\times \frac{1}{2^{j-1}}\frac{(2k-3)!}{(j-1)!(2k-2j-1)!} \times (2k-2) \ (-1)^{j}(q^{2})^{(j-1)} \ (q\cdot p_{A})^{(2k-2j-1)} \ (p_{A}^{2})^{j}$$
(A.92)

$$= -\frac{4p_{A\mu}q_{\nu}}{Q^4} \frac{(Q^2)^{2k}}{2^{2k}} \sum_{j=1}^{k-1} \frac{(2k-j)!(2k-2)!}{(2k)!(j-1)!(2k-2j-1)!} \left(\frac{M_A^2}{Q^2}\right)^j x_A^{-(2k-2j-1)} .$$
(A.93)

Case (e) follows from case (d) but with $\mu \leftrightarrow \nu$ exchanged.

After assembling all five cases and regrouping, we find

$$\begin{split} \Delta \tilde{T}_{2\mu\nu}^{A} &= \frac{2p_{A\mu}p_{A\nu}}{Q^{2}} \times (-4i) \sum_{k=1}^{\infty} \left[C_{2}^{2k} A_{\tau=2}^{2k} \right] \\ &\times \sum_{j=0}^{k-1} \frac{(2k-j)!(2k-2)!}{(2k)!j!(2k-2j-2)!} \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} x_{A}^{-(2k-2j-2)} \\ &-g_{\mu\nu} \times (-4i) \sum_{k=1}^{\infty} \left[C_{2}^{2k} A_{\tau=2}^{2k} \right] \sum_{j=1}^{k} \frac{(2k-j)!(2k-2)!}{(2k)!(j-1)!(2k-2j)!} \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} x_{A}^{-(2k-2j)} \\ &+ \frac{2q_{\mu}q_{\nu}}{Q^{2}} \times (-4i) \sum_{k=2}^{\infty} \left[C_{2}^{2k} A_{\tau=2}^{2k} \right] \sum_{j=2}^{k} \frac{(2k-j)!(2k-2)!}{(2k)!(j-2)!(2k-2j)!} \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} x_{A}^{-(2k-2j)} \\ &- \frac{2(p_{A\mu}q_{\nu} + p_{A\nu}q_{\mu})}{Q^{2}} \times (-4i) \sum_{k=2}^{\infty} \left[C_{2}^{2k} A_{\tau=2}^{2k} \right] \\ &\times \sum_{j=1}^{k-1} \frac{(2k-j)!(2k-2)!}{(2k)!(j-1)!(2k-2j-1)!} \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} x_{A}^{-(2k-2j-1)} + \mathcal{O}(\tau > 2) . \end{split}$$
(A.94)

The result implies that $\Delta \tilde{T}_{2\mu\nu}^{A}$, and hence the Wilson coefficient C_{2}^{2k} , contributes to ΔT_{1}^{A} , ΔT_{2}^{A} , ΔT_{4}^{A} , and ΔT_{5}^{A} in Eq. (A.69). The absence of contributions to ΔT_{3}^{A} and ΔT_{6}^{A} , which are asymmetric under $\mu \leftrightarrow \nu$ exchange, can be traced to the tensor coefficient multiplying C_{2}^{2k} in $c_{\mu\nu\mu_{1},\ldots,\mu_{2k}}^{\tau=2,\iota}$, which is symmetric under $\mu \leftrightarrow \nu$ exchange.

Appendix A.4. Nuclear structure functions from the OPE II: structure-function mixing Using the expressions for $\Delta \tilde{T}^A_{\iota \ \mu\nu}$, we finally obtain the quantities $\Delta \tilde{T}^A_i$ in Eq. (A.69). For $\Delta \tilde{T}^A_1$ and $M_A \neq 0$, we collect like-terms from $\Delta \tilde{T}^A_{1\mu\nu}$ and $\Delta \tilde{T}^A_{2\mu\nu}$ to obtain

$$\begin{split} \Delta \tilde{T}_{1}^{A} &= (-4i) \sum_{k=1}^{\infty} \left[C_{1}^{2k} A_{\tau=2}^{2k} \right] \sum_{j=0}^{k} \frac{(2k-j)!(2k)!}{(2k)!j!(2k-2j)!} \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} x_{A}^{-(2k-2j)} \\ &+ (-4i) \sum_{k=1}^{\infty} \left[C_{2}^{2k} A_{\tau=2}^{2k} \right] \sum_{j=1}^{k} \frac{(2k-j)!(2k-2)!}{(2k)!(j-1)!(2k-2j)!} \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} x_{A}^{-(2k-2j)} + \mathcal{O}(\tau > 2) \quad (A.95) \\ &= (-4i) \sum_{l=0}^{\infty} x_{A}^{-(2l+1)} \sum_{j=0}^{\infty} \frac{(2l+j+1)!}{j!(2l+1)!} \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} \\ &\times \left[\left(C_{1}^{2l+2j+1} A_{\tau=2}^{2l+2j+1} \right) + \frac{j \left(C_{2}^{2l+2j+1} A_{\tau=2}^{2l+2j+1} \right)}{(2l+2j+1)(2l+2j)} \right] + \mathcal{O}(\tau > 2) \quad (A.96) \end{split}$$

To reach Eq. (A.96), a factor of 1 = (2k)(2k-1)j/(2k)(2k-1)j is introduced into the C_2 term to complete the factorials. The summation in the C_2 term can then be extended to include j = 0 since the contribution is zero. We also re-index the summation over k into a summation over 2l = 2k-2j-1. After this shift it becomes manifest that the contribution from the C_2 Wilson coefficient vanishes when the summation over j is truncated at j = 0. It also becomes clear that structure-function mixing is due to TMCs since neglecting terms where j > 0 is equivalent to taking the $(M_A^2/Q^2) \rightarrow 0$ limit.
The coefficient $\Delta \tilde{T}_2^A$ is sourced entirely from $\Delta \tilde{T}_{\mu\nu2}^A$, and is given by

$$\begin{split} \Delta \tilde{T}_{2}^{A} &= (-4i) \; \frac{2M_{A}^{2}}{Q^{2}} \; \sum_{k=1}^{\infty} \left[C_{2}^{2k} A_{\tau=2}^{2k} \right] \; \sum_{j=0}^{k-1} \; \frac{(2k-j)!(2k-2)!}{(2k)!j!(2k-2j-2)!} \; \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} \; x_{A}^{-(2k-2j-2)} \\ &+ \mathcal{O}(\tau > 2) \\ &= (-4i) \; \frac{2M_{A}^{2}}{Q^{2}} \; \sum_{l=0}^{\infty} \; x_{A}^{-(2l-1)} \; \sum_{j=0}^{\infty} \; \frac{(2l+j+1)!}{j!(2l+1)!} \; \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} (2l+1)(2l) \; \frac{\left(C_{2}^{(2l+2j+1)} A_{\tau=2}^{(2l+2j+1)} \right)}{(2l+2j+1)(2l+2j)} \\ &+ \mathcal{O}(\tau > 2) \; . \end{split}$$
(A.97)

In reaching Eq. (A.98) we again completed the factorials and set 2l = 2k - 2j - 1.

Similarly, $\Delta \tilde{T}_3^A$ for $M_A \neq 0$ is sourced entirely from $\Delta \tilde{T}_{\mu\nu3}^A$. It is given by

$$\Delta \tilde{T}_{3}^{A} = (-4i) \frac{M_{A}^{2}}{Q^{2}} \sum_{k=1}^{\infty} \left[C_{3}^{2k} A_{\tau=2}^{2k} \right] \sum_{j=0}^{k-1} \frac{(2k-j)!(2k-1)!}{(2k)!j!(2k-2j-1)!} \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} x_{A}^{-(2k-2j-1)} + \mathcal{O}(\tau > 2)$$

$$= (-4i) \frac{M_{A}^{2}}{Q^{2}} \sum_{l=0}^{\infty} x_{A}^{-(2l-1)} \sum_{j=0}^{\infty} \frac{(2l+j)!}{j!(2l)!} \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} (2l) \frac{\left(C_{3}^{(2l+2j)} A_{\tau=2}^{(2l+2j)} \right)}{(2l+2j)} + \mathcal{O}(\tau > 2) , \qquad (A.100)$$

where the index l = k - j was used after completing the factorials.

For $\Delta \tilde{T}_4^A$ and $M_A \neq 0$, we collect terms from $\Delta \tilde{T}_{\mu\nu4}^A$, $\Delta \tilde{T}_{\mu\nu2}^A$, and $\Delta \tilde{T}_{\mu\nu6}^A$. The result is

$$\begin{split} \Delta \tilde{T}_{4}^{A} &= (-4i) \; \frac{2M_{A}^{2}}{Q^{2}} \; \sum_{k=1}^{\infty} \; \left[C_{4}^{2k} A_{\tau=2}^{2k} \right] \; \sum_{j=0}^{k} \; \frac{(2k-j)!(2k)!}{(2k)!j!(2k-2j)!} \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} \; x_{A}^{-(2k-2j)} \; . \\ + (-4i) \; \frac{2M_{A}^{2}}{Q^{2}} \; \sum_{k=2}^{\infty} \left[C_{2}^{2k} A_{\tau=2}^{2k} \right] \; \sum_{j=2}^{k} \; \frac{(2k-j)!(2k-2)!}{(2k)!(j-2)!(2k-2j)!} \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} \; x_{A}^{-(2k-2j)} \\ - (-4i) \; \frac{4M_{A}^{2}}{Q^{2}} \; \sum_{k=1}^{\infty} \; \left[C_{5}^{2k} A_{\tau=2}^{2k} \right] \; \sum_{j=1}^{k} \; \frac{(2k-j)!(2k-1)!}{(2k)!(j-1)!(2k-2j)!} \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} \; x_{A}^{-(2k-2j)} \\ + \mathcal{O}(\tau > 2) \; (A.101) \\ = (-4i) \; \frac{2M_{A}^{2}}{Q^{2}} \; \sum_{l=0}^{\infty} x_{A}^{-(2l+1)} \; \sum_{j=0}^{\infty} \frac{(2l+j+1)!}{j!(2l+1)!} \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} \\ \times \left[\left(C_{4}^{(2l+2j+1)} A_{\tau=2}^{(2l+2j+1)} \right) + \; \frac{j \; (j-1) \; \left(C_{2}^{(2l+2j+1)} A_{\tau=2}^{(2l+2j+1)} \right)}{(2l+2j+1)(2l+2j)} - \frac{2j \; \left(C_{5}^{(2l+2j+1)} A_{\tau=2}^{(2l+2j+1)} \right)}{(2l+2j+1)} \right] \\ + \mathcal{O}(\tau > 2) \; . \end{split}$$
(A.102)

We complete the factorials using factors of 1 = (2k)j/(2j)j, etc., and use 2l = 2k-2j-1 to re-index the summation over k. The summations over j can also be extended to j = 0 since the additional terms are proportional to j or j(j-1), i.e., are zero at j = 0 and/or j = 1, and hence do not contribute.

The coefficient $\Delta \tilde{T}_5^A$ for $M_A \neq 0$ is generated by $\Delta \tilde{T}_{2\mu\nu}^A$ and $\Delta \tilde{T}_{5\mu\nu}^A$. The result is

$$\begin{split} \Delta \tilde{T}_{5}^{A} &= (-4i) \; \frac{2M_{A}^{2}}{Q^{2}} \; \sum_{k=1}^{\infty} \; \left[C_{5}^{2k} A_{\tau=2}^{2k} \right] \; \sum_{j=0}^{k-1} \; \frac{(2k-j)!(2k-1)!}{(2k)!j!(2k-2j-1)!} \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} \; x_{A}^{-(2k-2j-1)} \\ &- (-4i) \; \frac{2M_{A}^{2}}{Q^{2}} \; \sum_{k=2}^{\infty} \left[C_{2}^{2k} A_{\tau=2}^{2k} \right] \; \sum_{j=1}^{k-1} \; \frac{(2k-j)!(2k-2)!}{(2k)!(j-1)!(2k-2j-1)!} \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} \; x_{A}^{-(2k-2j-1)} \\ &+ \; \mathcal{O}(\tau > 2) \qquad (A.103) \\ &= (-4i) \; \frac{2M_{A}^{2}}{Q^{2}} \; \sum_{l=0}^{\infty} x_{A}^{-2l} \; \sum_{j=0}^{\infty} \frac{(2l+j+1)!}{j!(2l+1)!} \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} \\ &\times \left[\frac{(2l+1) \; \left(C_{5}^{(2l+2j+1)} A_{\tau=2}^{(2l+2j+1)} \right)}{(2l+2j+1)} \; - \frac{j \; (2l+1) \; \left(C_{2}^{(2l+2j+1)} A_{\tau=2}^{(2l+2j+1)} \right)}{(2l+2j+1)(2l+2j)} \right] \\ &+ \mathcal{O}(\tau > 2) \; , \qquad (A.104) \end{split}$$

where the index reassignment is 2l = 2k - 2j - 1.

Finally, the coefficient $\Delta \tilde{T}_6^A$ is generated solely by $\Delta \tilde{T}_{6\mu\nu}^A$. It is given by

$$\begin{split} \Delta \tilde{T}_{6}^{A} = (-4i) \; \frac{2M_{A}^{2}}{Q^{2}} \; \sum_{k=1}^{\infty} \; \left[C_{6}^{2k} A_{\tau=2}^{2k} \right] \; \sum_{j=0}^{k-1} \; \frac{(2k-j)!(2k-1)!}{(2k)!j!(2k-2j-1)!} \; \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} \; x_{A}^{-(2k-2j-1)} \\ &+ \; \mathcal{O}(\tau > 2) \\ = (-4i) \; \frac{2M_{A}^{2}}{Q^{2}} \; \sum_{l=0}^{\infty} \; x_{A}^{-(2l-1)} \; \sum_{j=0}^{\infty} \; \frac{(2l+j)!}{j!(2l)!} \; \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} \; (2l) \; \frac{\left(C_{6}^{(2l+2j)} A_{\tau=2}^{(2l+2j)} \right)}{(2l+2j)} \\ &+ \; \mathcal{O}(\tau > 2) \; , \end{split}$$
(A.105)

where the index l = k - j was used after completing the factorials.

We briefly stress that the (M_A^2/Q^2) prefactor in all $\Delta \tilde{T}_i^A$ is conventional. They exist to match the normalization of \tilde{T}_i^A in Eq. (A.69) and structure functions \tilde{W}_i^A in Eq. (A.31). The massless target limit should be understood as truncating the summation over j at j = 0. In addition, the re-indexing from k to l in the above expressions is not random. Different choices of relabeling can be found in the literature and depend on the precise interpretation of the quantities to the right of the x_A^{-N} factor. Our choices are motivated by the attempt to align the ratio of factorials into binomial distributions, e.g., $\binom{k+j}{i} = (k+j)!/j!k!$.

Appendix A.5. Nuclear structure functions from the OPE III: massless and massive targets

Massless nuclear targets

As a final preliminary step to obtaining TMCs at leading twist, we establish the connection between the effective Wilson coefficients C_{ι} , the reduced matrix element $A_{\tau=2}$, and structure functions in the limit that $(M_A^2/Q^2) \rightarrow 0$; in this limit, the TMCs will not be present, and we label this result as "No TMC" in Eq.(A.107). For each $\Delta \tilde{T}_i^A$ above, truncating terms with j > 0 and using the Taylor expansion of $\Delta \tilde{T}_i^A$ from Eq. (A.61) gives the following:

$$\Delta \tilde{T}_{1}^{A} \bigg|_{j=0} = -4i \sum_{l=0}^{\infty} \left. x_{A}^{-(2l+1)} \left(C_{1}^{2l+1} A_{\tau=2}^{2l+1} \right) + \mathcal{O}(\tau > 2) \right. = \sum_{l=0}^{\infty} \left. \omega_{A}^{2l+1} \tilde{F}_{1}^{A(2l+1)} \right|_{\text{No TMC}}, \quad (A.107a)$$

$$\left(\frac{Q^2}{2x_A M_A^2}\right) \left. \Delta \tilde{T}_2^A \right|_{j=0} = -4i \sum_{l=0}^{\infty} \left. x_A^{-2l} \left(C_2^{2l+1} A_{\tau=2}^{2l+1} \right) \right. + \mathcal{O}(\tau > 2) = \sum_{l=0}^{\infty} \left. \omega_A^{2l} \left. \tilde{F}_2^{A(2l)} \right|_{\text{No TMC}}, \tag{A.107b}$$

$$\left(\frac{Q^2}{x_A M_A^2}\right) \left. \Delta \tilde{T}_3^A \right|_{j=0} = -4i \sum_{l=0}^{\infty} \left. x_A^{-2l} \left(C_3^{2l} A_{\tau=2}^{2l} \right) + \mathcal{O}(\tau > 2) = \sum_{l=0}^{\infty} \left. \omega_A^{2l} \left. \tilde{F}_3^{A(2l)} \right|_{\text{No TMC}},$$
(A.107c)

$$\left(\frac{Q^2}{2M_A^2}\right)\Delta \tilde{T}_4^A \bigg|_{j=0} = -4i\sum_{l=0}^{\infty} \left. x_A^{-(2l+1)} \left(C_4^{2l+1} A_{\tau=2}^{2l+1} \right) + \mathcal{O}(\tau > 2) \right. \\ = \sum_{l=0}^{\infty} \left. \omega_A^{2l+1} \tilde{F}_4^{A(2l+1)} \right|_{\text{No TMC}}, \quad (A.107d)$$

$$\left(\frac{Q^2}{2x_A M_A^2}\right) \left. \Delta \tilde{T}_5^A \right|_{j=0} = -4i \sum_{l=0}^{\infty} \left. x_A^{-(2l+1)} \left(C_5^{2l+1} A_{\tau=2}^{2l+1} \right) + \mathcal{O}(\tau > 2) \right|_{l=0} \left. \omega_A^{2l+1} \tilde{F}_5^{A(2l+1)} \right|_{\text{No TMC}}, \quad (A.107e)$$

$$\left(\frac{Q^2}{2x_A M_A^2}\right) \left. \Delta \tilde{T}_6^A \right|_{j=0} = -4i \sum_{l=0}^{\infty} \left. x_A^{-2l} \left(C_6^{2l} A_{\tau=2}^{2l} \right) \right. + \mathcal{O}(\tau > 2) = \sum_{l=0}^{\infty} \left. \omega_A^{2l} \left. \tilde{F}_6^{A(2l)} \right|_{\text{No TMC}} \right.$$
(A.107f)

The conclusion is that, in the massless target limit, one can identify the leading power of the product of the N^{th} Wilson coefficient C_i^N (note the index shift from ι to i) and hadronic matrix element $A_{\tau=2}^N$ as a Mellin moment of structure functions themselves. This is summarized by

$$\tilde{F}_{i}^{AN}\Big|_{\text{No TMC}} = C_{i}^{N} A_{\tau=2}^{N} + \mathcal{O}(\tau > 2) \quad \text{for} \quad i = 1, 3 - 6 , \qquad (A.108a)$$

$$\tilde{F}_{2}^{A(N-1)} \bigg|_{\text{No TMC}} = C_{2}^{N} A_{\tau=2}^{N} + \mathcal{O}(\tau > 2) .$$
(A.108b)

Massive nuclear targets

Given the relationship between the moments of $\tilde{F}_i^A|_{\text{No TMC}}$ and the product $(C_i A_{\tau=2})$ in the massless target limit, we are now in position to bootstrap expressions for \tilde{F}_i^A with target mass corrections. The key to this is comparing the expansion of $\Delta \tilde{T}_i^A$ in Eq. (A.61) with the full expressions for \tilde{T}_i^A in Eqs. (A.96)-(A.106). One can identify the moments of structure functions with TMCs as everything encapsulated by the summation over j, i.e., the coefficients multiplying the $x_A^{-2l(\pm 1)}$ factor. Using Eq. (A.108), the unknown/non-perturbative ($C_i A_{\tau=2}$) can then be replaced by the measurable $\tilde{F}_i^A|_{\text{No TMC}}$. Expressions in x_A -space are obtained by inverse Mellin transformations. The transformations and summations over j are solvable using various identities.

 \tilde{F}_2^A : To build \tilde{F}_2^A for a non-zero target mass, we start from $\Delta \tilde{T}_2^A$ in Eq. (A.98). Using Eq. (A.61) (or Eq. (A.107)), we identify everything to the right of $x_A^{-(2l-1)}$ in Eq. (A.98) as the $(2l)^{th}$ moment of $\tilde{F}_2^A(x_A, Q^2)$ with TMCs. This allows us to write

$$\tilde{F}_{2}^{A(2l)} = \sum_{j=0}^{\infty} \frac{(2l+j+1)!}{j!(2l+1)!} \left(\frac{M_{A}^{2}}{Q^{2}}\right)^{j} (2l+1)(2l) \frac{\left(C_{2}^{(2l+2j+1)}A_{\tau=2}^{(2l+2j+1)}\right)}{(2l+2j+1)(2l+2j)} .$$
(A.109)

For an arbitrary integrable function B(y) over $y \in [0,1]$ and $m \ge 0$, we have the identities

$$\frac{1}{(m+1)} \int_0^1 dy \ y^{m+1} \ B(y) = \int_0^1 dy \ y^m \ H(y), \tag{A.110a}$$

$$\frac{1}{(m+2)(m+1)} \int_0^1 dy \ y^{(m+2)} \ B(y) = \int_0^1 dy \ y^m \ G(y), \text{ where}$$
(A.110b)

$$H(y) = \int_{y}^{1} dy' B(y'),$$
 and (A.110c)

$$G(y) = \int_{y}^{1} dy' \int_{y'}^{1} dy'' B(y'') = \int_{y}^{1} dy' (y' - y) B(y')$$
(A.110d)

These identities will be used repeatedly for the different structure functions. Schematically, the identities show that each factor of "1/m," which originates from completing the (numerator) factorials in $\Delta \tilde{T}_i^A$, corresponds to an additional integral over a structure function. Using the second identity and Eq. (A.108), we have

$$\frac{C_2^{(2l+2j+1)}A_{\tau=2}^{(2l+2j+1)}A_{\tau=2}^{(2l+2j+1)}}{(2l+2j+1)(2l+2j)} = \frac{1}{(2l+2j+1)(2l+2j)} \int_0^1 dy \ y^{2l+2j+1} \cdot y^{-2}\tilde{F}_2^A(y) \bigg|_{\text{No TMCs}}$$
(A.111)

$$= \int_0^1 dy \ y^{2l+2j-1} \ \tilde{g}_2(y), \quad \text{where}$$
 (A.112)

$$\tilde{g}_2(y) \equiv \int_y^1 dy' \int_{y'}^1 dy'' (y'')^{-2} \tilde{F}_2^A(y'') \bigg|_{\text{No TMCs}} .$$
(A.113)

This allows allows us to evaluate the summation over j:

$$\tilde{F}_{2}^{A(2l)} = \int_{0}^{1} dy \ y^{2l-1} \ \tilde{g}_{2}(y) \ (2l+1)(2l) \ \sum_{j=0}^{\infty} \ \frac{(2l+j+1)!}{j!(2l+1)!} \ \left(\frac{y^{2}M_{A}^{2}}{Q^{2}}\right)^{j}$$
(A.114)

$$= \int_0^1 dy \ y^{-1} \ \tilde{g}_2(y) \ \frac{(2l+1)(2l) \ y^{2l}}{(1-y^2 M_A^2/Q^2)^{2l+2}} \ , \tag{A.115}$$

where we used the first of the following identities, which also hold for z > 1:

$$\sum_{j=0}^{\infty} \frac{(n+j)!}{j! n!} z^j = \frac{1}{(1-z)^{(n+1)}} , \qquad (A.116a)$$

$$\sum_{j=1}^{\infty} j \frac{(n+j)!}{j! n!} z^j = \frac{(n+1) z}{(1-z)^{(n+2)}} , \qquad (A.116b)$$

$$\sum_{j=2}^{\infty} j (j-1) \frac{(n+j)!}{j! n!} z^j = \frac{(n+1) (n+2) z^2}{(1-z)^{(n+3)}} .$$
(A.116c)

The first identity is derived by Taylor expanding the right-hand side and completing the factorial. The second follows from taking the derivative of the first identity and multiplying by z; similarly, the third by taking two derivatives and scaling by z^2 . After relabeling N = 2l, and using the second of the following identities

$$(n+a)z^{-n} = -z^{(1+a)} \frac{d}{dz} z^{-(n+a)}$$
, (A.117a)

$$(n+a)(n+a+1)z^{-n} = z^{(2+a)}\frac{d^2}{dz^2} z^{-(n+a)}$$
, (A.117b)

we can take the inverse Mellin transformation of \tilde{F}_2^{AN} . We remark that these derivative identities will be used repeatedly for the different structure functions. Schematically, they show that each factor of "n" or "(n + 1)," which originates from completing the factorials in $\Delta \tilde{T}_i^A$, corresponds to a derivative over the generating functions G(y) and H(y).

Inserting the derivative identity and taking the inverse Mellin transformation gives

$$\tilde{F}_{2}^{A}(x_{A}) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} dN \ x_{A}^{-N} \ \tilde{F}_{2}^{AN}$$
(A.118)

$$=\frac{x_A^2}{2\pi i}\frac{d^2}{dx_A^2}\int_0^1 dy \;\frac{\tilde{g}_2(y)}{y(1-y^2M_A^2/Q^2)^2}\;\int_{c-i\infty}^{c+i\infty} dN\;\frac{(y/x_A)^N}{(1-y^2M_A^2/Q^2)^N}\tag{A.119}$$

$$= x_A^2 \frac{d^2}{dx_A^2} \int_0^1 dy \; \frac{\tilde{g}_2(y)}{y(1 - y^2 M_A^2/Q^2)^2} \; \delta \left[\log \left(\frac{(y/x_A)}{(1 - y^2 M_A^2/Q^2)} \right) \right] \;, \tag{A.120}$$

where we employed the contour integral

$$\int_{c-i\infty}^{c+i\infty} dN \ z^N = (2\pi i) \ \delta\left[\log(z)\right] \ . \tag{A.121}$$

To evaluate the δ function, we use the decomposition formula

$$\delta[g(z)] = \sum_{z_k} \frac{1}{|g'(z)|} \,\delta(z - z_k), \quad \text{such that} \quad g(z = z_k) = 0. \tag{A.122}$$

This allows us to express the δ function in $\tilde{F}_2^A(x_A)$ in terms of the zeros of the logarithm, i.e., when $(y/x) = (1 - y^2 M_A^2/Q^2)$. While the logarithm vanishes for two values of y, only one is physical. The unphysical solution y_2 lies outside the domain $y \in [0, 1]$, meaning the contribution from $\delta(y - y_2)$ is always zero. The physical solution is the Nachtmann variable [67]:

$$\xi_A = \frac{2x_A}{1+r} , \quad \text{where} \quad r \equiv \sqrt{1 + 4x_A^2 M_A^2/Q^2} = \frac{(1 + \xi_A^2 M_A^2/Q^2)}{(1 - \xi_A^2 M_A^2/Q^2)} , \quad (A.123)$$

and corresponds to the following solution for the δ function:

$$\delta \left[\log \left(\frac{(y/x_A)}{(1 - y^2 M_A^2/Q^2)} \right) \right] = \left[\frac{y(1 - y^2 M_A^2/Q^2)}{(1 + y^2 M_A^2/Q^2)} \right] \delta(\xi_A - y) .$$
(A.124)

Some identities that are useful for mapping between results in the literature include

$$\frac{x_A}{r\xi_A} = \frac{1}{\left(1 + \xi_A^2 M_A^2/Q^2\right)} \quad \text{and} \quad \frac{x_A}{\xi_A} = \frac{1}{\left(1 - \xi_A^2 M_A^2/Q^2\right)} \quad (A.125)$$

Replacing the δ function with Eq. (A.124) allows us to finally write

$$\tilde{F}_{2}^{A}(x_{A}) = x_{A}^{2} \frac{d^{2}}{dx_{A}^{2}} \int_{0}^{1} dy \frac{\tilde{g}_{2}(y)}{y(1-y^{2}M_{A}^{2}/Q^{2})^{2}} \left[\frac{y(1-y^{2}M_{A}^{2}/Q^{2})}{(1+y^{2}M_{A}^{2}/Q^{2})}\right] \delta(\xi-y)$$
(A.126)

$$\tilde{F}_{2}^{A}(x_{A}) = x_{A}^{2} \frac{d^{2}}{dx_{A}^{2}} \left[\frac{(1+r)^{2}}{4r} \, \tilde{g}_{2}(\xi_{A}) \right] .$$
(A.127)

Allowing the derivative to act on the quantities inside the bracket recovers Eq. (3.24b). The final expression is summarized in Eq. (A.161).

 \tilde{F}_3^A : To build \tilde{F}_3^A , we similarly use Eq. (A.61) and identify everything to the right of $x_A^{-(2l-1)}$ in Eq. (A.100) as the $2l^{th}$ moment of $\tilde{F}_3^A(x_A, Q^2)$ with TMCs. We then write

$$\tilde{F}_{3}^{A(2l)} = \sum_{j=0}^{\infty} \frac{(2l+j)!}{j!(2l)!} \left(\frac{M_{A}^{2}}{Q^{2}}\right)^{j} (2l) \frac{\left(C_{3}^{(2l+2j)}A_{\tau=2}^{(2l+2j)}\right)}{(2l+2j)} .$$
(A.128)

Using the integral identity of Eq. (A.110a) and the relationship in Eq. (A.108) between the moment of F_3^A and the product $(C_3A_{\tau=2})$ when $(M_A^2/Q^2) \rightarrow 0$, we have

$$\frac{C_3^{(2l+2j)}A_{\tau=2}^{(2l+2j)}}{(2l+2j)} = \frac{1}{(2l+2j)} \int_0^1 dy \ y^{2l+2j} \cdot y^{-1}\tilde{F}_3^A(y) \bigg|_{\text{No TMCs}}$$
(A.129)

$$= \int_0^1 dy \ y^{2l+2j-1} \ \tilde{h}_3(y), \quad \text{where}$$
 (A.130)

$$\tilde{h}_3(y) \equiv \int_y^1 dy' (y')^{-1} \tilde{F}_3^A(y') \bigg|_{\text{No TMCs}} .$$
(A.131)

Using this, the summation over j is then given by

$$\tilde{F}_{3}^{A(2l)} = \int_{0}^{1} dy \ y^{2l-1} \ \tilde{h}_{3}(y) \ (2l) \ \sum_{j=0}^{\infty} \ \frac{(2l+j)!}{j!(2l)!} \ \left(\frac{y^{2}M_{A}^{2}}{Q^{2}}\right)^{j}$$
(A.132)

$$= \int_0^1 dy \ y^{-1} \ \tilde{h}_3(y) \ \frac{(2l) \ y^{2l}}{(1 - y^2 M_A^2/Q^2)^{2l+1}} \ . \tag{A.133}$$

After relabeling N = 2l, using the first derivative identity in Eq. (A.117), the contour integral of Eq. (A.121), the δ function decomposition of Eq. (A.124), and the identities for ξ_A , we get as

the inverse Mellin transformation of \tilde{F}_3^A

$$\tilde{F}_{3}^{A}(x_{A}) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} dN \ x_{A}^{-N} \ \tilde{F}_{3}^{AN}$$
(A.134)

$$= \frac{-x_A}{2\pi i} \frac{d}{dx_A} \int_0^1 dy \; \frac{h_3(y)}{y(1-y^2 M_A^2/Q^2)} \; \int_{c-i\infty}^{c+i\infty} dN \; \frac{(y/x_A)^N}{(1-y^2 M_A^2/Q^2)^N} \tag{A.135}$$

$$= -x_A \frac{d}{dx_A} \int_0^1 dy \; \frac{\tilde{h}_3(y)}{y(1-y^2 M_A^2/Q^2)} \left[\frac{y(1-y^2 M_A^2/Q^2)}{(1+y^2 M_A^2/Q^2)} \right] \delta(\xi_A - y) \;, \qquad (A.136)$$

$$\tilde{F}_{3}^{A}(x_{A}) = -x_{A} \frac{d}{dx_{A}} \left[\frac{(1+r)}{2r} \tilde{h}_{3}(\xi_{A}) \right]$$
(A.137)

Allowing the derivative to act on the quantities inside the bracket recovers Eq. (3.24c). The final expression is summarized in Eq. (A.161).

 \tilde{F}_6^A : Building \tilde{F}_6^A with TMCs proceeds identically to \tilde{F}_3^A . We first identify everything to the right of $x_A^{-(2l-1)}$ in Eq. (A.106) as the $2l^{th}$ moment of $\tilde{F}_6^A(x_A, Q^2)$. Explicitly,

$$\tilde{F}_{3}^{A(2l)} = \sum_{j=0}^{\infty} \frac{(2l+j)!}{j!(2l)!} \left(\frac{M_{A}^{2}}{Q^{2}}\right)^{j} (2l) \frac{\left(C_{6}^{(2l+2j)}A_{\tau=2}^{(2l+2j)}\right)}{(2l+2j)} .$$
(A.138)

The procedure for this case is similar to \tilde{F}_3^A . Using the relationship in Eq. (A.108), we obtain

$$\tilde{F}_{6}^{A}(x_{A}) = -x \frac{d}{dx_{A}} \left[\frac{(1+r)}{2r} \tilde{h}_{6}(y) \right], \quad \text{where} \quad \tilde{h}_{6}(y) \equiv \int_{y}^{1} dy' y'^{-1} \left. \tilde{F}_{6}^{A}(y') \right|_{\text{No TMCs}}.$$
(A.139)

The final expression is summarized in Eq. (A.161). To our knowledge, the TMCs to the \tilde{F}_6 structure function have not been previously reported in the literature.

 \tilde{F}_1^A : In principle, building \tilde{F}_1^A follows the same procedure as the previous cases. However, the nuance here is that there are contributions from $(C_1A_{\tau=2})$ and $(C_2A_{\tau=2})$ when j > 0. The solution is to decompose these into the moments of \tilde{F}_1^A and \tilde{F}_2^A , respectively. Using Eqs. (A.61) and (A.107), we identify everything to the right of $x_A^{-(2l+1)}$ in Eq. (A.96) as the $(2l+1)^{st}$ moment of $\tilde{F}_1^A(x_A, Q^2)$ with TMCs. Symbolically, this is given by

$$\tilde{F}_{1}^{A(2l+1)} = \sum_{j=0}^{\infty} \frac{(2l+j+1)!}{j! (2l+1)!} \left(\frac{M_{A}^{2}}{Q^{2}}\right)^{j} \left[\left(C_{1}^{2l+2j+1}A_{\tau=2}^{2l+2j+1}\right) + \frac{j\left(C_{2}^{2l+2j+1}A_{\tau=2}^{2l+2j+1}\right)}{(2l+2j+1)(2l+2j)} \right]$$
(A.140)

$$= \int_0^1 dy \, \sum_{j=0}^\infty \frac{(2l+j+1)!}{j! \, (2l+1)!} \, \left(\frac{M_A^2}{Q^2}\right)^j \left[y^{2l+2j} \tilde{F}_1^A(y) \Big|_{\text{No TMCs}} + j \, y^{2l+2j-1} \tilde{g}_2(y) \right], \quad (A.141)$$

where we expressed $(C_2A_{\tau=2})$ as an integral over $\tilde{g}_2(y)$ according to Eq. (A.111). Using Eq. (A.116),

we evaluate the summation of each term separately, giving

$$\tilde{F}_{1}^{A(2l+1)} = \int_{0}^{1} dy \; \frac{\tilde{F}_{1}^{A}(y)|_{\text{No TMCs}}}{y(1-y^{2}M_{A}^{2}/Q^{2})} \left[\frac{y^{2l+1}}{(1-y^{2}M_{A}^{2}/Q^{2})^{2l+1}} \right] \\
+ \left(\frac{M_{A}^{2}}{Q^{2}} \right) \int_{0}^{1} dy \; \frac{\tilde{g}_{2}(y)}{(1-y^{2}M_{A}^{2}/Q^{2})^{2}} \left[\frac{(2l+2)y^{2l+1}}{(1-y^{2}M_{A}^{2}/Q^{2})^{2l+1}} \right].$$
(A.142)

We now set N = (2l + 1), take the inverse Mellin transformation with respect to N, and use the first derivative identity in Eq. (A.117) to remove the (N + 1). The result is

$$\tilde{F}_{1}^{A}(x_{A}) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} dN \ x_{A}^{-N} \ \tilde{F}_{1}^{AN}$$

$$= \frac{1}{2\pi i} \int_{c-i\infty}^{1} dy \ \frac{\tilde{F}_{1}^{A}(y)|_{\text{No TMCs}}}{(1-2M^{2}/C^{2})} \int_{c-i\infty}^{c+i\infty} dN \left[\frac{(y/x_{A})^{N}}{(1-2M^{2}/C^{2})^{N}} \right]$$
(A.143)

$$= \left[\frac{M_A^2 x_A^2}{2\pi i Q^2}\right] \frac{d}{dx_A} \int_0^1 dy \, \frac{\tilde{g}_2(y)}{(1 - y^2 M_A^2/Q^2)^2} \int_{c-i\infty}^{c+i\infty} dN \, \left[\frac{(y/x_A)^N \, x_A^{-1}}{(1 - y^2 M_A^2/Q^2)^N}\right]$$
(A.144)
$$= \left[\frac{\tilde{F}_1^A(y)|_{\text{No TMCs}}}{(1 + y^2 M_A^2/Q^2)}\right]_{u=\xi_A}$$

$$-\left(\frac{M_A^2 x_A^2}{Q^2}\right) \frac{d}{dx_A} \left[\frac{\tilde{g}_2(y) \ x_A^{-1} \ y}{(1-y^2 M_A^2/Q^2)(1+y^2 M_A^2/Q^2)}\right]_{y=\xi_A} .$$
(A.145)

To reach the last line, we used the contour integral of Eq. (A.121) and the δ function decomposition of Eq. (A.124). Using the ξ_A identities, the final expressions is

$$\left. \tilde{F}_{1}^{A}(x_{A}) = \frac{x_{A}}{r\xi_{A}} \left. \tilde{F}_{1}^{A}(\xi_{A}) \right|_{\text{No TMCs}} - \left(\frac{M_{A}^{2} x_{A}^{2}}{Q^{2}} \right) \left. \frac{d}{dx_{A}} \left[\frac{(1+r)}{2r} \, \tilde{g}_{2}(\xi_{A}) \right] \right.$$
(A.146)

Applying the derivative to the quantities inside the bracket recovers Eq. (3.24a). Note that the factor of 1/2 in the square brackets is absent in Eq. (14) of Ref. [60]. This appears to be a typo in Ref. [60] as the final result matches Eq. (3.24a). Furthermore, after employing ξ_A identities, we find agreement with Eq. (3.1) of Ref. [54]. The final expression is summarized in Eq. (A.161).

 \tilde{F}_5^A : Building \tilde{F}_5^A with TMCs, which involves mixing with \tilde{F}_2^A , is similar to \tilde{F}_1^A . Subsequently, we identify everything to the right of $x_A^{-(2l)}$ in Eq. (A.104) as the $(2l+1)^{th}$ moment of $\tilde{F}_5^A(x_A, Q^2)$:

$$\begin{split} \tilde{F}_{5}^{A(2l+1)} &= \sum_{j=0}^{\infty} \frac{(2l+j+1)!}{j! \ (2l+1)!} \ \left(\frac{M_{A}^{2}}{Q^{2}}\right)^{j} \\ \times \left[\frac{(2l+1) \ \left(C_{5}^{(2l+2j+1)} A_{\tau=2}^{(2l+2j+1)}\right)}{(2l+2j+1)} - \frac{j \ (2l+1) \ \left(C_{2}^{(2l+2j+1)} A_{\tau=2}^{(2l+2j+1)}\right)}{(2l+2j+1)(2l+2j)}\right] + \mathcal{O}(\tau > 2) \quad (A.147) \\ &= \int_{0}^{1} dy \ \sum_{j=0}^{\infty} \frac{(2l+j+1)!}{j! \ (2l+1)!} \ \left(\frac{M_{A}^{2}}{Q^{2}}\right)^{j} \left[\frac{(2l+1)}{2} \ y^{2l+2j} \ \tilde{h}_{5}(y) - j(2l+1) \ y^{2l+2j-1} \ \tilde{g}_{2}(y)\right] \\ &+ \mathcal{O}(\tau > 2), \quad (A.148) \end{split}$$

where we used Eqs. (A.108) and (A.111) to decompose both $(C_i A_{\tau=2})$ products as integrals over $\tilde{F}_i^A(y)|_{\text{No TMCs}}$, and have defined via the identity Eq. (A.110a) the generating integral

$$\tilde{h}_5(y) \equiv \int_y^1 dy' \, (y')^{-1} \left. 2\tilde{F}_5^A(y') \right|_{\text{No TMCs}}.$$
(A.149)

Note the factor of 2 in the definition of \tilde{h}_5 , which follows the convention of Eq. (3.15) in Ref. [54]. Using Eq. (A.116), we evaluate the summation of each term separately. This gives

$$\tilde{F}_{5}^{A(2l+1)} = \int_{0}^{1} dy \; \frac{\tilde{h}_{5}(y)}{2y(1-y^{2}M_{A}^{2}/Q^{2})} \left[\frac{(2l+1)\; y^{2l+1}}{(1-y^{2}M_{A}^{2}/Q^{2})^{(2l+1)}} \right] \\ - \left(\frac{M_{A}^{2}}{Q^{2}}\right) \; \int_{0}^{1} dy \; \frac{\tilde{g}_{2}(y)}{(1-y^{2}M_{A}^{2}/Q^{2})^{2}} \left[\frac{(2l+2)\; (2l+1)\; y^{2l+1}}{(1-y^{2}M_{A}^{2}/Q^{2})^{(2l+1)}} \right] + \mathcal{O}(\tau > 2).$$
(A.150)

Setting N = (2l + 1), taking the inverse Mellin transform with respect to x_A , and using the derivative identities of Eq. (A.117) allows us to obtain $\tilde{F}_5^A(x_A)$. It is given by

$$\begin{split} \tilde{F}_{5}^{A}(x_{A}) &= \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} dN \ x_{A}^{-N} \ \tilde{F}_{5}^{A(N)} \end{split} \tag{A.151} \\ &= \left(\frac{-x_{A}}{4\pi i}\right) \frac{d}{dx_{A}} \int_{0}^{1} dy \ \frac{\tilde{h}_{5}(y)}{y \ (1-y^{2}M_{A}^{2}/Q^{2})} \ \int_{c-i\infty}^{c+i\infty} dN \ \left[\frac{(y/x_{A})^{N}}{(1-y^{2}M_{A}^{2}/Q^{2})^{N}}\right] \\ &- \left(\frac{x_{A}^{2}M_{A}^{2}}{2\pi i Q^{2}}\right) \frac{d^{2}}{dx_{A}^{2}} \int_{0}^{1} dy \ \frac{\tilde{g}_{2}(y)}{(1-y^{2}M_{A}^{2}/Q^{2})^{2}} \ \int_{c-i\infty}^{c+i\infty} dN \ \left[\frac{(y/x_{A})^{N}}{(1-y^{2}M_{A}^{2}/Q^{2})^{N}}\right] \\ &+ \mathcal{O}(\tau > 2). \end{aligned} \tag{A.152}$$

Applying the contour integral of Eq. (A.121) and the identity of Eq. (A.124), we get

$$\tilde{F}_{5}^{A}(x_{A}) = \left(\frac{-x_{A}}{2}\right) \frac{d}{dx_{A}} \left[\frac{(1+r)}{2r} \tilde{h}_{5}(\xi_{A})\right] - \left(\frac{x_{A}^{2}M_{A}^{2}}{Q^{2}}\right) \frac{d^{2}}{dx_{A}^{2}} \left[\frac{(1+r)^{2}}{4r} \xi_{A} \tilde{g}_{2}(\xi_{A})\right] + \mathcal{O}(\tau > 2) .$$
(A.153)

After accounting for the difference between \tilde{W}_5^A and \tilde{F}_5^A , this agrees with Eq. (3.5) of Ref. [54]. The final expression is summarized in Eq. (A.161).

 \tilde{F}_4^A : Building \tilde{F}_4^A follows the same procedure as above with the complication that the result is sourced by three contributions for j > 0. Using Eqs. (A.61) and (A.107), we identify everything to the right of $x_A^{-(2l+1)}$ in Eq. (A.102) as the $(2l+1)^{st}$ moment of $\tilde{F}_4^A(x_A, Q^2)$. This is given by

$$\begin{split} \tilde{F}_{4}^{A(2l+1)} &= \sum_{j=0}^{\infty} \frac{(2l+j+1)!}{j! (2l+1)!} \left(\frac{M_{A}^{2}}{Q^{2}}\right)^{j} \\ &\times \left[\left(C_{4}^{(2l+2j+1)} A_{\tau=2}^{(2l+2j+1)} \right) + \frac{j(j-1) \left(C_{2}^{(2l+2j+1)} A_{\tau=2}^{(2l+2j+1)} \right)}{(2l+2j+1) (2l+2j)} - \frac{2j \left(C_{5}^{(2l+2j+1)} A_{\tau=2}^{(2l+2j+1)} \right)}{(2l+2j+1)} \right] \\ &+ \mathcal{O}(\tau > 2), \quad (A.154) \\ &= \int_{0}^{1} dy \sum_{j=0}^{\infty} \frac{(2l+j+1)!}{j! (2l+1)!} \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{j} \\ &\times \left[y^{2l+2j} \tilde{F}_{4}^{A}(y) |_{\text{No TMC}} + j(j-1) y^{2l+2j-1} \tilde{g}_{2}(y) - j y^{2l+2j} \tilde{h}_{5}(y) \right] + \mathcal{O}(\tau > 2) . \quad (A.155) \end{split}$$

Here, we again used the Eqs. (A.108), (A.111), and (A.149) to rewrite $(C_i A_{\tau=2})$. Distributing the summations and using the identities listed in Eq. (A.116), we obtain

$$\begin{split} \tilde{F}_{4}^{A(2l+1)} &= \int_{0}^{1} dy \left[y^{2l} \; \tilde{F}_{4}^{A}(y) |_{\text{No TMC}} \sum_{j=0}^{\infty} \frac{(2l+j+1)!}{j! \; (2l+1)!} \left(\frac{M_{A}^{2}y^{2}}{Q^{2}} \right)^{j} \right. \\ &+ y^{2l-1} \; \tilde{g}_{2}(y) \sum_{j=0}^{\infty} j(j-1) \frac{(2l+j+1)!}{j! \; (2l+1)!} \left(\frac{M_{A}^{2}y^{2}}{Q^{2}} \right)^{j} \\ &- y^{2l} \; \tilde{h}_{5}(y) \; \sum_{j=0}^{\infty} j \frac{(2l+j+1)!}{j! \; (2l+1)!} \left(\frac{M_{A}^{2}y^{2}}{Q^{2}} \right)^{j} \right] + \; \mathcal{O}(\tau > 2) \quad (A.156) \\ &= \int_{0}^{1} dy \left[\frac{\tilde{F}_{4}^{A}(y)|_{\text{No TMC}}}{y(1-y^{2}M_{A}^{2}/Q^{2})} \left[\frac{y^{2l+1}}{(1-y^{2}M_{A}^{2}/Q^{2})^{2l+1}} \right] \\ &+ \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{2} \; \frac{y^{2} \; \tilde{g}_{2}(y)}{(1-y^{2}M_{A}^{2}/Q^{2})^{3}} \left[\frac{(2l+2)(2l+3)y^{2l+1}}{(1-y^{2}M_{A}^{2}/Q^{2})^{2l+1}} \right] \\ &- \left(\frac{M_{A}^{2}}{Q^{2}} \right) \; \frac{y \; \tilde{h}_{5}(y)}{(1-y^{2}M_{A}^{2}/Q^{2})} \left[\frac{(2l+2)y^{2l+1}}{(1-y^{2}M_{A}^{2}/Q^{2})^{2l+1}} \right] \\ &+ \; \mathcal{O}(\tau > 2) \; . \quad (A.157) \end{split}$$

Relabeling N = (2l + 1) and using the derivative identities in Eq. (A.117) gives

$$\begin{split} \tilde{F}_{4}^{A}(x_{A}) &= \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} dN \; x_{A}^{-N} \; \tilde{F}_{4}^{AN} \quad (A.158) \\ &= \frac{1}{2\pi i} \; \int_{0}^{1} dy \; \frac{\tilde{F}_{4}^{A}(y)|_{\text{No TMC}}}{y(1-y^{2}M_{A}^{2}/Q^{2})} \; \int_{c-i\infty}^{c+i\infty} dN \; \left[\frac{(y/x_{A})^{N}}{(1-y^{2}M_{A}^{2}/Q^{2})^{N}} \right] \\ &+ \left(\frac{M_{A}^{2}}{Q^{2}} \right)^{2} \; \left(\frac{x_{A}^{3}}{2\pi i} \right) \; \frac{d^{2}}{dx_{A}^{2}} \; \int_{0}^{1} dy \; \frac{y^{2} \; x_{A}^{-1} \; \tilde{g}_{2}(y)}{(1-y^{2}M_{A}^{2}/Q^{2})^{3}} \; \int_{c-i\infty}^{c+i\infty} dN \; \left[\frac{(y/x_{A})^{N}}{(1-y^{2}M_{A}^{2}/Q^{2})^{N}} \right] \\ &+ (-1)^{2} \; \left(\frac{M_{A}^{2}}{Q^{2}} \right) \; \left(\frac{x_{A}^{2}}{2\pi i} \right) \; \frac{d}{dx_{A}} \; \int_{0}^{1} dy \; \frac{y \; x_{A}^{-1} \; \tilde{h}_{5}(y)}{(1-y^{2}M_{A}^{2}/Q^{2})^{2}} \; \int_{c-i\infty}^{c+i\infty} dN \; \left[\frac{(y/x_{A})^{N}}{(1-y^{2}M_{A}^{2}/Q^{2})^{N}} \right] \\ &+ \; \mathcal{O}(\tau > 2) \; . \end{split}$$

Finally, using contour integral of Eq. (A.121) and the ξ_A identities, we get

$$\tilde{F}_{4}^{A}(x_{A}) = \frac{(1+r)}{2r} \tilde{F}_{4}^{A}(\xi_{A}) \Big|_{\text{No TMC}} + \left(\frac{M_{A}^{2}}{Q^{2}}\right)^{2} x_{A}^{3} \frac{d^{2}}{dx_{A}^{2}} \left[\left(\frac{x_{A}}{r}\right) \tilde{g}_{2}(\xi_{A})\right] \\
+ \left(\frac{M_{A}^{2}}{Q^{2}}\right) x_{A}^{2} \frac{d}{dx_{A}} \left[\left(\frac{x_{A}}{r}\right) \tilde{h}_{5}(\xi_{A})\right] + \mathcal{O}(\tau > 2) .$$
(A.160)

After accounting for differences between \tilde{W}_4^A and \tilde{F}_4^A , this agrees with Eq. (3.4) of Ref. [54]. The final expression is summarized in Eq. (A.161).

Summary

We now give the full expressions for all six structure functions with TMCs. These are obtained by successive applications of the chain rule to the several expressions above. The final results are:

$$\begin{split} \tilde{F}_{1}^{A}(x_{A}) &= \frac{x_{A}}{\xi_{A}r} \left. \tilde{F}_{1}^{A}(\xi_{A}) \right|_{\text{No TMCs}} + \left(\frac{M_{A}^{2}x_{A}^{2}}{Q^{2}r^{2}} \right) \tilde{h}_{2}(\xi_{A}) + \left(\frac{2M_{A}^{4}x_{A}^{3}}{Q^{4}r^{3}} \right) \tilde{g}_{2}(\xi_{A}) + \mathcal{O}(\tau > 2) , \quad (A.161a) \\ \tilde{F}_{2}^{A}(x_{A}) &= \frac{x_{A}^{2}}{\xi_{A}^{2}r^{3}} \left. \tilde{F}_{2}^{A}(\xi_{A}) \right|_{\text{No TMCs}} + \left(\frac{6M_{A}^{2}x_{A}^{3}}{Q^{2}r^{4}} \right) \tilde{h}_{2}(\xi_{A}) + \left(\frac{12M_{A}^{4}x_{A}^{4}}{Q^{4}r^{5}} \right) \tilde{g}_{2}(\xi_{A}) \\ &+ \mathcal{O}(\tau > 2) , \qquad (A.161b) \end{split}$$

$$\tilde{F}_{3}^{A}(x_{A}) = \frac{x_{A}}{\xi_{A}r_{A}^{2}} \tilde{F}_{3}^{A}(\xi_{A}) \bigg|_{\text{No TMCs}} + \left(\frac{2M_{A}^{2}x_{A}^{2}}{Q^{2}r^{3}}\right) \tilde{h}_{3}(\xi_{A}) + \mathcal{O}(\tau > 2) , \qquad (A.161c)$$

$$\tilde{F}_{4}^{A}(x_{A}) = \frac{x_{A}}{\xi_{A}r} \left. \tilde{F}_{4}^{A}(\xi_{A}) \right|_{\text{No TMCs}} - \left(\frac{2M_{A}^{2}x_{A}^{2}}{Q^{2}r^{2}} \right) \left. \tilde{F}_{5}^{A}(\xi_{A}) \right|_{\text{No TMCs}} + \left(\frac{M_{A}^{4}x_{A}^{3}}{Q^{4}r^{3}} \right) \left. \tilde{F}_{2}^{A}(\xi_{A}) \right|_{\text{No TMCs}} + \left(\frac{M_{A}^{2}x_{A}^{2}}{Q^{2}r^{3}} \right) \left. \tilde{h}_{5}^{A}(\xi_{A}) - \left(\frac{2M_{A}^{4}x_{A}^{4}}{Q^{4}r^{4}} \right) \left(2 - \xi^{2}M_{A}^{2}/Q^{2} \right) \left. \tilde{h}_{2}^{A}(\xi_{A}) + \left(\frac{2M_{A}^{4}x_{A}^{3}}{Q^{4}r^{5}} \right) \left(1 - 2x_{A}^{2}M_{A}^{2}/Q^{2} \right) \left. \tilde{g}_{2}^{A}(\xi_{A}) + \mathcal{O}(\tau > 2) \right. \right. \tag{A.161d}$$

$$\tilde{F}_{5}^{A}(x_{A}) = \frac{x_{A}}{\xi_{A}r^{2}} \tilde{F}_{5}^{A}(\xi_{A}) \Big|_{\text{No TMCs}} - \left(\frac{M_{A}^{2}x_{A}^{2}}{Q^{2}r^{3}\xi_{A}}\right) \tilde{F}_{2}^{A}(\xi_{A}) \Big|_{\text{No TMCs}} \\
+ \left(\frac{M_{A}^{2}x_{A}^{2}}{Q^{2}r^{3}}\right) \tilde{h}_{5}^{A}(\xi_{A}) - \left(\frac{2M_{A}^{2}x_{A}^{2}}{Q^{2}r^{4}}\right) \left(1 - x\xi M_{A}^{2}/Q^{2}\right) \tilde{h}_{2}^{A}(\xi_{A}) \\
+ \left(\frac{6M_{A}^{4}x_{A}^{3}}{Q^{4}r^{5}}\right) \tilde{g}_{2}^{A}(\xi_{A}) + \mathcal{O}(\tau > 2) ,$$
(A.161e)

$$\tilde{F}_{6}^{A}(x_{A}) = \frac{x_{A}}{\xi_{A}r_{A}^{2}} \tilde{F}_{6}^{A}(\xi_{A}) \bigg|_{\text{No TMCs}} + \left(\frac{2M_{A}^{2}x_{A}^{2}}{Q^{2}r^{3}}\right) \tilde{h}_{6}(\xi_{A}) + \mathcal{O}(\tau > 2) .$$
(A.161f)

Importantly, the above expressions for *nuclear* structure functions $\tilde{F}_1, \ldots, \tilde{F}_5$ agree with those for *nucleon* structure functions F_1, \ldots, F_5 in Eq. (3.17) of Ref. [60]. This is a main conclusion of our work: that TMCs for unpolarized protons and neutrons are the same for unpolarized nuclei.

Appendix B. Derivation of the Full/Leading TMC Parameterization

In this appendix, we derive our parameterizations of Eq. (7.8) by approximating the general form of the structure functions in the TMC region. First, we assume the structure functions $F_a^{(0)}(x)$, a = 1, 2, 3, vanish at x = 1. This assumption follows from the fact that the PDFs vanish at x = 1.

Next, we use a finite difference formula to approximate derivatives of $F_a^{j,(0)}(x = \xi)$ as

$$F_{a}^{'(0)}(u=\xi) \approx -\frac{\gamma_{a}F_{a}^{(0)}(\xi)}{1-\xi}$$
(B.1)

$$F_a^{j,(0)}(u=\xi) \approx \frac{0 - F_a^{j-1,(0)}(\xi)}{(1-\xi)} = (-1)^j \frac{\gamma_a F_a^{(0)}(\xi)}{(1-\xi)^j} \quad . \tag{B.2}$$

Here, γ_a is a universal correction factor for the first derivative which we assume to be independent of x and the underlying PDFs, and to have a mild Q dependence. We can then expand $F_a^{(0)}(u)$ about $u = \xi$ as

$$F_a^{(0)}(u) = \sum_{j=0}^{\infty} \frac{1}{j!} F_a^{j,(0)}(\xi) (u - \xi)^j$$
(B.3)

$$\approx F_a^{(0)}(\xi) \left(1 + \sum_{j=1}^{\infty} \frac{1}{j!} (-1)^j \frac{\gamma_a}{(1-\xi)^j} (u-\xi)^j \right)$$
(B.4)

$$\equiv F_a^{(0)}(\xi) K_a(u,\xi,\gamma_a) \quad . \tag{B.5}$$

Here,

$$K_a(u,\xi,\gamma_a) = 1 + \sum_{j=1}^{\infty} \frac{1}{j!} (-1)^j \frac{\gamma_a}{(1-\xi)^j} (u-\xi)^j .$$
(B.6)

As $K_a(u,\xi,\gamma)$ is independent of the PDFs, this implies that the ratios

$$\frac{h_a(\xi)}{F_a^{(0)}(\xi)} = \int_{\xi}^{1} L_a(u) K_a(\xi, x, \gamma) du$$
(B.7)

$$\frac{g_2(\xi)}{F_2^{(0)}(\xi)} = \int_{\xi}^{1} \frac{u-\xi}{u^2} K_a(\xi, x, \gamma) du$$
(B.8)

are also independent of the underlying PDFs. Here we have defined $L_1(u) = 2/u$, $L_2(u) = 1/u^2$, and $L_3(u) = 1/u$.

Evaluating the explicit expression for $K_a(u,\xi,\gamma_a)$ in the above relations, we obtain (7.8a), (7.8b), and (7.8d). To calculate $h_2(\xi)/F_1^{(0)}(\xi)$ and $g_2(\xi)/F_1^{(0)}(\xi)$, which are needed to calculate the ratio $F_1^{TMC}(x)/F_1^{Leading-TMC}$, one can assume an approximate Callan-Gross relation. Deviations from the Callan-Gross relation can be absorbed into the fitted $\gamma_1(Q)$. We note that in practice, we calculate the massless structure function using Eqs. (6.10), namely :

$$F_1^{(0)}(\xi) = F_1^{TMC-ACOT}(x),$$
(B.9a)

$$F_2^{(0)}(\xi) = \frac{\xi r^2}{x} F_2^{TMC-ACOT}(x),$$
(B.9b)

$$F_3^{(0)}(\xi) = r F_3^{TMC-ACOT}(x), \tag{B.9c}$$

Assuming Callan-Gross relation at the level of the TMC-ACOT structure functions, implies

$$F_2^{(0)}(\xi) = 2\xi r^2 F_1^{(0)}(\xi) \tag{B.10}$$

Using (B.10), one can easily obtain $h_2(\xi)/F_1^{(0)}(\xi)$ and $g_2(\xi)/F_1^{(0)}(\xi)$ from $h_2(\xi)/F_2^{(0)}(\xi)$ and $g_2(\xi)/F_2^{(0)}(\xi)$, as shown in (7.8c) and (7.8e).

If we perform a single parameter fit and take γ_a as a constant, we find reasonable agreement with the exact TMC result at the level of $\leq 0.75\%$ for Q = 1.3 GeV and Q = 2 GeV. We can do even better if we perform a 2-parameter fit and replace γ_a with:

$$\gamma_a \to \gamma_a(Q) = \lambda_a \ln(Q)^{\delta_a}$$
 (B.11)

Thus, we give a mild Q dependence to the parameter γ_a . This 2-parameter fit is superior to the one parameter fit as the agreement is $\leq 0.6\%$ for Q = 1.3 GeV, and $\leq 0.2\%$ for Q = 2 GeV which is a typical kinematic cut for the global PDF fits.

References

- [1] R. E. Taylor, "Deep inelastic scattering: The Early years," Rev. Mod. Phys. 63 (1991) 573–595.
- [2] H. W. Kendall, "Deep inelastic scattering: Experiments on the proton and the observation of scaling," *Rev. Mod. Phys.* 63 (1991) 597-614.
- [3] J. I. Friedman, "Deep inelastic scattering: Comparisons with the quark model," *Rev. Mod. Phys.* 63 (1991) 615–629.
- [4] R. Devenish and A. Cooper-Sarkar, *Deep inelastic scattering*. Oxford University Press, 2004.
- [5] T.-J. Hou et al., "New CTEQ global analysis of quantum chromodynamics with high-precision data from the LHC," arXiv:1912.10053 [hep-ph].
- [6] A. Accardi, L. T. Brady, W. Melnitchouk, J. F. Owens, and N. Sato, "Constraints on large-x parton distributions from new weak boson production and deep-inelastic scattering data," *Phys. Rev. D* 93 no. 11, (2016) 114017, arXiv:1602.03154 [hep-ph].
- [7] S. Bailey, T. Cridge, L. A. Harland-Lang, A. D. Martin, and R. S. Thorne, "Parton distributions from LHC, HERA, Tevatron and fixed target data: MSHT20 PDFs," *Eur. Phys. J. C* 81 no. 4, (2021) 341, arXiv:2012.04684 [hep-ph].
- [8] H1, ZEUS Collaboration, H. Abramowicz et al., "Combination of measurements of inclusive deep inelastic e[±]p scattering cross sections and QCD analysis of HERA data," Eur. Phys. J. C 75 no. 12, (2015) 580, arXiv:1506.06042 [hep-ex].
- [9] NNPDF Collaboration, R. D. Ball et al., "Parton distributions from high-precision collider data," Eur. Phys. J. C77 no. 10, (2017) 663, arXiv:1706.00428 [hep-ph].
- [10] S. Alekhin, J. Blümlein, S. Moch, and R. Placakyte, "Parton distribution functions, α_s, and heavy-quark masses for LHC Run II," *Phys. Rev. D* 96 no. 1, (2017) 014011, arXiv:1701.05838 [hep-ph].
- [11] M. Hirai, S. Kumano, and T. H. Nagai, "Determination of nuclear parton distribution functions and their uncertainties in next-to-leading order," *Phys. Rev. C* 76 (2007) 065207, arXiv:0709.3038 [hep-ph].
- [12] D. de Florian, R. Sassot, P. Zurita, and M. Stratmann, "Global Analysis of Nuclear Parton Distributions," *Phys. Rev.* D85 (2012) 074028, arXiv:1112.6324 [hep-ph].
- [13] K. Eskola, H. Paukkunen, and C. Salgado, "EPS09: A New Generation of NLO and LO Nuclear Parton Distribution Functions," JHEP 04 (2009) 065, arXiv:0902.4154 [hep-ph].
- [14] K. J. Eskola, P. Paakkinen, H. Paukkunen, and C. A. Salgado, "EPPS16: Nuclear parton distributions with LHC data," *Eur. Phys. J.* C77 no. 3, (2017) 163, arXiv:1612.05741 [hep-ph].
- [15] K. J. Eskola, P. Paakkinen, H. Paukkunen, and C. A. Salgado, "EPPS21: a global QCD analysis of nuclear PDFs," *Eur. Phys. J. C* 82 no. 5, (2022) 413, arXiv:2112.12462 [hep-ph].
- [16] I. Schienbein, J. Y. Yu, K. Kovarik, C. Keppel, J. G. Morfin, F. Olness, and J. F. Owens, "PDF Nuclear Corrections for Charged and Neutral Current Processes," *Phys. Rev. D* 80 (2009) 094004, arXiv:0907.2357 [hep-ph].
- [17] K. Kovařík et al., "nCTEQ15 Global analysis of nuclear parton distributions with uncertainties in the CTEQ framework," Phys. Rev. D93 no. 8, (2016) 085037, arXiv:1509.00792 [hep-ph].
- [18] A. Kusina, F. Lyonnet, D. B. Clark, E. Godat, T. Jezo, K. Kovařík, F. I. Olness, I. Schienbein, and J. Y. Yu, "Vector boson production in pPb and PbPb collisions at the LHC and its impact on nCTEQ15 PDFs," *Eur. Phys. J.* C77 no. 7, (2017) 488, arXiv:1610.02925 [nucl-th].

- [19] E. P. Segarra *et al.*, "Extending nuclear PDF analyses into the high-x, low-Q² region," *Phys. Rev. D* 103 no. 11, (2021) 114015, arXiv:2012.11566 [hep-ph].
- [20] A. Kusina et al., "Impact of LHC vector boson production in heavy ion collisions on strange PDFs," Eur. Phys. J. C 80 no. 10, (2020) 968, arXiv:2007.09100 [hep-ph].
- [21] P. Duwentäster, L. A. Husová, T. Ježo, M. Klasen, K. Kovařík, A. Kusina, K. F. Muzakka, F. I. Olness, I. Schienbein, and J. Y. Yu, "Impact of inclusive hadron production data on nuclear gluon PDFs," arXiv:2105.09873 [hep-ph].
- [22] P. Duwentäster, T. Ježo, M. Klasen, K. Kovařík, A. Kusina, K. F. Muzakka, F. I. Olness, R. Ruiz, I. Schienbein, and J. Y. Yu, "Impact of heavy quark and quarkonium data on nuclear gluon PDFs," arXiv:2204.09982 [hep-ph].
- [23] K. F. Muzakka et al., "Compatibility of Neutrino DIS Data and Its Impact on Nuclear Parton Distribution Functions," arXiv:2204.13157 [hep-ph].
- [24] NNPDF Collaboration, R. Abdul Khalek, J. J. Ethier, and J. Rojo, "Nuclear parton distributions from lepton-nucleus scattering and the impact of an electron-ion collider," *Eur. Phys. J.* C79 no. 6, (2019) 471, arXiv:1904.00018 [hep-ph].
- [25] R. Abdul Khalek, J. J. Ethier, J. Rojo, and G. van Weelden, "nNNPDF2.0: Quark Flavor Separation in Nuclei from LHC Data," arXiv:2006.14629 [hep-ph].
- [26] R. A. Khalek, R. Gauld, T. Giani, E. R. Nocera, T. R. Rabemananjara, and J. Rojo, "nNNPDF3.0: Evidence for a modified partonic structure in heavy nuclei," arXiv:2201.12363 [hep-ph].
- [27] M. Walt, I. Helenius, and W. Vogelsang, "Open-source QCD analysis of nuclear parton distribution functions at NLO and NNLO," *Phys. Rev.* D100 no. 9, (2019) 096015, arXiv:1908.03355 [hep-ph].
- [28] H. Khanpour, M. Soleymaninia, S. Atashbar Tehrani, H. Spiesberger, and V. Guzey, "Nuclear parton distribution functions with uncertainties in a general mass variable flavor number scheme," *Phys. Rev.* D 104 no. 3, (2021) 034010, arXiv:2010.00555 [hep-ph].
- [29] A. Accardi et al., "Electron Ion Collider: The Next QCD Frontier: Understanding the glue that binds us all," Eur. Phys. J. A 52 no. 9, (2016) 268, arXiv:1212.1701 [nucl-ex].
- [30] R. Abdul Khalek et al., "Science Requirements and Detector Concepts for the Electron-Ion Collider: EIC Yellow Report," arXiv:2103.05419 [physics.ins-det].
- [31] J. L. Feng et al., "The Forward Physics Facility at the High-Luminosity LHC," arXiv:2203.05090 [hep-ex].
- [32] J. Arrington et al., "Physics with CEBAF at 12 GeV and future opportunities," Prog. Part. Nucl. Phys. 127 (2022) 103985, arXiv:2112.00060 [nucl-ex].
- [33] DUNE Collaboration, R. Acciarri et al., "Long-Baseline Neutrino Facility (LBNF) and Deep Underground Neutrino Experiment (DUNE): Conceptual Design Report, Volume 2: The Physics Program for DUNE at LBNF," arXiv:1512.06148 [physics.ins-det].
- [34] P. A. Machado, O. Palamara, and D. W. Schmitz, "The Short-Baseline Neutrino Program at Fermilab," Ann. Rev. Nucl. Part. Sci. 69 (2019) 363-387, arXiv:1903.04608 [hep-ex].
- [35] L. A. Ruso *et al.*, "Theoretical tools for neutrino scattering: interplay between lattice QCD, EFTs, nuclear physics, phenomenology, and neutrino event generators," arXiv:2203.09030 [hep-ph].
- [36] E. D. Bloom and F. J. Gilman, "Scaling, Duality, and the Behavior of Resonances in Inelastic electron-Proton Scattering," *Phys. Rev. Lett.* **25** (1970) 1140.

- [37] W. Melnitchouk, R. Ent, and C. Keppel, "Quark-hadron duality in electron scattering," *Phys. Rept.* 406 (2005) 127-301, arXiv:hep-ph/0501217.
- [38] K. G. Wilson, "Nonlagrangian models of current algebra," Phys. Rev. 179 (1969) 1499–1512.
- [39] R. A. Brandt and G. Preparata, "Operator product expansions near the light cone," Nucl. Phys. B 27 (1971) 541–567.
- [40] N. H. Christ, B. Hasslacher, and A. H. Mueller, "Light cone behavior of perturbation theory," *Phys. Rev. D* 6 (1972) 3543.
- [41] G. T. Bodwin, "Factorization of the Drell-Yan Cross-Section in Perturbation Theory," *Phys. Rev. D* 31 (1985) 2616. [Erratum: Phys.Rev.D 34, 3932 (1986)].
- [42] J. C. Collins, D. E. Soper, and G. F. Sterman, "Factorization for Short Distance Hadron Hadron Scattering," Nucl. Phys. B 261 (1985) 104–142.
- [43] J. C. Collins, D. E. Soper, and G. F. Sterman, "Soft Gluons and Factorization," Nucl. Phys. B 308 (1988) 833–856.
- [44] J. C. Collins and D. E. Soper, "The Theorems of Perturbative QCD," Ann. Rev. Nucl. Part. Sci. 37 (1987) 383–409.
- [45] J. C. Collins, D. E. Soper, and G. F. Sterman, "Factorization of Hard Processes in QCD," Adv. Ser. Direct. High Energy Phys. 5 (1989) 1-91, arXiv:hep-ph/0409313.
- [46] J. C. Collins, "Hard scattering factorization with heavy quarks: A General treatment," *Phys. Rev. D* 58 (1998) 094002, arXiv:hep-ph/9806259.
- [47] J. Collins, Foundations of perturbative QCD, vol. 32. Cambridge University Press, 11, 2013.
- [48] E. L. Berger and F. Coester, "Nuclear Effects in Deep Inelastic Lepton Scattering," Ann. Rev. Nucl. Part. Sci. 37 (1987) 463–491.
- [49] H. Georgi and H. D. Politzer, "Freedom at Moderate Energies: Masses in Color Dynamics," *Phys. Rev.* D 14 (1976) 1829.
- [50] H. Georgi and H. D. Politzer, "Precocious Scaling, Rescaling and xi Scaling," *Phys. Rev. Lett.* 36 (1976) 1281. [Erratum: Phys.Rev.Lett. 37, 68 (1976)].
- [51] R. Barbieri, J. R. Ellis, M. K. Gaillard, and G. G. Ross, "Mass Corrections to Scaling in Deep Inelastic Processes," *Nucl. Phys. B* 117 (1976) 50–76.
- [52] R. Barbieri, J. R. Ellis, M. K. Gaillard, and G. G. Ross, "A Quest for a Wholly Scaling Variable," *Phys. Lett. B* 64 (1976) 171–176.
- [53] A. De Rujula, H. Georgi, and H. D. Politzer, "Demythification of Electroproduction, Local Duality and Precocious Scaling," Annals Phys. 103 (1977) 315.
- [54] S. Kretzer and M. H. Reno, "Target mass corrections to electro-weak structure functions and perturbative neutrino cross sections," *Phys. Rev.* D69 (2004) 034002, hep-ph/0307023.
- [55] R. K. Ellis, W. Furmanski, and R. Petronzio, "Unraveling Higher Twists," Nucl. Phys. B 212 (1983) 29.
- [56] A. Accardi, T. Hobbs, and W. Melnitchouk, "Hadron mass corrections in semi-inclusive deep inelastic scattering," JHEP 11 (2009) 084, arXiv:0907.2395 [hep-ph].
- [57] M. A. G. Aivazis, F. I. Olness, and W.-K. Tung, "Leptoproduction of heavy quarks. 1. General formalism and kinematics of charged current and neutral current production processes," *Phys. Rev. D* 50 (1994) 3085–3101, arXiv:hep-ph/9312318.

- [58] M. A. G. Aivazis, J. C. Collins, F. I. Olness, and W.-K. Tung, "Leptoproduction of heavy quarks. 2. A Unified QCD formulation of charged and neutral current processes from fixed target to collider energies," *Phys. Rev. D* 50 (1994) 3102–3118, arXiv:hep-ph/9312319.
- [59] S. Kretzer and M. H. Reno, "Tau neutrino deep inelastic charged current interactions," Phys. Rev. D66 (2002) 113007, hep-ph/0208187.
- [60] I. Schienbein et al., "A Review of Target Mass Corrections," J. Phys. G35 (2008) 053101, arXiv:0709.1775 [hep-ph].
- [61] R. L. Jaffe, "Deep Inelastic Scattering with Application to Nuclear Targets: Lectures at the 1985 Los Alamos School on Relativistic Dynamics and Quark Nuclear Physics," in *Research Program at CEBAF* (I): Report of the 1985 Summer Study Group. 7, 1985. arXiv:2212.05616 [hep-ph].
- [62] J.-w. Qiu, "QCD factorization and rescattering in proton nucleus collisions," arXiv:hep-ph/0305161.
- [63] T. Muta, Foundations of Quantum Chromodynamics: An Introduction to Perturbative Methods in Gauge Theories, (2nd ed.), vol. 57 of World scientific Lecture Notes in Physics. World Scientific, Hackensack, N.J., 2nd ed., 1998.
- [64] T. Muta, Foundations of Quantum Chromodynamics: An Introduction to Perturbative Methods in Gauge Theories, (3rd ed.), vol. 78 of World scientific Lecture Notes in Physics. World Scientific, Hackensack, N.J., 3rd ed., 2010.
- [65] I. Schienbein, J. Y. Yu, C. Keppel, J. G. Morfin, F. Olness, and J. F. Owens, "Nuclear parton distribution functions from neutrino deep inelastic scattering," *Phys. Rev. D* 77 (2008) 054013, arXiv:0710.4897 [hep-ph].
- [66] K. Kovarik, I. Schienbein, F. I. Olness, J. Y. Yu, C. Keppel, J. G. Morfin, J. F. Owens, and T. Stavreva, "Nuclear Corrections in Neutrino-Nucleus DIS and Their Compatibility with Global NPDF Analyses," *Phys. Rev. Lett.* **106** (2011) 122301, arXiv:1012.0286 [hep-ph].
- [67] O. Nachtmann, "Positivity constraints for anomalous dimensions," Nucl. Phys. B 63 (1973) 237-247.
- [68] P. G. Blunden, W. Melnitchouk, and J. A. Tjon, "Two photon exchange and elastic electron proton scattering," *Phys. Rev. Lett.* **91** (2003) 142304, arXiv:nucl-th/0306076.
- [69] P. G. Blunden, W. Melnitchouk, and J. A. Tjon, "Two-photon exchange in elastic electron-nucleon scattering," *Phys. Rev. C* 72 (2005) 034612, arXiv:nucl-th/0506039.
- [70] J. Arrington, W. Melnitchouk, and J. A. Tjon, "Global analysis of proton elastic form factor data with two-photon exchange corrections," *Phys. Rev. C* 76 (2007) 035205, arXiv:0707.1861 [nucl-ex].
- [71] J. Arrington, P. G. Blunden, and W. Melnitchouk, "Review of two-photon exchange in electron scattering," Prog. Part. Nucl. Phys. 66 (2011) 782-833, arXiv:1105.0951 [nucl-th].
- [72] J. Blumlein and A. Tkabladze, "Target mass corrections for polarized structure functions and new sum rules," Nucl. Phys. B 553 (1999) 427-464, arXiv:hep-ph/9812478.
- [73] C. A. Aidala, S. D. Bass, D. Hasch, and G. K. Mallot, "The Spin Structure of the Nucleon," *Rev. Mod. Phys.* 85 (2013) 655-691, arXiv:1209.2803 [hep-ph].
- [74] A. Deshpande, R. Milner, R. Venugopalan, and W. Vogelsang, "Study of the fundamental structure of matter with an electron-ion collider," Ann. Rev. Nucl. Part. Sci. 55 (2005) 165–228, arXiv:hep-ph/0506148.
- [75] Particle Data Group Collaboration, P. A. Zyla *et al.*, "Review of Particle Physics," *PTEP* 2020 no. 8, (2020) 083C01.

- [76] S. Alekhin et al., "A facility to Search for Hidden Particles at the CERN SPS: the SHiP physics case," *Rept. Prog. Phys.* 79 no. 12, (2016) 124201, arXiv:1504.04855 [hep-ph].
- [77] FASER Collaboration, H. Abreu *et al.*, "Detecting and Studying High-Energy Collider Neutrinos with FASER at the LHC," *Eur. Phys. J. C* 80 no. 1, (2020) 61, arXiv:1908.02310 [hep-ex].
- [78] SHiP Collaboration, C. Ahdida et al., "SND@LHC," arXiv:2002.08722 [physics.ins-det].
- [79] P. Hoodbhoy, R. L. Jaffe, and A. Manohar, "Novel Effects in Deep Inelastic Scattering from Spin 1 Hadrons," Nucl. Phys. B 312 (1989) 571–588.
- [80] I. Schienbein, "Two photon processes and photon structure," Annals Phys. 301 (2002) 128–156, arXiv:hep-ph/0205301.
- [81] W. Detmold, "Target mass effects in deep-inelastic scattering on the deuteron," *Phys. Lett. B* 632 (2006) 261-269, arXiv:hep-ph/0509011.
- [82] Y. Kitadono, K. Sasaki, T. Ueda, and T. Uematsu, "Target Mass Corrections for the Virtual Photon Structure Functions to the Next-to-next-to-leading Order in QCD," *Phys. Rev. D* 77 (2008) 054019, arXiv:0801.0937 [hep-ph].
- [83] R. K. Ellis, W. Furmanski, and R. Petronzio, "Power Corrections to the Parton Model in QCD," Nucl. Phys. B 207 (1982) 1–14.
- [84] R. L. Jaffe and M. Soldate, "Twist Four in Electroproduction: Canonical Operators and Coefficient Functions," *Phys. Rev. D* 26 (1982) 49–68.
- [85] J. C. Collins, Renormalization: An Introduction to Renormalization, The Renormalization Group, and the Operator Product Expansion, vol. 26 of Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 1986.
- [86] A. Vogt, "Efficient evolution of unpolarized and polarized parton distributions with QCD-PEGASUS," *Comput. Phys. Commun.* 170 (2005) 65–92, arXiv:hep-ph/0408244.
- [87] J. L. Miramontes and J. Sanchez Guillen, "Understanding higher twist: operator approach to power corrections," Z. Phys. C 41 (1988) 247.
- [88] E. Moffat, W. Melnitchouk, T. C. Rogers, and N. Sato, "What are the low-Q and large-x boundaries of collinear QCD factorization theorems?," *Phys. Rev. D* 95 no. 9, (2017) 096008, arXiv:1702.03955 [hep-ph].
- [89] R. P. Feynman, "Very high-energy collisions of hadrons," Phys. Rev. Lett. 23 (1969) 1415-1417.
- [90] J. D. Bjorken and E. A. Paschos, "Inelastic Electron Proton and gamma Proton Scattering, and the Structure of the Nucleon," *Phys. Rev.* 185 (1969) 1975–1982.
- [91] F. I. Olness, R. J. Scalise, and W.-K. Tung, "Heavy quark hadroproduction in perturbative QCD," *Phys. Rev. D* 59 (1999) 014506, arXiv:hep-ph/9712494.
- [92] R. S. Thorne and R. G. Roberts, "A Variable number flavor scheme for charged current heavy flavor structure functions," *Eur. Phys. J. C* 19 (2001) 339–349, arXiv:hep-ph/0010344.
- [93] S. Alekhin, J. Blümlein, and S. Moch, "Heavy-flavor PDF evolution and variable-flavor number scheme uncertainties in deep-inelastic scattering," *Phys. Rev. D* 102 no. 5, (2020) 054014, arXiv:2006.07032 [hep-ph].
- [94] S. Alekhin, J. Blumlein, S. Klein, and S. Moch, "The 3, 4, and 5-flavor NNLO Parton from Deep-Inelastic-Scattering Data and at Hadron Colliders," *Phys. Rev. D* 81 (2010) 014032, arXiv:0908.2766 [hep-ph].

- [95] V. N. Gribov and L. N. Lipatov, "Deep inelastic e p scattering in perturbation theory," Sov. J. Nucl. Phys. 15 (1972) 438–450.
- [96] G. Altarelli and G. Parisi, "Asymptotic Freedom in Parton Language," Nucl. Phys. B 126 (1977) 298–318.
- [97] Y. L. Dokshitzer, "Calculation of the Structure Functions for Deep Inelastic Scattering and e⁺e[−] Annihilation by Perturbation Theory in Quantum Chromodynamics.," Sov. Phys. JETP 46 (1977) 641–653.
- [98] MARATHON Collaboration, D. Abrams et al., "Measurement of the Nucleon F₂ⁿ/F₂^p Structure Function Ratio by the Jefferson Lab MARATHON Tritium/Helium-3 Deep Inelastic Scattering Experiment," arXiv:2104.05850 [hep-ex].
- [99] C. Cocuzza, C. E. Keppel, H. Liu, W. Melnitchouk, A. Metz, N. Sato, and A. W. Thomas, "Isovector EMC effect from global QCD analysis with MARATHON data," arXiv:2104.06946 [hep-ph].
- [100] M. Sajjad Athar and J. G. Morfín, "Neutrino(antineutrino)-nucleus interactions in the shallow- and deep-inelastic scattering regions," J. Phys. G 48 no. 3, (2021) 034001, arXiv:2006.08603 [hep-ph].
- [101] I. Niculescu *et al.*, "Direct observation of quark-hadron duality in the free neutron F₂ structure function," *Phys. Rev. C* 91 no. 5, (2015) 055206, arXiv:1501.02203 [hep-ex].
- [102] L. L. Frankfurt and M. I. Strikman, "Hard Nuclear Processes and Microscopic Nuclear Structure," *Phys. Rept.* 160 (1988) 235–427.
- [103] L. L. Frankfurt, M. I. Strikman, D. B. Day, and M. Sargsian, "Evidence for short range correlations from high Q**2 (e, e-prime) reactions," *Phys. Rev. C* 48 (1993) 2451–2461.
- [104] L. L. Frankfurt and M. I. Strikman, "High-Energy Phenomena, Short Range Nuclear Structure and QCD," Phys. Rept. 76 (1981) 215–347.
- [105] M. M. Sargsian et al., "Hadrons in the nuclear medium," J. Phys. G 29 no. 3, (2003) R1-R45, arXiv:nucl-th/0210025.
- [106] J. Rozynek and M. C. Birse, "Nucleon Nucleon Correlation Effects on Deeply Inelastic Lepton Scattering in the Region X > 1," *Phys. Rev. C* 38 (1988) 2201–2204.
- [107] I. Niculescu, J. Arrington, R. Ent, and C. E. Keppel, "Moments of nuclear and nucleon structure functions at low Q**2 and the momentum sum rule," *Phys. Rev. C* 73 (2006) 045206, arXiv:hep-ph/0509241.
- [108] L. A. Kondratyuk and M. Shmatikov, "EMC effect and multi quark bags in nuclei," Yad. Fiz. 41 (1985) 222–235.
- [109] M. M. Sargsian, "Superfast quarks in the nuclear medium," Nucl. Phys. A 782 (2007) 199–206.
- [110] A. J. Freese, M. M. Sargsian, and M. I. Strikman, "Probing superfast quarks in nuclei through dijet production at the LHC," *Eur. Phys. J. C* 75 no. 11, (2015) 534, arXiv:1411.6605 [hep-ph].
- [111] A. J. Freese, W. Cosyn, and M. M. Sargsian, "QCD evolution of superfast quarks," *Phys. Rev. D* 99 no. 11, (2019) 114019, arXiv:1511.06044 [hep-ph].
- [112] A. Bodek and J. L. Ritchie, "Fermi Motion Effects in Deep Inelastic Lepton Scattering from Nuclear Targets," *Phys. Rev. D* 23 (1981) 1070.
- [113] A. Bodek and J. L. Ritchie, "Further Studies of Fermi Motion Effects in Lepton Scattering from Nuclear Targets," *Phys. Rev. D* 24 (1981) 1400.
- [114] CCFR Collaboration, M. Vakili et al., "Nuclear structure functions in the large x large Q**2 kinematic region in neutrino deep inelastic scattering," Phys. Rev. D 61 (2000) 052003, arXiv:hep-ex/9905052.

- [115] S. Rock, "Elastic electron proton scattering at high Q**2," 11, 1986.
- [116] J. Arrington et al., "Inclusive Scattering from Nuclei at x>1 in the quasielastic and deeply inelastic regimes," JLab Proposal PR12-06-105 (2006) 1-21. https://www.jlab.org/exp_prog/proposals/06/PR12-06-105.pdf.
- [117] N. Fomin *et al.*, "Scaling of the F_2 structure function in nuclei and quark distributions at x > 1," *Phys. Rev. Lett.* **105** (2010) 212502, arXiv:1008.2713 [nucl-ex].
- [118] J. Arrington et al., "Inclusive electron scattering from nuclei at x approximately = 1," Phys. Rev. C 53 (1996) 2248-2251, arXiv:nucl-ex/9504003.
- [119] B. W. Filippone et al., "Nuclear structure functions at x > 1," Phys. Rev. C 45 (1992) 1582–1585.
- [120] BCDMS Collaboration, A. C. Benvenuti *et al.*, "Nuclear structure function in carbon near x = 1," Z. Phys. C 63 (1994) 29–36.
- [121] S. A. Kulagin and R. Petti, "Global study of nuclear structure functions," Nucl. Phys. A 765 (2006) 126-187, arXiv:hep-ph/0412425.
- [122] L. T. Brady, A. Accardi, T. J. Hobbs, and W. Melnitchouk, "Next-to leading order analysis of target mass corrections to structure functions and asymmetries," *Phys. Rev. D* 84 (2011) 074008, arXiv:1108.4734 [hep-ph]. [Erratum: Phys.Rev.D 85, 039902 (2012)].
- [123] M. A. G. Aivazis, F. I. Olness, and W.-K. Tung, "QCD formulation of charm production in deep inelastic scattering and the sea quark - gluon dichotomy," *Phys. Rev. Lett.* 65 (1990) 2339–2342.
- [124] B. A. Kniehl, G. Kramer, I. Schienbein, and H. Spiesberger, "Collinear subtractions in hadroproduction of heavy quarks," *Eur. Phys. J. C* 41 (2005) 199–212, arXiv:hep-ph/0502194.
- [125] H1, ZEUS Collaboration, H. Abramowicz et al., "Combination of measurements of inclusive deep inelastic e[±]p scattering cross sections and QCD analysis of HERA data," Eur. Phys. J. C 75 no. 12, (2015) 580, arXiv:1506.06042 [hep-ex].
- [126] F. M. Steffens, M. D. Brown, W. Melnitchouk, and S. Sanches, "Parton distributions in the presence of target mass corrections," *Phys. Rev. C* 86 (2012) 065208, arXiv:1210.4398 [hep-ph].
- [127] T. Becher, M. Neubert, and L. Rothen, "Factorization and N^3LL_p +NNLO predictions for the Higgs cross section with a jet veto," *JHEP* **10** (2013) 125, arXiv:1307.0025 [hep-ph].